


Review on sensitivity to neutrino mass

Nicola Rossi and A. Cocco, N. D'Ambrosio,
M. Messina, F. Pofi and F. Virzi (*LNGS Group*)



PTOLEMY ITA Meeting
Roma, Feb 18th 2024



Part I

What we know about
neutrino mass

Neutrino mass

1. $m_\beta^2 = \sum_i |U_{ei}|^2 m_i^2$ [beta decay]
2. $m_{\beta\beta} = |\sum_i U_{ei}^2 m_i|$ [double beta decay]
3. $\Sigma = \sum_i m_i$ [cosmology]



Depending on

- Mass ordering
- Oscillation parameters
- Cosmological model

Reference formulae

					ing ($\Delta\chi^2 = 2.3$)
					3σ range
					$0.270 \rightarrow 0.341$
					$31.31 \rightarrow 35.74$
without SK atmospheric data	$\sin^2 \theta_{12}$	$0.348^{+0.018}_{-0.023}$	$0.406 \rightarrow 0.620$	$0.578^{+0.016}_{-0.021}$	$0.412 \rightarrow 0.623$
	$\theta_{12}/^\circ$	$33.8^{+1.2}_{-0.7}$	$-40.1^{+1.0}_{-0.6} \rightarrow 51.0$	$40.5^{+0.9}_{-0.5}$	$20.0^{+0.9}_{-0.5} \rightarrow 52.1$
$m_{\nu_e}^{\text{eff}}$	$\sin \theta_1$	$0.2396^{+0.018}_{-0.023}$	$0.406 \rightarrow 0.620$	$0.578^{+0.016}_{-0.021}$	$0.412 \rightarrow 0.623$
	δ_{CP}	$90^{+1.0}_{-0.6}$	$-40.1^{+1.0}_{-0.6} \rightarrow 51.0$	$40.5^{+0.9}_{-0.5}$	$20.0^{+0.9}_{-0.5} \rightarrow 52.1$
m_{ee}	10^{-3} eV^2	$+2.011^{+0.011}_{-0.027}$	$+2.428 \rightarrow +2.597$	$-2.498^{+0.025}_{-0.025}$	$-2.081 \rightarrow -2.408$
		Normal Ordering (best fit)		Inverted Ordering ($\Delta\chi^2 = 6.4$)	
with SK atmospheric data	$\frac{\Delta m_{21}^2}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	$6.82 \rightarrow 8.03$	$7.41^{+0.21}_{-0.20}$	$6.82 \rightarrow 8.03$
	$\frac{\Delta m_{3\ell}^2}{10^{-3} \text{ eV}^2}$	$+2.507^{+0.026}_{-0.027}$	$+2.427 \rightarrow +2.590$	$-2.486^{+0.025}_{-0.028}$	$-2.570 \rightarrow -2.406$

About ordering

N.O.
I.O.



Y.E.S.

Yearningly
Expected
Spectrum

[F. Vissani, Neutrino 2018]

Beta decay ent-point

KATRIN (Tritium)


$m_\beta < 0.45 \text{ meV}$ (90% CL)
(Lokhov-Tkachov)

$m_\beta < 0.31 \text{ meV}$ (90% CL)
(Feldman-Cousins)

$Q_\beta = 18575.0 \pm 0.3 \text{ meV}$



Neutrino-less double beta decay




KamLAND-Zen

$T_{1/2} > 3.8 \times 10^{26} \text{ y}$
 $m_{\beta\beta} < 28\text{-}122 \text{ meV}$

LEGEND-200


$T_{1/2} > 1.9 \times 10^{26} \text{ y}$
 $m_{\beta\beta} < 79\text{-}180 \text{ meV}$

Cosmology





Neutrinos affects the growth of the cosmic clustering → CMB, LSS

- Planck+ACT+BAO+RSD
- Planck+ACT_new
- Planck+BAO_sample5eV_new
- CMB+ACT DR6 lensing+DESI



$m_{\beta\beta}$ Bootstrap



Next generation 0v $\beta\beta$
(KamLAND-Zen, LEGEND-1000, CUPID)
 $m_{\beta\beta} \sim 15$ meV

$(m_0, m_{\beta\beta}) = f(\text{oscillation, Majorana, Cosmology})$

Chance for discovery:



Planck + ACT: 66%

Planck + BAO: 41%

Planck + BAO + ACT + RDS: 36%

Planck + BAO + DESI: 16%

m_β Bootstrap



Most conservative choice (CMB only)

$\langle m_\beta \rangle = 31 \text{ meV (NO)}$

$\langle m_\beta \rangle = 72 \text{ meV (IO)}$

(challenging)

Dirac vs Majorana

Race for Majorana Neutrino

$T_{1/2} > 10^{28}$ y

LEGEND-1000

(1t of ^{76}Ge , 500 M\$, >10 y)

nEXO


(5t of ^{136}Ge , 500 M\$, >10 y)

CUPID

(1t of ^{100}Mo , 100 M\$, >10 y)

Expected rate

- **Dirac** 0.4 events/g/y
- **Majorana** 0.8 events/g/y




Problem to face



1 g
5000 m²

We urgently need
source compactification!!!



100 g
500000 m²


Part II

Experimental sensitivity
to neutrino mass

Reference source

$$N_{dec} = \frac{1}{2} \left(\frac{m_S \mathcal{N}_A}{A_{({}^3H)}} (1 - e^{-t/\tau}) \right) \simeq 10^{16}$$

(50%
Efficiency
for total
events)
in 3 y




$\rho = 0.2 \text{ mg/m}^2$
(full loading)


$\rightarrow 1 \mu\text{g}$

716 MBq
(19.3 mCi)

Origin of sensitivity




Spectrum distortion



Region of interest

Counting analysis




$$N_{\sigma,m} = \int_{E_{th}} S_{\sigma,m}(E) dE$$

$$N_{\sigma,0} = \int_{E_{th}} S_{\sigma,0}(E) dE$$


Sensitivity 90 %CL :=

$$\frac{N_{\sigma,0} - N_{\sigma,m}}{\sqrt{N_{\sigma,0} + N_{\sigma,m}}} = 1.64$$


Sensitivity from counting



Codice: PtSens®




Profile likelihood



$$\mathcal{L}_0 = -2 \log \frac{\mathcal{L}(S_{\sigma,0} | \text{data}_0)}{\mathcal{L}(S_{\sigma,m} | \text{data}_0)}$$


$$\mathcal{L} = -2 \log \frac{\mathcal{L}(S_{\sigma,0} | \text{data}_m)}{\mathcal{L}(S_{\sigma,m} | \text{data}_m)}$$



[Cowan et al. (2013)]








Profile likelihood with spectral fit: 1 µg

Profile Likelihood




Example
 $m = 200 \text{ meV}$,
 $\sigma = 50 \text{ meV}$

L_1/L_0 as a function of m_{eff}




For different resolution
from 50 meV to 1 eV

Resulting sensitivity 1 μg





Counting analysis



Profile Likelihood



Effect of releasing Q_0 - Ex 2/2



Q fixed

$M = 175 \text{ meV}$

$\sigma = 500 \text{ meV}$





Q free

$M = 175 \text{ meV}$

$\sigma = 500 \text{ meV}$



Effect of releasing Q_0 - Ex 1/2



Q fixed

$M = 100 \text{ meV}$

$\sigma = 500 \text{ meV}$




Q free

$M = 100 \text{ meV}$


$\sigma = 500 \text{ meV}$

Sensitivity to (unknown) smearing



**Zero-point
Field (ZPF)
Effect**

Spectrum Deconvolution



Region-1: dominated by ZPF effect
Region-2: mass distortion region

Observed spectrum

$$S(E) = S_m(E) \otimes G_{\text{Res}} \otimes G_{\text{ZPF}}$$

Strategy:


1. Deconvolution in Region-1

$$\tilde{G}_{\text{ZPF}} \simeq S(E) \otimes G_{\text{Res}}^{-1} \otimes S_0^{-1}(E)$$

2. Penalty fit in Region-2

$$S_m(E) = S(E) \otimes \tilde{G}_{\text{ZPF}}^{-1} G_{\text{Res}}^{-1}$$

Deconvolution example



Events in Region-1 ($\sim 10^{15}$) are 1/10 of events needed for Region-2 fit ($\sim 10^{16}$)

Simulated:

$$\sigma_{\text{zpF}} = 500 \text{ meV}$$

meff = 200 meV

Deconvoltuion in Region-1 (-20, -3):


$$\sigma_{\Delta} = 0.49 \pm 0.06 \text{ eV (12\%)}$$

Mass with prior from fit in (-3, 2):

$$m_v = 0.196 \pm 0.026 \text{ eV (13\%)}$$

Example 2/2

Profile Likelihood




Simulated:


$$\sigma_\Delta = 500 \text{ meV}$$

$$m_\nu = 125 \text{ meV}$$


$$\sigma = 200 \text{ meV}$$



Sensitivity to (known) smearing




Quantum levels




Total spectrum

Region of interest




S_m (200 meV)
 S_0 (0 meV)



Random sampling
10 μg sample

Profile likelihood

Profile Likelihood ($N_{\text{sim}}=1000$), $m_{\text{eff}}=0.2$ [eV], target=1. μg




Source = 1 μg

$m = 200$ meV

$\sigma = 50$ meV

Profile likelihood




Source = 10 μg

$m = 200$ meV

$\sigma = 50$ meV

Profile likelihood





Source = 20 μg

$m = 200$ meV



$\sigma = 50$ meV

Profile likelihood



Source = $50 \mu\text{g}$
m = 200 meV
 σ = 50 meV

Profile likelihood





Source = 100 µg

$m = 200 \text{ meV}$

$\sigma = 50 \text{ meV}$

From P. Campana (Bayesian)

- The dependence of the upper limits on m_ν from the number of collected events N has been fitted with a power law.
- In each case, the results are consistent between different choices of CL.



Conclusions

- Ptolemy could be **competitive with KATRIN** in the neutrino mass exploration with a very small target ($1\mu\text{g}$)
- (Tough) **expectations for m_β** are not really promising
- Gaussian unknown or complex know **smearing** could be reasonably handled

Thank you very much!