

COHERENT STATES DETECTION FOR LIGHT DARK MATTER SEARCHES USING MULTI-QUBIT SENSOR

WP8: Application for Quantum Sensing and Machine Learning

QUART&T Kick-Off Meeting


February 21, 2025 | Online

Marco Gobbo

University of Milano-Bicocca

INFN – Milano-Bicocca

Bicocca Quantum Technologies (BiQuTe) Centre




DARK MATTER

“There are a lot of things we understand about the universe, but the fun is in all the things we do not.”

The evidence of existence of dark matter is based on astronomical observations of its gravitational interactions.

- Velocity discrepancy in galaxy clusters (1933 - Zwicky)
- Galaxy rotation curves (1970s - Rubin and Ford)
- Gravitational lensing (1980s - present)
- Large-scale structure of the universe
- Cosmic microwave background (CMB)
- Baryon acoustic oscillations (BAO)
- Collision of galaxy clusters
- ... and so on



Credit: xkcd

QUANTUM SENSING – LIGHT DARK MATTER SEARCHES

AXION


Strong CP problem:

- QCD does not conserve CP (theory);
- QCD conserves CP (experiment).


$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \theta \frac{g^2}{32\pi^2}F^{\mu\nu}\tilde{F}_{\mu\nu} + \bar{\psi}(i\gamma^\mu D_\mu - me^{i\theta'\gamma_5})\psi$$

A possible solution to conserve CP: **axion** (pseudo-Goldstone boson):

$$\mathcal{L} \supset -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \frac{1}{2}\partial_\mu a\partial^\mu a - \frac{1}{2}m_a^2 a^2 - \frac{1}{8}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu}$$



Inverse Primakoff effect



[10.5281/zenodo.3932430](https://zenodo.org/record/3932430)



QUANTUM SENSING – LIGHT DARK MATTER SEARCHES

DARK PHOTON

Some anomalies in astrophysics could be explained through the interactions between the **dark matter** and **dark photons**.

- Introducing a new gauge $U(1)$ symmetry in the SM;
- **Kinetic mixing** with the electromagnetic field.

$$\mathcal{L} \supset -\frac{1}{4}F'^{\mu\nu}F'_{\mu\nu} + \frac{1}{2}m_{A'}A'^{\mu}A'_{\mu} + \epsilon e A'^{\mu}J_{\mu}^{EM}$$



[10.5281/zenodo.3932430](https://zenodo.org/record/3932430)

DM DETECTION USING QUBITS: DIRECT EXCITATION

Assumption: the DM candidate generates a weak coherent effective electromagnetic field.

If the field is resonant with the qubit $|g\rangle \rightarrow |e\rangle$ transition it can trigger the qubit to be in an excited state.

Qubit state $|\psi(t)\rangle = \psi_g(t)|g\rangle + e^{-i\omega_q t}\psi_e(t)|e\rangle$

Hamiltonian for an axion-induced electric field $H = \omega_q|e\rangle\langle e| + 2\eta \cos(\mathbf{m}_a t - \alpha) (|e\rangle\langle g| + |g\rangle\langle e|)$

$$p_{eg}(\tau) \simeq 0.11 \left(\frac{g_{a\gamma\gamma}}{10^{-10} \text{ GeV}^{-1}} \right)^2 \left(\frac{m_a}{1 \text{ } \mu\text{eV}} \right)^2 \left(\frac{B_0}{1 \text{ T}} \right)^2 \left(\frac{\tau}{100 \text{ } \mu\text{s}} \right)^2 \kappa^2 \left(\frac{C}{0.1 \text{ pF}} \right) \left(\frac{d}{100 \text{ } \mu\text{m}} \right)^2 \left(\frac{\rho_{\text{DM}}}{0.45 \text{ GeV/cm}^3} \right)$$

Hamiltonian for a dark photon-induced electric field $H = \omega_q|e\rangle\langle e| + 2\eta \sin \mathbf{m}_{A'} t (|e\rangle\langle g| + |g\rangle\langle e|)$

$$p_{eg}(\tau) \simeq 0.12 \kappa^2 \cos \Theta \left(\frac{\epsilon}{10^{-11}} \right)^2 \left(\frac{f}{1 \text{ GHz}} \right) \left(\frac{\tau}{100 \text{ } \mu\text{s}} \right)^2 \left(\frac{C}{0.1 \text{ pF}} \right) \left(\frac{d}{100 \text{ } \mu\text{m}} \right)^2 \left(\frac{\rho_{\text{DM}}}{0.45 \text{ GeV/cm}^3} \right)$$



[10.1103/PhysRevD.110.115021](https://arxiv.org/abs/10.1103/PhysRevD.110.115021), [10.1103/PhysRevLett.131.211001](https://arxiv.org/abs/10.1103/PhysRevLett.131.211001)

DM DETECTION USING QUBITS: QND MEASUREMENTS

Qubit state dependent on cavity population



$$H = \omega_c a^\dagger a + \frac{1}{2}(\omega_q + 2\chi a^\dagger a)\sigma_z$$

1



[10.1103/PhysRevLett.126.141302](https://doi.org/10.1103/PhysRevLett.126.141302)

2



[10.1038/s41567-018-0066-3](https://doi.org/10.1038/s41567-018-0066-3)

EXTENDING THE QND DETECTION SCHEME



Increasing the number of qubits (N) in a photon counter device provides several advantages:

1. Reduction of readout errors

$$p_{\text{err}}(\otimes_i^N |e\rangle_i) = \prod_{i=1}^N p_{\text{err}}(|e\rangle) \quad p_{\text{err}}^{\text{best}}(|e\rangle) = 0.3\%$$

2. Improved detection efficiency;
3. More effective quantum non-demolition (QND) measurement;
4. Reduced scan time in experimental searches.

$$\kappa_{\text{ext}} \ll 2 \sum_{i=1}^N \chi_i \quad \longrightarrow \quad \text{Total phase shift } \phi = 2\pi$$




[10.3390/app14041478](https://arxiv.org/abs/10.3390/app14041478)

SIMULATIONS OF MULTI-QUBIT CHIP DEVICE

Using an input-output theory we can model:


- The interaction between an input field and the cavity
- The interaction between the cavity and the qubits



From the model, it is possible to evaluate the conditional probability


$$P(|q\rangle | |1\rangle) = \frac{P(|q\rangle)P(|1\rangle | |q\rangle)}{P(|1\rangle)}$$
$$P(|n\rangle) = \frac{|\alpha|^{2n}}{n!} e^{-|\alpha|^2}$$

SIMULATIONS OF MULTI-QUBIT CHIP DEVICE – 2 QUBITS




Total phase shift $\phi = 2\pi$

3D double-qubit single-photon detector




[10.1016/j.nima.2024.170010](https://doi.org/10.1016/j.nima.2024.170010)

SIMULATIONS OF MULTI-QUBIT CHIP DEVICE – 2 QUBITS




Total phase shift $\phi = 2\pi$




Total phase shift $\phi = \pi$

SIMULATIONS OF MULTI-QUBIT CHIP DEVICE – 4 QUBITS



Total phase shift $\phi = 2\pi$



Total phase shift $\phi = \pi$

PHASE DISTRIBUTION

Simplest case: single qubit in the absence of excitation, relaxation, and decoherence phenomena.

$$|g\rangle \xrightarrow{\gamma/2} \frac{|g\rangle + |e\rangle}{\sqrt{2}} \xrightarrow{\gamma} \frac{|g\rangle + e^{i\phi}|e\rangle}{\sqrt{2}} \xrightarrow{-\gamma/2} \frac{1}{2}[(e^{i\phi} + 1)|g\rangle + (e^{i\phi} - 1)|e\rangle]$$

$$P(|g\rangle) = \frac{1 + \cos \phi}{2}$$


$$P(|e\rangle) = \frac{1 - \cos \phi}{2}$$

Extending to a system of independent N qubits without entanglement, in which the total phase is equally distributed $\phi_i = \phi/N$


$$P(k) = \binom{N}{k} P(|g\rangle)^k P(|e\rangle)^{N-k}$$

k is the number of qubits in the $|g\rangle$ in a system of N qubits.


LOW-ENTANGLEMENT (?)



$\phi = 2\pi$ ← Total phase shift → $\phi = \pi$



CONCEPTUAL PLANAR CHIP DESIGN



PRELIMINARY CONCLUSIONS

- The $N = 2$ qubit model works correctly and does not introduce any entanglement effects;
- For $N \geq 3$ qubits, some low entanglement begins to emerge which:
 - Requires a more refined theoretical model for a complete description of the system's dynamics;
 - Suggests considering the use of entangled initial states, such as GHZ states, with a proper detection protocol;

$$|\text{GHZ}\rangle = \frac{|g\rangle^{\otimes N} + |e\rangle^{\otimes N}}{\sqrt{2}}$$

- Further analysis is needed for cases with different χ values, particularly when dealing with non-uniform phase distributions.

NEXT-STEPS

- In progress** Refine the theoretical model through the input-output theory;
- Done** Design of 3D version with 2 qubits;
- Planning** Design of 2D version with 2 qubits;
- Not started** Fabrication of the devices;
- Not started** Characterization of the devices;
- Not started** Axion and dark photon measurement campaigns.

COLLABORATION

UNIMIB/INFN-MIB group

Matteo Borghesi, Pietro Campana, Rodolfo Carobene, Alessandro Cattaneo, Hervè Corti, Marco Faverzani, Elena Ferri, Sara Gamba, Andrea Giachero, **Marco Gobbo**, Danilo Labranca, Roberto Moretti, Angelo Nucciotti, Luca Origo

INFN-LNF group

Fabio Chiarello (CNR), Alessandro D'Elia, Claudio Gatti, Alex Stephane Piedjou Komnang, Davide Milillo, Alessio Rettaroli, Simone Tocci

UNIFI/INFN-FI group

Leonardo Banchi, Alessandro Cuccoli

