




Marco Radici



Pedagogical overview of SIDIS and TMD phenomenology





Useful references



• Lecture notes

- V. Barone - Cabelo School https://www.fe.infn.it/cabelo_school/2010/cabelo_school_2010.pdf
- A. Bacchetta - Trento School https://www2.pv.infn.it/~bacchett/teaching/Bacchetta_Trento2012.pdf
- R. Jaffe - Erice School <https://arxiv.org/pdf/hep-ph/9602236.pdf>
- P. Mulders - GGI School <http://www.nat.vu.nl/~mulders/tmdreview-vs3.pdf>

• Books

- V. Barone, P. Ratcliffe - *Transverse Spin Physics*
- J. Collins - *Foundations of perturbative QCD*
- R. Devenish, A. Cooper-Sarkar - *Deep Inelastic Scattering*
- T. Muta - *Foundations of Quantum Chromodynamics*



• Papers

- EPJ-A topical issue: The 3D structure of the nucleon
https://link.springer.com/journal/10050/topicalCollection/AC_628286e999d9a60c9a780398df15f93d
- M. Diehl - *Introduction to GPDs and TMDs* <https://inspirehep.net/literature/1408303>
- A. Metz, A. Vossen - *Parton fragmentation functions* <https://inspirehep.net/literature/1475000>



- **Why SIDIS?**




SIDIS: I) access fragmentation







SIDIS: I) access fragmentation



inclusive DIS: no sensitivity to fragmentation



SIDIS




e^+e^- annihilation

check universality of FFs

(simpler picture than hadronic collisions)




SIDIS: 2) access intrinsic partonic \perp motion




inclusive DIS: - hard scale $Q^2 = -q^2 \gg M^2$ to “see” partons
- no further scale to probe proton interior



SIDIS: 2) access intrinsic partonic \perp motion




inclusive DIS: - hard scale $Q^2 = -q^2 \gg M^2$ to “see” partons
 - no further scale to probe proton interior



semi-inclusive DIS (SIDIS):

- hard scale $Q^2 = -q^2 \gg M^2$ to “see” partons
 - soft scale: detect hadron h with $P_{hT}^2 \sim M^2 \ll Q^2$




with these two scales, the process is factorizable into a hard photon-quark vertex and a quark \rightarrow hadron fragmentation

$$\mathbf{P}_{hT} = z\mathbf{k}_\perp + \mathbf{P}_\perp + \mathcal{O}(k_\perp^2/Q^2) \quad z = \text{fractional energy of } h \text{ (analogous of } x)$$


hadron P_{hT} arises from struck quark k_\perp and transverse momentum P_\perp generated during fragmentation



SIDIS: 2) access intrinsic partonic \perp motion




inclusive DIS: - hard scale $Q^2 = -q^2 \gg M^2$ to “see” partons
 - no further scale to probe proton interior



semi-inclusive DIS (SIDIS):

- hard scale $Q^2 = -q^2 \gg M^2$ to “see” partons
 - soft scale: detect hadron h with $P_{hT}^2 \sim M^2 \ll Q^2$



with these two scales, the process is factorizable into a hard photon-quark vertex and a quark \rightarrow hadron fragmentation


$$\mathbf{P}_{hT} = z\mathbf{k}_{\perp} + \mathbf{P}_{\perp} + \mathcal{O}(k_{\perp}^2/Q^2) \quad z = \text{fractional energy of } h \text{ (analogous of } x)$$

hadron P_{hT} arises from struck quark k_{\perp} and transverse momentum P_{\perp} generated during fragmentation

measure $P_{hT} \rightarrow$ get to k_{\perp}



SIDIS: 3) access chiral-odd structures



chirality = helicity for a spin-1/2 object
chiral-odd structures mix quark helicities:



$$\langle + | \dots | - \rangle, \langle - | \dots | + \rangle$$

hence, chiral-odd structures can appear
only paired to another chiral-odd structure
because cross section is chiral even

chiral-odd structures suppressed in DIS



SIDIS: 3) access chiral-odd structures




DIS

chirality = helicity for a spin-1/2 object
chiral-odd structures mix quark helicities:

$$\langle + | \dots | - \rangle, \langle - | \dots | + \rangle$$

hence, chiral-odd structures can appear only paired to another chiral-odd structure because cross section is chiral even

chiral-odd structures suppressed in DIS



SIDIS

chiral-odd structures possible by pairing with a chiral-odd fragmentation function



- A short recap of inclusive DIS and collinear factorization



“Deep-Inelastic” kinematics



Internal hadron structure is best explored with a powerful “microscopic lens”
need a process with a hard scale; example: inclusive lepton-proton scattering

$$\ell + N(P) \rightarrow \ell' + X$$

Kinematic invariants

$$P^2 = M^2$$


$$Q^2 = -q^2 \approx 2EE'(1 - \cos\theta_e) = 4EE'\sin^2\theta_e/2$$

$$\nu = \frac{P \cdot q}{M} \stackrel{\text{TRF}}{=} E - E' \quad \text{transferred energy}$$

$$y = \frac{P \cdot q}{P \cdot \ell} \stackrel{\text{TRF}}{=} \frac{E - E'}{E} \quad \text{fraction of } \nu \quad 0 \leq y \leq 1$$

$$x_b = \frac{Q^2}{2P \cdot q} \stackrel{\text{TRF}}{=} \frac{Q^2}{2M\nu} \quad \text{inelastic } 0 < x \leq 1 \text{ elastic}$$

$$W^2 = (P + q)^2 = M^2 + Q^2(1/x - 1) \geq M^2 \quad \text{invariant mass}$$



$$P = (M, 0, 0, 0) \quad \text{initially at rest}$$

$$q = (\nu, 0, 0, |\mathbf{q}|)$$

$$P' = (\sqrt{M^2 + P_z'^2}, 0, 0, P_z')$$



“Deep-Inelastic” kinematics




Internal hadron structure is best explored with a powerful “microscopic lens” need a process with a hard scale; example: inclusive lepton-proton scattering

$$\ell + N(P) \rightarrow \ell' + X$$

Kinematic invariants

$$\begin{aligned}
 P^2 &= M^2 \\
 Q^2 &= -q^2 \approx 2EE'(1 - \cos\theta_e) = 4EE'\sin^2\theta_e/2 \\
 \nu &= \frac{P \cdot q}{M} \stackrel{\text{TRF}}{=} E - E' \quad \text{transferred energy} \\
 y &= \frac{P \cdot q}{P \cdot \ell} \stackrel{\text{TRF}}{=} \frac{E - E'}{E} \quad \text{fraction of " " } 0 \leq y \leq 1 \\
 x_B &= \frac{Q^2}{2P \cdot q} \stackrel{\text{TRF}}{=} \frac{Q^2}{2M\nu} \quad \text{inelastic } 0 < x \leq 1 \text{ elastic} \\
 W^2 &= (P + q)^2 = M^2 + Q^2(1/x - 1) \geq M^2 \quad \text{invariant mass}
 \end{aligned}$$




$$\begin{aligned}
 P &= (M, 0, 0, 0) \quad \text{initially at rest} \\
 q &= (\nu, 0, 0, |\mathbf{q}|) \\
 P' &= (\sqrt{M^2 + P_z'^2}, 0, 0, P_z') \\
 P_z' &= |\mathbf{q}| \gg M
 \end{aligned}$$

Deep-Inelastic regime: $\begin{cases} Q^2 \rightarrow \infty \\ x_B = \frac{Q^2}{2P \cdot q} \text{ finite} \end{cases}$

Light-Cone coordinates:


$$\begin{aligned}
 a^\mu &= (a_0, a_1, a_2, a_3) = (a_+, a_-, \mathbf{a}_\perp) \\
 a_\pm &= \frac{a_0 \pm a_3}{\sqrt{2}} \quad \mathbf{a}_\perp = (a_1, a_2)
 \end{aligned}$$



$$\begin{aligned}
 P' &\approx (|\mathbf{q}|, 0, 0, |\mathbf{q}|) \\
 P' &\approx (P'_+ = \sqrt{2}|\mathbf{q}|, P'_- \approx 0, \mathbf{0}_\perp) \\
 \text{in general } P'_+ &\gg P'_-
 \end{aligned}$$



“Deep-Inelastic” kinematics




Internal hadron structure is best explored with a powerful “microscopic lens” need a process with a hard scale; example: inclusive lepton-proton scattering

$$\ell + N(P) \rightarrow \ell' + X$$

Kinematic invariants

$$\begin{aligned}
 P^2 &= M^2 \\
 Q^2 &= -q^2 \approx 2EE'(1 - \cos\theta_e) = 4EE'\sin^2\theta_e/2 \\
 \nu &= \frac{P \cdot q}{M} \stackrel{\text{TRF}}{=} E - E' \quad \text{transferred energy} \\
 y &= \frac{P \cdot q}{P \cdot \ell} \stackrel{\text{TRF}}{=} \frac{E - E'}{E} \quad \text{fraction of } \nu \quad 0 \leq y \leq 1 \\
 x_B &= \frac{Q^2}{2P \cdot q} \stackrel{\text{TRF}}{=} \frac{Q^2}{2M\nu} \quad \text{inelastic } 0 < x \leq 1 \text{ elastic} \\
 W^2 &= (P + q)^2 = M^2 + Q^2(1/x - 1) \geq M^2 \quad \text{invariant mass}
 \end{aligned}$$



$$\begin{aligned}
 P &= (M, 0, 0, 0) \quad \text{initially at rest} \\
 q &= (\nu, 0, 0, |\mathbf{q}|) \\
 P' &= (\sqrt{M^2 + P_z'^2}, 0, 0, P_z') \\
 P_z' &= |\mathbf{q}| \gg M
 \end{aligned}$$


Deep-Inelastic regime:

$$\begin{cases} Q^2 \rightarrow \infty \\ x_B = \frac{Q^2}{2P \cdot q} \text{ finite} \end{cases}$$

$$P' \approx (|\mathbf{q}|, 0, 0, |\mathbf{q}|)$$

Light-Cone coordinates:


$$\begin{aligned}
 a^\mu &= (a_0, a_1, a_2, a_3) = (a_+, a_-, \mathbf{a}_\perp) \\
 a_\pm &= \frac{a_0 \pm a_3}{\sqrt{2}} \quad \mathbf{a}_\perp = (a_1, a_2)
 \end{aligned}$$




$$P' \approx (P'_+ = \sqrt{2}|\mathbf{q}|, P'_- \approx 0, \mathbf{0}_\perp)$$

in general $P'_+ \gg P'_-$

In this regime, only one single dominant component of proton momentum, P_+




inclusive Deep-Inelastic Scattering (DIS): **1 dominant direction of momenta**
→ **all partons collinear to proton**




Basics of Feynman parton model:

- DIS regime and relativistic corrections:
the virtual photon probes a frozen ensemble of partons
- **factorisation** between hard collision and proton structure





inclusive Deep-Inelastic Scattering (DIS): **1 dominant direction of momenta**
→ **all partons collinear to proton**




Basics of Feynman parton model:

- DIS regime and relativistic corrections: the virtual photon probes a frozen ensemble of partons
- **factorisation** between hard collision and proton structure






inclusive Deep-Inelastic Scattering (DIS): **1 dominant direction of momenta**
→ all partons collinear to proton



Basics of Feynman parton model:

- DIS regime and relativistic corrections: the virtual photon probes a frozen ensemble of partons
- **factorisation** between hard collision and proton structure
- **1D imaging of proton structure**, parametrised by collinear Parton Distribution Functions **PDF(x, Q²)**







inclusive DIS



More rigorously:

one photon-exchange approximation






inclusive DIS





More rigorously:

one photon-exchange approximation



optical theorem



cut-diagram notation:

cross section = product of two amplitudes
particles entering cut are on-shell




inclusive DIS





More rigorously:

one photon-exchange approximation



optical theorem



cut-diagram notation:

cross section = product of two amplitudes
particles entering cut are on-shell

$$\frac{d\sigma}{dx_B dy d\phi_S} = \frac{\alpha^2 y}{2Q^4} L_{\mu\nu}(\ell, \ell', \lambda_e) W^{\mu\nu}(q, P, S)$$

leptonic tensor

hadronic tensor

calculable in QED

linear combination of all tensor structures with q, P, S , subject to Hermiticity, gauge-, parity- and time reversal- invariance
→ parametrised with **four** structure functions



Collinear factorization theorem



$$\frac{d\sigma}{dx_B dy d\phi_S} = \frac{2\alpha^2}{x_{BY} Q^2} \left\{ A(y) F_{UU,T} + B(y) F_{UU,L} + \lambda_e S_L C(y) F_{LL} + \lambda_e |\mathbf{S}_T| D(y) \cos \phi_S F_{LT} \right\}$$

each F. (x, Q²)
F_{XY,Z}
ℓ P Y*



Collinear factorization theorem



$$\frac{d\sigma}{dx_B dy d\phi_S} = \frac{2\alpha^2}{x_B y Q^2} \left\{ A(y)F_{UU,T} + B(y)F_{UU,L} + \lambda_e S_L C(y)F_{LL} + \lambda_e |\mathbf{S}_T| D(y) \cos \phi_S F_{LT} \right\} \text{ each } F. (x, Q^2)$$

connection to standard notation

$$F_{UU,T} = 2x_B F_1$$

$$F_{UU,L} = F_2 - 2x_B F_1 + \mathcal{O}(\gamma^2) F_2$$

$$F_{LL} = 2x_B g_1 + \mathcal{O}(\gamma^2) g_2$$

$$F_{LT} = \mathcal{O}(\gamma)(g_1 + g_2)$$

$$\gamma = \frac{2xM}{Q} \text{ target mass correction}$$



Collinear factorization theorem



$$\frac{d\sigma}{dx_B dy d\phi_S} = \frac{2\alpha^2}{x_B y Q^2} \left\{ A(y)F_{UU,T} + B(y)F_{UU,L} + \lambda_e S_L C(y)F_{LL} + \lambda_e |S_T| D(y) \cos \phi_S F_{LT} \right\} \text{ each F. (x, Q}^2\text{)}$$

connection to standard notation

$$F_{UU,T} = 2x_B F_1$$

$$F_{UU,L} = F_2 - 2x_B F_1 + \mathcal{O}(\gamma^2)F_2$$

$$F_{LL} = 2x_B g_1 + \mathcal{O}(\gamma^2)g_2$$

$$F_{LT} = \mathcal{O}(\gamma)(g_1 + g_2)$$

$$\gamma = \frac{2xM}{Q} \text{ target mass correction}$$

unpolarized: $\lambda_e = S_L = 0$

$$\frac{d\sigma}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} [A(y) F_2(x, Q^2) - y^2 F_L(x, Q^2)] \quad Q^2 = sxy$$

$$F_i(x, Q^2) = \sum_f e_f^2 \int_x^1 \frac{d\xi}{\xi} d\hat{\sigma}_{i,f} \left(\alpha_s, \frac{x}{\xi}, \frac{Q^2}{\mu_F^2} \right) \phi_f(\alpha_s, \xi, \mu_F) \equiv \sum_f e_f^2 d\hat{\sigma}_{i,f} \otimes \phi_f$$

usually $\mu_R^2 = \mu_F^2 = Q^2$

$$d\hat{\sigma}_{i,f} = d\hat{\sigma}_{i,f}^{(0)} + \frac{\alpha_s}{4\pi} d\hat{\sigma}_{i,f}^{(1)} + \dots$$



Collinear factorization theorem



$$\frac{d\sigma}{dx_B dy d\phi_S} = \frac{2\alpha^2}{x_B y Q^2} \left\{ A(y)F_{UU,T} + B(y)F_{UU,L} + \lambda_e S_L C(y)F_{LL} + \lambda_e |S_T| D(y) \cos \phi_S F_{LT} \right\} \text{ each } F_{i,j}(\mathbf{x}, Q^2)$$

connection to standard notation

$$F_{UU,T} = 2x_B F_1$$

$$F_{UU,L} = F_2 - 2x_B F_1 + \mathcal{O}(\gamma^2)F_2$$

$$F_{LL} = 2x_B g_1 + \mathcal{O}(\gamma^2)g_2$$

$$F_{LT} = \mathcal{O}(\gamma)(g_1 + g_2)$$

$$\gamma = \frac{2xM}{Q} \text{ target mass correction}$$

unpolarized: $\lambda_e = S_L = 0$

$$\frac{d\sigma}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} [A(y)F_2(x, Q^2) - y^2 F_L(x, Q^2)] \quad Q^2 = sxy$$

$$F_i(x, Q^2) = \sum_f e_f^2 \int_x^1 \frac{d\xi}{\xi} d\hat{\sigma}_{i,f} \left(\alpha_s, \frac{x}{\xi}, \frac{Q^2}{\mu_F^2} \right) \phi_f(\alpha_s, \xi, \mu_F) \equiv \sum_f e_f^2 d\hat{\sigma}_{i,f} \otimes \phi_f$$

usually $\mu_R^2 = \mu_F^2 = Q^2$

$$d\hat{\sigma}_{i,f} = d\hat{\sigma}_{i,f}^{(0)} + \frac{\alpha_s}{4\pi} d\hat{\sigma}_{i,f}^{(1)} + \dots$$

QCD collinear factorization theorem at scale μ_F , valid at all orders

Physics does not depend on fictitious scale μ_F : DGLAP evolution equations




Operator Product Expansion



Factorization from another point of view: the OPE

1. Justification



consider the inclusive DIS
scattering amplitude


$$\mathcal{M} = \bar{u}(k') \gamma_\mu u(k) \frac{e^2}{Q^2} \langle P_X | J^\mu(0) | P \rangle$$

phase space

$$dR = (2\pi)^4 \delta(P + q - P_X) d^4 P_X d^4 k'$$


cross section

$$d\sigma \propto \int_X |\mathcal{M}|^2 dR = \frac{e^4}{Q^4} L_{\mu\nu} W^{\mu\nu} d^4 k'$$



2

optical
theorem



$$L_{\mu\nu} = 2k_\mu k'_\nu + 2k'_\nu k_\mu - Q^2 g_{\mu\nu} \quad \text{leptonic tensor}$$

$$W^{\mu\nu} = \int d^4 P_X (2\pi)^4 \delta(P_X - P - q) \times \langle P | \hat{J}^\mu(0) | P_X \rangle \langle P_X | J^\nu(0) | P \rangle \quad \text{hadronic tensor}$$



Operator Product Expansion



Factorization from another point of view: the OPE

1. Justification

consider the inclusive DIS scattering amplitude



$$\mathcal{M} = \bar{u}(k') \gamma_\mu u(k) \frac{e^2}{Q^2} \langle P_X | J^\mu(0) | P \rangle$$

phase space

$$dR = (2\pi)^4 \delta(P + q - P_X) d^4P_X d^4k'$$


cross section

$$d\sigma \propto \int_X |\mathcal{M}|^2 dR = \frac{e^4}{Q^4} L_{\mu\nu} W^{\mu\nu} d^4k'$$



2

optical theorem



$$L_{\mu\nu} = 2k_\mu k'_\nu + 2k'_\nu k_\mu - Q^2 g_{\mu\nu} \quad \text{leptonic tensor}$$

$$W^{\mu\nu} = \int d^4P_X (2\pi)^4 \delta(P_X - P - q) \times \langle P | \hat{J}^\mu(0) | P_X \rangle \langle P_X | \hat{J}^\nu(0) | P \rangle \quad \text{hadronic tensor}$$

matrix element of bilocal operator

$$= \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle$$

$$\hat{J}^\mu = \bar{\psi} \gamma^\mu \psi$$

parton e.m. current

$$\text{check } \hat{J}^{\mu\dagger} = \hat{J}^\mu$$




Operator Product Expansion



Factorization from another point of view: the OPE

1. Justification



consider the inclusive DIS
scattering amplitude


$$\mathcal{M} = \bar{u}(k') \gamma_\mu u(k) \frac{e^2}{Q^2} \langle P_X | J^\mu(0) | P \rangle$$

phase space

$$dR = (2\pi)^4 \delta(P + q - P_X) d^4 P_X d^4 k'$$


cross section

$$d\sigma \propto \int_X |\mathcal{M}|^2 dR = \frac{e^4}{Q^4} L_{\mu\nu} W^{\mu\nu} d^4 k'$$



2

optical
theorem



$$L_{\mu\nu} = 2k_\mu k'_\nu + 2k'_\nu k_\mu - Q^2 g_{\mu\nu} \quad \text{leptonic tensor}$$

$$W^{\mu\nu} = \int d^4 P_X (2\pi)^4 \delta(P_X - P - q) \times \langle P | \hat{J}^\mu(0) | P_X \rangle \langle P_X | \hat{J}^\nu(0) | P \rangle \quad \text{hadronic tensor}$$

matrix element of bilocal operator

$$= \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle$$

dominated by time-like short distances $\xi^2 \rightarrow 0$, but ill defined !

$$\hat{J}^\mu = \bar{\psi} \gamma^\mu \psi$$

parton e.m. current
check $\hat{J}^{\mu\dagger} = \hat{J}^\mu$





Operator Product Expansion



DIS regime:

$$Q^2 \rightarrow \infty$$

$$x = \frac{Q^2}{2P \cdot q} \Big|_{\text{TRF}} = \frac{Q^2}{2M\nu} \text{ fixed}$$

Target Rest Frame $\Rightarrow \nu \rightarrow \infty$

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle$$



Operator Product Expansion



DIS regime:

$$Q^2 \rightarrow \infty$$

$$x = \frac{Q^2}{2P \cdot q} \Big|_{\text{TRF}} = \frac{Q^2}{2M\nu} \text{ fixed}$$

Target Rest Frame $\Rightarrow \nu \rightarrow \infty$

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle$$

Riemann - Lebesgue theorem:

for $|q \cdot \xi| \rightarrow \infty$, large oscillations and cancelations;

integral is dominated by terms with $|q \cdot \xi| \leq K$ constant



Operator Product Expansion



DIS regime:

$$Q^2 \rightarrow \infty$$

$$x = \frac{Q^2}{2P \cdot q} \Big|_{\text{TRF}} = \frac{Q^2}{2M\nu} \text{ fixed}$$

Target Rest Frame $\Rightarrow \nu \rightarrow \infty$

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle$$

Riemann - Lebesgue theorem:

for $|q \cdot \xi| \rightarrow \infty$, large oscillations and cancelations;

integral is dominated by terms with $|q \cdot \xi| \leq K$ constant

Then, $(q \cdot \xi)|_{\text{cm}} = \nu \xi^0 \leq K \Rightarrow \xi^0 \leq \frac{K}{\nu} \xrightarrow{\nu \rightarrow \infty} 0$

space-like distances $\xi^2 < 0$ are forbidden by causality;

for time-like distances $\xi^2 \geq 0$, $\xi^2 = (\xi^0)^2 - \vec{\xi}^2 \geq 0 \Rightarrow (\xi^0)^2 \geq \vec{\xi}^2 \xrightarrow{\nu \rightarrow \infty} 0$

The integral is dominated by short time-like distances $\xi^2 \rightarrow 0$, but in this limit the bilocal operator is ill defined. Example: free neutron scalar field $\phi(x)$ with propagator $\Delta(x-y)$



Operator Product Expansion



DIS regime:

$$Q^2 \rightarrow \infty$$

$$x = \frac{Q^2}{2P \cdot q} \Big|_{\text{TRF}} = \frac{Q^2}{2M\nu} \text{ fixed}$$

Target Rest Frame $\Rightarrow \nu \rightarrow \infty$

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle$$

Riemann - Lebesgue theorem:

for $|q \cdot \xi| \rightarrow \infty$, large oscillations and cancelations;

integral is dominated by terms with $|q \cdot \xi| \leq K$ constant

Then, $(q \cdot \xi)|_{\text{cm}} = \nu \xi^0 \leq K \Rightarrow \xi^0 \leq \frac{K}{\nu} \xrightarrow{\nu \rightarrow \infty} 0$

space-like distances $\xi^2 < 0$ are forbidden by causality;

for time-like distances $\xi^2 \geq 0$, $\xi^2 = (\xi^0)^2 - \vec{\xi}^2 \geq 0 \Rightarrow (\xi^0)^2 \geq \vec{\xi}^2 \xrightarrow{\nu \rightarrow \infty} 0$

The integral is dominated by short time-like distances $\xi^2 \rightarrow 0$, but in this limit the bilocal operator is ill defined. Example: free neutron scalar field $\phi(x)$ with propagator $\Delta(x-y)$

$$\langle 0 | \mathcal{T}[\phi(x)\phi(y)] | 0 \rangle = -i \Delta(x-y) = i \int \frac{d^4p}{(2\pi)^4} \frac{e^{-ip \cdot (x-y)}}{p^2 - m^2 + i\epsilon} \quad \text{for } x \rightarrow y, \text{ the integral is divergent :}$$

$$= \frac{m}{4\pi^2} \frac{K_1 \left(m \sqrt{-(x-y)^2 + i\epsilon} \right)}{\sqrt{-(x-y)^2 + i\epsilon}} - \frac{i}{4\pi} \delta((x-y)^2) \xrightarrow{x \rightarrow y} \infty$$

K_1 modified Bessel
funct. of 2^o kind



2. Definition


$$\hat{A}(x) \hat{B}(y) = \sum_{i=0}^{\infty} C_i(x-y) \hat{O}_i\left(\frac{x+y}{2}\right)$$

local operators,
regular for $x \rightarrow y$,
typically $\hat{O}_0 = \mathbf{I}$

Wilson coefficients, singular for $x \rightarrow y$,
ordered in decreasing singularity



Operator Product Expansion



2. Definition

$$\hat{A}(x) \hat{B}(y) = \sum_{i=0}^{\infty} C_i(x-y) \hat{O}_i\left(\frac{x+y}{2}\right)$$

local operators,
regular for $x \rightarrow y$,
typically $\hat{O}_0 = \mathbf{I}$

Wilson coefficients, singular for $x \rightarrow y$,
ordered in decreasing singularity

Example: the Wick theorem

$$\begin{aligned} \lim_{x \rightarrow y} \mathcal{T} [\phi(x) \phi(y)] &= : \phi(x) \phi(y) : + \langle 0 | \mathcal{T} [\phi(x) \phi(y)] | 0 \rangle \\ &= 1 \cdot \hat{O}_1 + C_0(x-y) \mathbf{I} \end{aligned}$$



2. Definition

$$\hat{A}(x) \hat{B}(y) = \sum_{i=0}^{\infty} C_i(x-y) \hat{O}_i\left(\frac{x+y}{2}\right)$$

local operators, regular for $x \rightarrow y$, typically $\hat{O}_0 = \mathbf{I}$

Wilson coefficients, singular for $x \rightarrow y$, ordered in decreasing singularity

Example: the Wick theorem

$$\lim_{x \rightarrow y} \mathcal{T} [\phi(x) \phi(y)] = : \phi(x) \phi(y) : + \langle 0 | \mathcal{T} [\phi(x) \phi(y)] | 0 \rangle$$

$$= 1 \cdot \hat{O}_1 + C_0(x-y) \mathbf{I}$$

3. Application to inclusive DIS

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle = \sum_{\{\alpha\}} C_{\{\alpha\}}^{\mu\nu} \left(\frac{M}{Q}\right)^{t-2}$$

twist $t =$ canonical dimension - spin of operator \hat{O}_i



Operator Product Expansion



2. Definition

$$\hat{A}(x) \hat{B}(y) = \sum_{i=0}^{\infty} C_i(x-y) \hat{O}_i\left(\frac{x+y}{2}\right)$$

local operators, regular for $x \rightarrow y$, typically $\hat{O}_0 = \mathbf{I}$

Wilson coefficients, singular for $x \rightarrow y$, ordered in decreasing singularity

Example: the Wick theorem

$$\lim_{x \rightarrow y} \mathcal{T} [\phi(x) \phi(y)] = : \phi(x) \phi(y) : + \langle 0 | \mathcal{T} [\phi(x) \phi(y)] | 0 \rangle$$

$$= 1 \cdot \hat{O}_1 + C_0(x-y) \mathbf{I}$$

3. Application to inclusive DIS

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle = \sum_{\{\alpha\}} C_{\{\alpha\}}^{\mu\nu} \left(\frac{M}{Q}\right)^{t-2}$$

twist $t =$ canonical dimension - spin of operator \hat{O}_i

		perturbative expansion			
		leading order LO	NLO	NNLO	...
twist expansion	leading twist 2	1	α_s	α_s^2	...
	subleading twist 3	$1/Q$			
	twist 4	$1/Q^2$			
			

parton model $\xrightarrow{\text{perturbative QCD}}$

OPE \downarrow

OPE rigorous proof only for inclusive processes. It can be effectively extended to all semi-inclusive hard processes.



4. Factorization

By applying the same technique of Wick theorem, it can be shown that the dominant contribution to the hadronic tensor of inclusive DIS comes from the so-called “handbag” diagram:

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle \sim$$

The diagram illustrates the handbag diagram for inclusive DIS. It shows a target with momentum \vec{P} interacting with a virtual photon γ^* of momentum q . The interaction is mediated by a parton with momentum k . The parton is shown as a red line connecting two vertices (blue circles). The target is represented by a yellow box labeled Φ . The diagram is divided into two regions by a vertical dashed line. The region to the right of the dashed line is labeled "hard interaction" in a cyan box. The region to the left of the dashed line is labeled "structure" in a yellow box.



4. Factorization

By applying the same technique of Wick theorem, it can be shown that the dominant contribution to the hadronic tensor of inclusive DIS comes from the so-called “handbag” diagram:

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle \sim$$

The diagram illustrates the handbag diagram for inclusive DIS. A target with momentum P is split into two parts, i and j , which are connected by a bilocal quark-quark correlator Φ . A hard interaction (represented by a blue circle) occurs between a photon γ^* with momentum q and a parton k from the target. The parton k is shown as a red line connecting the hard interaction to the structure function Φ . The structure function Φ is represented by a yellow box. The hard interaction is labeled "hard interaction" and the structure function is labeled "structure".

Φ bilocal quark-quark correlator:

$$\Phi_{ij}(k, P, S) = \int d^4 P_X \delta(P - k - P_X) \langle P, S | \bar{\psi}_f(0) | P_X \rangle \langle P_X | \psi_f(0) | P, S \rangle$$

$$= \int \frac{d^4 \xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$



4. Factorization

By applying the same technique of Wick theorem, it can be shown that the dominant contribution to the hadronic tensor of inclusive DIS comes from the so-called “handbag” diagram:

$$W^{\mu\nu} = \int d\xi e^{iq \cdot \xi} \langle P | [\hat{J}^\mu(\xi), \hat{J}^\nu(0)] | P \rangle \sim$$

The diagram illustrates the handbag diagram for the hadronic tensor. It shows a target with momentum P and a parton with momentum k interacting with a photon with momentum q via a hard interaction. The interaction is represented by a blue circle with a wavy line labeled γ^* . The parton is represented by a red line with momentum k . The structure function Φ is represented by a yellow box with labels i and j . The diagram is labeled "hard interaction" and "structure".

Φ bilocal quark-quark correlator:

$$\Phi_{ij}(k, P, S) = \int d^4 P_X \delta(P - k - P_X) \langle P, S | \bar{\psi}_f(0) | P_X \rangle \langle P_X | \psi_f(0) | P, S \rangle$$

$$= \int \frac{d^4 \xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$


By taking suitable projections, one can extract from the “structure” the leading-twist part, the subleading part at twist 3, at twist 4, etc..



parton-parton correlator



$$\Phi_f(k, P, S) = \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$




non-local correlator



parton-parton correlator



$$\Phi_f(k, P, S) = \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$



non-local correlator


➔ $\Phi(k, P, S)$ = linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)



parton-parton correlator



$$\Phi_f(k, P, S) = \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$



non-local correlator

- ➔ $\Phi(k, P, S)$ = linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)
- ➔ DIS regime $\rightarrow P^+$ dominant component; OPE on $\Phi(k, P, S) \rightarrow$ expansion in powers of M/P_+

Caveat


canonical OPE on local operators $\hat{\mathcal{O}}$; expansion in twist = $\dim(\hat{\mathcal{O}}) - \text{spin}(\hat{\mathcal{O}})$
Here, Φ is non-local, but can be expanded in local operators of same twist
“working” definition of twist = 2 + powers of M/P_+



parton-parton correlator



$$\Phi_f(k, P, S) = \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$



non-local correlator

➔ $\Phi(k, P, S)$ = linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)

➔ DIS regime $\rightarrow P^+$ dominant component; OPE on $\Phi(k, P, S) \rightarrow$ expansion in powers of M/P_+

Caveat


canonical OPE on local operators $\hat{\mathcal{O}}$; expansion in twist = $\dim(\hat{\mathcal{O}}) - \text{spin}(\hat{\mathcal{O}})$

Here, Φ is non-local, but can be expanded in local operators of same twist

“working” definition of twist = 2 + powers of M/P_+

➔ P^+ dominant component \rightarrow take partons collinear, $k^+ = xP^+$, and integrate other components:

$$\begin{aligned} \Phi(x, S) &= \int dk_+ dk_- d\mathbf{k}_T \delta(k_+ - xP_+) \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) \psi(0) | P, S \rangle \\ &= \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0} \end{aligned}$$




non-locality along “-” direction



parton-parton correlator



$$\Phi_f(k, P, S) = \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_f(\xi) \psi_f(0) | P, S \rangle$$



non-local correlator

➔ $\Phi(k, P, S)$ = linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)

➔ DIS regime $\rightarrow P^+$ dominant component; OPE on $\Phi(k, P, S) \rightarrow$ expansion in powers of M/P_+

Caveat


canonical OPE on local operators $\hat{\mathcal{O}}$; expansion in twist = $\dim(\hat{\mathcal{O}}) - \text{spin}(\hat{\mathcal{O}})$

Here, Φ is non-local, but can be expanded in local operators of same twist

“working” definition of twist = 2 + powers of M/P_+

➔ P^+ dominant component \rightarrow take partons collinear, $k^+ = xP^+$, and integrate other components:

$$\begin{aligned} \Phi(x, S) &= \int dk_+ dk_- d\mathbf{k}_T \delta(k_+ - xP_+) \int \frac{d^4\xi}{(2\pi)^4} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) \psi(0) | P, S \rangle && \Phi(k, P, S) \\ &= \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0} && \text{expanded in} \\ &&& \text{powers of } M/P_+ \end{aligned}$$



non-locality along “-” direction



OPE on $\Phi(k, P, S) \rightarrow$ expansion in powers of $M/P_+ \rightarrow$ keeping only leading twist

$$\Phi(x, S) = \int dk_+ dk_- d\mathbf{k}_T \delta(k_+ - xP_+) \Phi(k, P, S)$$

$$= \frac{1}{2} \left[f_1(x) \gamma_- + \right.$$

$$g_1(x) S_L \gamma_5 \gamma_- +$$

$$\left. \sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu] \quad h_1(x) i \sigma_{-\nu} \gamma_5 S_T^\nu \right]$$



OPE on $\Phi(k, P, S) \rightarrow$ expansion in powers of $M/P_+ \rightarrow$ keeping only leading twist

$$\begin{aligned}\Phi(x, S) &= \int dk_+ dk_- d\mathbf{k}_T \delta(k_+ - xP_+) \Phi(k, P, S) \\ &= \frac{1}{2} \left[f_1(x) \gamma_- + \right. \\ &\quad \left. g_1(x) S_L \gamma_5 \gamma_- + \right. \\ \sigma^{\mu\nu} &= \frac{i}{2} [\gamma^\mu, \gamma^\nu] \quad \left. h_1(x) i\sigma_{-\nu} \gamma_5 S_T^\nu \right]\end{aligned}$$

$$f_1(x) = \frac{1}{2} \text{Tr}[\Phi \gamma_+] \equiv \Phi^{[\gamma_+]}$$

$$S_L g_1(x) = \frac{1}{2} \text{Tr}[\Phi \gamma_+ \gamma_5] \equiv \Phi^{[\gamma_+ \gamma_5]}$$

$$(S_T)_i h_1(x) = \frac{1}{2} \text{Tr}[\Phi i\sigma_{+i} \gamma_5] \equiv \Phi^{[i\sigma_{+i} \gamma_5]}$$

Let's define

$$\Phi^{[\Gamma]}(x) = \frac{1}{P_+} \int dk^- d\mathbf{k}_T \text{Tr} \left[(\Phi(k, P, S))_{ji} (\Gamma)_{ij} \right] \Big|_{k^+ = xP^+}$$



OPE on $\Phi(k, P, S) \rightarrow$ expansion in powers of $M/P_+ \rightarrow$ keeping only leading twist

$$\Phi(x, S) = \int dk_+ dk_- d\mathbf{k}_T \delta(k_+ - xP_+) \Phi(k, P, S)$$

$$= \frac{1}{2} \left[f_1(x) \gamma_- + \right. \quad \text{unpolarized Parton Distribution Function (PDF)}$$

$$g_1(x) S_L \gamma_5 \gamma_- + \quad \text{longitudinally polarized PDF (requires hadron long. pol. } S_L)$$

$$\left. \sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu] \quad h_1(x) i\sigma_{-\nu} \gamma_5 S_T^\nu \right] \quad \text{transversely polarized PDF (requires hadron transv. pol. } S_T)$$


$$f_1(x) = \frac{1}{2} \text{Tr}[\Phi \gamma_+] \equiv \Phi^{[\gamma_+]} \quad \text{(fractional) momentum distribution}$$

$$S_L g_1(x) = \frac{1}{2} \text{Tr}[\Phi \gamma_+ \gamma_5] \equiv \Phi^{[\gamma_+ \gamma_5]} \quad \text{helicity distribution}$$

$$(S_T)_i h_1(x) = \frac{1}{2} \text{Tr}[\Phi i\sigma_{+i} \gamma_5] \equiv \Phi^{[i\sigma_{+i} \gamma_5]} \quad \text{transversity distribution}$$



The PDF table





PDFs ($x; Q^2$) at leading twist
for a spin-1/2 hadron (Nucleon)

polarizations
quark


•



nucleon




		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		
	L		$g_1 = \odot \rightarrow - \odot \rightarrow$	
	T			$h_1 = \odot \uparrow - \odot \uparrow$




PDFs ($x; Q^2$) at leading twist for a spin-1/2 hadron (Nucleon)

polarizations
nucleon



quark



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		
	L		$g_1 = \odot \rightarrow - \odot \rightarrow$	
	T			$h_1 = \odot \uparrow - \odot \uparrow$

probabilistic interpretation

probability density of finding an unpol. quark in an unpol. nucleon

probability density of finding a long. pol. quark in a long. pol. nucleon

probability density of finding a transv. pol. quark in a transv. pol. nucleon

N.B. Using appropriate projections $\Phi^{[\Gamma]}$ one can extract also subleading-twist PDFs, but no probabilistic interpretation



connection of PDFs with measurable structure functions

at leading order $\mathcal{O}(\alpha_s^0)$ and leading twist


$$F_{UU,T}(x_B, Q^2) = x_B \sum_q e_q^2 f_1^q(x_B, Q^2)$$

$$F_{UU,L}(x_B, Q^2) \approx 0$$

$$F_{LL}(x_B, Q^2) = x_B \sum_q e_q^2 g_1^q(x_B, Q^2)$$

$$F_{LT}(x_B, Q^2) \approx 0$$

hard cross section $d\hat{\sigma} = 1 + c_1\alpha_s + \dots$
produce $F_L, F_{LT} \neq 0$






connection of PDFs with measurable structure functions

at leading order $\mathcal{O}(\alpha_s^0)$ and leading twist

$$F_{UU,T}(x_B, Q^2) = x_B \sum_q e_q^2 f_1^q(x_B, Q^2) \quad F_{UU,L}(x_B, Q^2) \approx 0$$

$$F_{LL}(x_B, Q^2) = x_B \sum_q e_q^2 g_1^q(x_B, Q^2) \quad F_{LT}(x_B, Q^2) \approx 0$$

hard cross section $d\hat{\sigma} = 1 + c_1\alpha_s + \dots$
produce $F_L, F_{LT} \neq 0$



Transversity PDF does not appear in inclusive DIS cross section!

It happens because transverse polarization mixes quark helicities:

$$\langle \uparrow | \dots | \uparrow \rangle \propto \langle + | \dots | - \rangle, \langle - | \dots | + \rangle$$

chirality = helicity for a spin-1/2 object; hence, $h_1(x)$ is a chiral-odd PDF and can appear in the cross section only paired to another chiral-odd structure.

**Transversity is not suppressed (as expected in perturbative QCD as m_q/Q),
it can be extracted in processes with at least two hadrons**



The gauge link



$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$

this non-local operator is not color-gauge invariant under $\psi(x) \rightarrow e^{i\alpha^a(x)t^a} \psi(x) \equiv U(x) \psi(x)$



The gauge link



$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$


this non-local operator is not color-gauge invariant under $\psi(x) \rightarrow e^{i\alpha^a(x)t^a} \psi(x) \equiv U(x) \psi(x)$



$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) U_{[\xi,0]} \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$

gauge-link operator $U_{[a,b]} = \mathcal{P} \exp \left[-ig \int_a^b d\eta_\mu A^\mu(\eta) \right]$

it transforms as $U_{[\xi,0]} \rightarrow U(\xi) U_{[\xi,0]} U^\dagger(0)$ so that $\Phi(k, P, S)$ is invariant





The gauge link




$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$

this non-local operator is not color-gauge invariant under $\psi(x) \rightarrow e^{i\alpha^a(x)t^a} \psi(x) \equiv U(x) \psi(x)$



$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) U_{[\xi, 0]} \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$




gauge-link operator $U_{[a,b]} = \mathcal{P} \exp \left[-ig \int_a^b d\eta_\mu A^\mu(\eta) \right]$

it transforms as $U_{[\xi, 0]} \rightarrow U(\xi) U_{[\xi, 0]} U^\dagger(0)$ so that $\Phi(k, P, S)$ is invariant

$\Phi(x, S)$ involves only the LC “-“ direction: $\int d\xi_- \dots |_{\xi_+, \xi_T = 0}$

trick: $\Phi(x, S) \propto \langle P, S | \bar{\psi}(\xi_-) U_{[\xi_-, 0]} \psi(0) | P, S \rangle = \langle P, S | \bar{\psi}(\xi_-) U_{[\xi_-, \infty_-]} U_{[\infty_-, 0]} \psi(0) | P, S \rangle \equiv \langle P, S | \{ \bar{\psi}(\xi_-) \} \{ \psi(0) \} | P, S \rangle$





The gauge link



$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$

this non-local operator is not color-gauge invariant under $\psi(x) \rightarrow e^{i\alpha^a(x)t^a} \psi(x) \equiv U(x) \psi(x)$





$$\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}_j(\xi) U_{[\xi, 0]} \psi_i(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$$

gauge-link operator $U_{[a,b]} = \mathcal{P} \exp \left[-ig \int_a^b d\eta_\mu A^\mu(\eta) \right]$

it transforms as $U_{[\xi, 0]} \rightarrow U(\xi) U_{[\xi, 0]} U^\dagger(0)$ so that $\Phi(k, P, S)$ is invariant

$\Phi(x, S)$ involves only the LC “-“ direction: $\int d\xi_- \dots |_{\xi_+, \xi_T = 0}$

trick: $\Phi(x, S) \propto \langle P, S | \bar{\psi}(\xi_-) U_{[\xi_-, 0]} \psi(0) | P, S \rangle = \langle P, S | \bar{\psi}(\xi_-) U_{[\xi_-, \infty_-]} U_{[\infty_-, 0]} \psi(0) | P, S \rangle \equiv \langle P, S | \{ \bar{\psi}(\xi_-) \} \{ \psi(0) \} | P, S \rangle$



factorisation is preserved


$$\{ \psi(x_-) \} = U_{[x_-, \infty_-]} \psi(x_-)$$



Recap



- hadron structure better explored in processes with a hard scale (much bigger than involved masses, $Q^2 \gg M^2$) ; on the Light-Cone, it implies one dominant direction \rightarrow collinear framework natural choice






Recap



- hadron structure better explored in processes with a hard scale (much bigger than involved masses, $Q^2 \gg M^2$) ; on the Light-Cone, it implies one dominant direction \rightarrow collinear framework natural choice
- Example: inclusive DIS, cross section $d\sigma \sim L_{\mu\nu} W^{\mu\nu}$ can be parametrised in terms of 4 structure functions (including polarization)






Recap



- hadron structure better explored in processes with a hard scale (much bigger than involved masses, $Q^2 \gg M^2$) ; on the Light-Cone, it implies one dominant direction \rightarrow collinear framework natural choice
- Example: inclusive DIS, cross section $d\sigma \sim L_{\mu\nu} W^{\mu\nu}$ can be parametrised in terms of 4 structure functions (including polarization)
- OPE on $W^{\mu\nu} \rightarrow$ factorisation of hadron structure in parton-parton non-local correlator Φ . It can be made color-gauge invariant by inserting proper gauge link









Recap



- hadron structure better explored in processes with a hard scale (much bigger than involved masses, $Q^2 \gg M^2$) ; on the Light-Cone, it implies one dominant direction \rightarrow collinear framework natural choice
- Example: inclusive DIS, cross section $d\sigma \sim L_{\mu\nu} W^{\mu\nu}$ can be parametrised in terms of 4 structure functions (including polarization)
- OPE on $W^{\mu\nu} \rightarrow$ factorisation of hadron structure in parton-parton non-local correlator Φ . It can be made color-gauge invariant by inserting proper gauge link
- Expansion of Φ in powers of M/Q (effective twist) contains operator-definition of collinear PDFs, that can be extracted by suitable projections









Recap



- hadron structure better explored in processes with a hard scale (much bigger than involved masses, $Q^2 \gg M^2$) ; on the Light-Cone, it implies one dominant direction \rightarrow collinear framework natural choice
- Example: inclusive DIS, cross section $d\sigma \sim L_{\mu\nu} W^{\mu\nu}$ can be parametrised in terms of 4 structure functions (including polarization)
- OPE on $W^{\mu\nu} \rightarrow$ factorisation of hadron structure in parton-parton non-local correlator Φ . It can be made color-gauge invariant by inserting proper gauge link
- Expansion of Φ in powers of M/Q (effective twist) contains operator-definition of collinear PDFs, that can be extracted by suitable projections
- Leading-twist PDFs have nice probabilistic interpretations, and can be connected to structure functions (except the chiral-odd transversity PDF)



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		
	L		$g_1 = \ominus \dashrightarrow \ominus$	
	T			$h_1 = \uparrow \dashrightarrow \uparrow$



- Why TMDs ?




Evidences of going beyond the collinear framework

Example #1: the “Spin Crisis”

*Ashmann et al. (EMC),
P.L. B206 (88) 364*


- In 1988, the EMC Collaboration at CERN measures the F_{LL} structure function in the polarized inclusive DIS process $\bar{\mu} + \vec{p} \rightarrow \mu' + X$. Surprisingly, the sum of quark helicities Δq contributes at most 25% of spin 1/2 of the proton (depending on Q^2).
- There has been an intense activity to measure the gluon helicity Δg , which is currently known with a large error \rightarrow there is room for contribution from the orbital motion of partons
- Contribution from the orbital angular momentum of partons $L^q, L^g \rightarrow$ need to be sensitive also to intrinsic transverse components of parton momentum

$$\frac{1}{2} = \sum_q \left(\frac{1}{2} \Delta q(Q^2) + L^q(Q^2) \right) + \Delta g(Q^2) + L^g(Q^2)$$






Example #2: elastic p-p scattering



$$A_N = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow}$$


$p^\uparrow p \rightarrow p p$ versus $p^\downarrow p \rightarrow p p$

for a review, see
Krisch, E.P.J. **A31** (07) 417





Example #2: elastic p-p scattering




$$A_N = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow}$$

$p^\uparrow p \rightarrow p p$ versus $p^\downarrow p \rightarrow p p$

correlation between spin of the proton
and k_T of partons

\leftrightarrow orbital motion

for a review, see
Krisch, E.P.J. **A31** (07) 417






Evidences to go beyond collinear



Example #3: semi-inclusive p-p collisions






$$p^\uparrow p \rightarrow \pi X$$



$$A_N = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow} \quad \text{single-spin asymmetry}$$

perturbative QCD $\propto \frac{m_q}{p_T} \alpha_s \sim \mathcal{O}(10^{-3})$

Kane, Pumplin, Repko,
P.R.L. **41** ('78) 1689



Persisting also up to
 $\sqrt{s} = 200 \text{ GeV}$




Adams *et al.* (STAR),
PRL **92** (04) 171801



Evidences to go beyond collinear








Example #3: semi-inclusive p-p collisions



perturbative QCD $\propto \frac{m_q}{p_T} \alpha_s \sim \mathcal{O}(10^{-3})$

Kane, Pumplin, Repko, P.R.L. **41** ('78) 1689

Instead, large asymmetries observed.
Evidence of correlation between spin of the proton and k_T and flavor of partons



Persisting also up to $\sqrt{s} = 200$ GeV




Adams *et al.* (STAR), PRL **92** (04) 171801




- The “TMD zoo”
 - factorisation theorem and general properties
(generalising same steps that lead to PDFs)
 - specific properties



Need semi-inclusive process



- inclusive DIS: - **hard** scale $Q^2 = -q^2 \gg M^2$ to “see” partons
 - factorisation \rightarrow isolate PDFs
 - no further scale to probe proton interior




- semi-inclusive DIS (SIDIS):
 - **hard** scale $Q^2 = -q^2 \gg M^2$ to “see” partons
 - **soft** scale: detect hadron h with $P_{hT}^2 \sim M^2 \ll Q^2$
 - factorisation \rightarrow isolate TMDs

Ji, Yuan, Ma, P.R. D71 (05)

Rogers & Aybat, P.R. D83 (11)

Collins, “Foundations of Perturbative QCD” (11)

Echevarria, Idilbi, Scimemi, JHEP 1207 (12)



with these **two** scales, the process is factorizable into a hard photon-quark vertex and a quark \rightarrow hadron fragmentation

$$\mathbf{P}_{hT} = z \mathbf{k}_\perp + \mathbf{P}_\perp + \mathcal{O}(k_\perp^2/Q^2)$$


z = fractional energy of h
(analogous of x)

hadron P_{hT} arises from struck quark k_\perp and transverse momentum P_\perp generated during fragmentation

measure $P_{hT} \rightarrow$ get to k_\perp




The TMD framework



ECG



cardio
MR



A new paradigm:
**3D imaging of
hadron structure**
(in momentum space)

parametrised by
Transverse-Momentum
Dependent PDFs
TMD PDF($x, k_{\perp}; Q^2$)


parton
"longitudinal"
momentum

$$k^+ = x P^+$$



hard
collision

P^+ proton
momentum



parton
transverse
impulse

partons


\perp plane





SIDIS



one photon-exchange approximation



optical theorem



same invariants as inclusive DIS plus

$$z_h = \frac{P \cdot P_h}{P \cdot q}$$

“energy fraction” of fragmenting parton carried by final hadron

$$\frac{d\sigma}{dx_B dy d\phi_S dz_h d\phi_h dP_{hT}^2} = \frac{\alpha^2 y}{2z_h Q^4} L_{\mu\nu}(\ell, \ell', \lambda_e) W^{\mu\nu}(q, P, S, P_h)$$

new dependence


(for unpolarized hadron, $S_h=0$)





Example : SIDIS



one photon-exchange approximation



optical theorem



same invariants as inclusive DIS plus

$$z_h = \frac{P \cdot P_h}{P \cdot q}$$

“energy fraction” of fragmenting parton carried by final hadron

$$\frac{d\sigma}{dx_B dy d\phi_S dz_h d\phi_h dP_{hT}^2} = \frac{\alpha^2 y}{2z_h Q^4} L_{\mu\nu}(\ell, \ell', \lambda_e) W^{\mu\nu}(q, P, S, P_h)$$

new dependence

(for unpolarized hadron, $S_h=0$)

leptonic tensor

hadronic tensor



parametrised with **8** structure functions at leading twist (**18** including subleading twist)



SIDIS cross section



$$\frac{d\sigma}{dx_B dy d\phi_S dz_h d\phi_h dP_{hT}^2} =$$


$$= \frac{\alpha^2}{x_B y Q^2} \left[A(y) F_{UU,T} + B(y) \cos 2\phi_h F_{UU}^{\cos 2\phi_h} \right.$$

$$+ S_L \sin 2\phi_h F_{UL}^{\sin 2\phi_h}$$

$$+ \lambda_e S_L C(y) F_{LL}$$

$$+ S_T \left[A(y) \sin(\phi_h - \phi_S) F_{UT,T}^{\sin(\phi_h - \phi_S)} + B(y) \sin(\phi_h + \phi_S) F_{UT}^{\sin(\phi_h + \phi_S)} + B(y) \sin(3\phi_h - \phi_S) F_{UT}^{\sin(3\phi_h - \phi_S)} \right]$$

$$+ \lambda_e S_T C(y) \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)} \left. \right] + \mathcal{O}\left(\frac{M}{Q}\right)$$



each $F_{XY,Z}(x_B, z_h, P_{hT}^2, Q^2)$



SIDIS cross section



$$\frac{d\sigma}{dx_B dy d\phi_S dz_h d\phi_h dP_{hT}^2} =$$


$$= \frac{\alpha^2}{x_B y Q^2} \left[A(y) F_{UU,T} + B(y) \cos 2\phi_h F_{UU}^{\cos 2\phi_h} \right.$$

$$+ S_L \sin 2\phi_h F_{UL}^{\sin 2\phi_h}$$

$$+ \lambda_e S_L C(y) F_{LL}$$

$$+ S_T \left[A(y) \sin(\phi_h - \phi_S) F_{UT,T}^{\sin(\phi_h - \phi_S)} + B(y) \sin(\phi_h + \phi_S) F_{UT}^{\sin(\phi_h + \phi_S)} + B(y) \sin(3\phi_h - \phi_S) F_{UT}^{\sin(3\phi_h - \phi_S)} \right]$$

$$+ \lambda_e S_T C(y) \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)} \left. \right] + \mathcal{O}\left(\frac{M}{Q}\right)$$




each $F_{XY,Z}(x_B, z_h, P_{hT}^2, Q^2)$



SIDIS : factorisation




OPE not possible, use diagrammatic approach (select dominant diagram by counting powers of divergences)



$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{C} \left[\text{Tr} \left[\Phi(x_B, \mathbf{k}_\perp, S) \gamma^\mu \Delta(z_h, \mathbf{P}_\perp) \gamma^\nu \right] \right] \quad \mathcal{C}[\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]$$



OPE not possible, use diagrammatic approach (select dominant diagram by counting powers of divergences)




$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{C} \left[\text{Tr} \left[\Phi(x_B, \mathbf{k}_\perp, S) \gamma^\mu \Delta(z_h, \mathbf{P}_\perp) \gamma^\nu \right] \right] \quad \mathcal{C}[\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]$$

non-local correlator:

from collinear $\Phi(x, S) = \int \frac{d\xi_-}{2\pi} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi,0]} \psi(0) | P, S \rangle_{\xi_+ = \xi_T = 0}$

to $\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi,0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$






SIDIS : factorisation




non-local correlators

$$z_h \sim z = \frac{P_{h-}}{k_-} \quad \begin{array}{l} \xi_+ \leftrightarrow k_- \\ \xi_T \leftrightarrow k_\perp \end{array}$$


$$\Delta(z, \mathbf{k}_\perp) = \sum_X \int \frac{d\xi_+ d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle 0 | \psi(0) | X, P_h \rangle \langle X, P_h | \bar{\psi}(\xi) | 0 \rangle_{\xi_- = 0}$$



hard cross section
 $d\hat{\sigma} = 1 + c_1 \alpha_s + \dots$



$$\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi, 0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$$



$$x_B \sim x = \frac{k_+}{P_+} \quad \begin{array}{l} \xi_- \leftrightarrow k_+ \\ \xi_T \leftrightarrow k_\perp \end{array}$$



SIDIS : factorisation





non-local correlators

$$z_h \sim z = \frac{P_{h-}}{k_-} \quad \begin{array}{l} \xi_+ \leftrightarrow k_- \\ \xi_T \leftrightarrow k_\perp \end{array}$$

$$\Delta(z, \mathbf{k}_\perp) = \sum_X \int \frac{d\xi_+ d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle 0 | \psi(0) | X, P_h \rangle \langle X, P_h | \bar{\psi}(\xi) | 0 \rangle_{\xi_- = 0}$$


hard cross section
 $d\hat{\sigma} = 1 + c_1 \alpha_s + \dots$



↑ flipping LC-dominant direction



$$\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi, 0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$$



$$x_B \sim x = \frac{k_+}{P_+} \quad \begin{array}{l} \xi_- \leftrightarrow k_+ \\ \xi_T \leftrightarrow k_\perp \end{array}$$



flipping LC-dominant direction

Definitions: $x = \frac{k^+}{P^+}$ $x_B = \frac{-q^2}{2P \cdot q} = \frac{-2q^+q^-}{2P^+q^-} = -\frac{q^+}{P^+}$

Parton model \rightarrow elastic kinematics $\rightarrow x \approx x_B$



flipping LC-dominant direction

Definitions: $x = \frac{k^+}{P^+} \quad x_B = \frac{-q^2}{2P \cdot q} = \frac{-2q^+q^-}{2P^+q^-} = -\frac{q^+}{P^+}$

Parton model \rightarrow elastic kinematics $\rightarrow x \approx x_B$

initial parton $k = \{k^+, k^-, \mathbf{k}_\perp\} = \left\{ xP^+, \frac{k^2 + \mathbf{k}_\perp^2}{2xP^+}, \mathbf{k}_\perp \right\} \approx \{xP^+, 0, \mathbf{k}_\perp\}$

$k^2 = 2k^+k^- - \mathbf{k}_\perp^2$

+ component dominant



flipping LC-dominant direction

Definitions: $x = \frac{k^+}{P^+} \quad x_B = \frac{-q^2}{2P \cdot q} = \frac{-2q^+q^-}{2P^+q^-} = -\frac{q^+}{P^+}$

Parton model \rightarrow elastic kinematics $\rightarrow x \approx x_B$

initial parton

$$k = \{k^+, k^-, \mathbf{k}_\perp\} = \left\{ xP^+, \frac{k^2 + \mathbf{k}_\perp^2}{2xP^+}, \mathbf{k}_\perp \right\} \approx \{xP^+, 0, \mathbf{k}_\perp\}$$

$k^2 = 2k^+k^- - \mathbf{k}_\perp^2$

+ component dominant

momentum transfer

$$q = \{q^+, q^-, \mathbf{0}_T\} = \left\{ -xP^+, \frac{Q^2}{2xP^+}, \mathbf{0}_T \right\}$$

$q^2 = 2q^+q^-$



flipping LC-dominant direction

Definitions: $x = \frac{k^+}{P^+} \quad x_B = \frac{-q^2}{2P \cdot q} = \frac{-2q^+q^-}{2P^+q^-} = -\frac{q^+}{P^+}$

Parton model \rightarrow elastic kinematics $\rightarrow x \approx x_B$

initial parton

$$k = \{k^+, k^-, \mathbf{k}_\perp\} = \left\{ xP^+, \frac{k^2 + \mathbf{k}_\perp^2}{2xP^+}, \mathbf{k}_\perp \right\} \approx \{xP^+, 0, \mathbf{k}_\perp\}$$

$k^2 = 2k^+k^- - \mathbf{k}_\perp^2$

+ component dominant

momentum transfer

$$q = \{q^+, q^-, \mathbf{0}_T\} = \left\{ -xP^+, \frac{Q^2}{2xP^+}, \mathbf{0}_T \right\}$$

$q^2 = 2q^+q^-$

final parton

$$k' = k + q = \left\{ 0, \frac{Q^2}{2xP^+}, \mathbf{k}_\perp \right\} \approx \{0, Q, \mathbf{k}_\perp\}$$


- component dominant



parton-parton correlator



linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)
 expansion of Φ in powers of M/P_+ . At leading twist:




$$\Phi(x, \mathbf{k}_\perp, S) = \frac{1}{2} \left[f_1 \gamma_- - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M} \gamma_- \right. \\
 + g_{1L} S_L \gamma_5 \gamma_- + g_{1T} \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} \gamma_5 \gamma_- \\
 \left. + h_{1T} i \sigma_{-\nu} \gamma_5 S_T^\nu + h_{1L}^\perp i \sigma_{-\nu} \gamma_5 S_L \frac{k_\perp^\nu}{M} \right. \\
 \left. + h_{1T}^\perp \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} i \sigma_{-\nu} \gamma_5 \frac{k_\perp^\nu}{M} - h_1^\perp \sigma_{-\nu} \frac{k_\perp^\nu}{M} \right]$$

$$\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$$


Notations:



parton-parton correlator



linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)
 expansion of Φ in powers of M/P_+ . At leading twist:



$$\Phi(x, \mathbf{k}_\perp, S) = \frac{1}{2} \left[f_1 \gamma_- - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M} \gamma_- \right. \\
 + g_{1L} S_L \gamma_5 \gamma_- + g_{1T} \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} \gamma_5 \gamma_- \\
 + h_{1T} i \sigma_{-\nu} \gamma_5 S_T^\nu + h_{1L}^\perp i \sigma_{-\nu} \gamma_5 S_L \frac{k_\perp^\nu}{M} \\
 \left. + h_{1T}^\perp \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} i \sigma_{-\nu} \gamma_5 \frac{k_\perp^\nu}{M} - h_1^\perp \sigma_{-\nu} \frac{k_\perp^\nu}{M} \right]$$

$$\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$$

Notations: $t_{1X}^{(\perp)}(x, \mathbf{k}_\perp^2)$ waited by k_\perp^i

leading twist $X = L$ longitudinally polarized hadron
 $X = T$ transversely polarized hadron


$t = f$ unpolarized parton
 $t = g$ longitudinally polarized parton
 $t = h$ transversely polarized parton



parton-parton correlator



linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)
 expansion of Φ in powers of M/P_+ . At leading twist:



$$\Phi(x, \mathbf{k}_\perp, S) = \frac{1}{2} \left[f_1 \gamma_- - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M} \gamma_- \right.$$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \equiv \Phi^{[t]} \rightarrow \text{2 TMDPDFs for unpol. parton}$$

$$+ g_{1L} S_L \gamma_5 \gamma_- + g_{1T} \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} \gamma_5 \gamma_-$$

$$\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$$

$$+ h_{1T} i \sigma_{-\nu} \gamma_5 S_T^\nu + h_{1L}^\perp i \sigma_{-\nu} \gamma_5 S_L \frac{k_\perp^\nu}{M}$$

$$+ h_{1T}^\perp \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} i \sigma_{-\nu} \gamma_5 \frac{k_\perp^\nu}{M} - h_1^\perp \sigma_{-\nu} \frac{k_\perp^\nu}{M} \left. \right]$$

Notations:

$$t_{1X}^{(\perp)}(x, \mathbf{k}_\perp^2) \quad \text{waited by } k_\perp^i$$

leading twist

$X = L$ longitudinally polarized hadron

$X = T$ transversely polarized hadron

$t = f$ unpolarized parton

$t = g$ longitudinally polarized parton


$t = h$ transversely polarized parton



parton-parton correlator



linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)
 expansion of Φ in powers of M/P_+ . At leading twist:



$$\Phi(x, \mathbf{k}_\perp, S) = \frac{1}{2} \left[f_1 \gamma_- - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M} \gamma_- \right.$$

$$\left. + g_{1L} S_L \gamma_5 \gamma_- + g_{1T} \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} \gamma_5 \gamma_- \right.$$

$$\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$$

$$\left. + h_{1T} i\sigma_{-\nu} \gamma_5 S_T^\nu + h_{1L}^\perp i\sigma_{-\nu} \gamma_5 S_L \frac{k_\perp^\nu}{M} \right.$$

$$\left. + h_{1T}^\perp \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} i\sigma_{-\nu} \gamma_5 \frac{k_\perp^\nu}{M} - h_1^\perp \sigma_{-\nu} \frac{k_\perp^\nu}{M} \right]$$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \equiv \Phi^{[\gamma_+]} \rightarrow \text{2 TMDPDFs for unpol. parton}$$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+ \gamma_5] \equiv \Phi^{[\gamma_+ \gamma_5]} \rightarrow \text{2 TMDPDFs for long. pol. parton}$$

$$\frac{1}{2} \text{Tr}[\Phi i\sigma_{+i} \gamma_5] \equiv \Phi^{[i\sigma_{+i} \gamma_5]} \rightarrow \text{4 TMDPDFs for transv. pol. parton along } i$$

Notations:

$$t_{1X}^{(\perp)}(x, \mathbf{k}_\perp^2) \quad \text{waited by } k_\perp^i$$

leading twist

$X = L$ longitudinally polarized hadron
 $X = T$ transversely polarized hadron

$t = f$ unpolarized parton

$t = g$ longitudinally polarized parton

$t = h$ transversely polarized parton




parton-parton correlator



linear combination of all tensor structures with k, P, S , subject to Hermiticity and parity-invariance (see later about time reversal)

expansion of Φ in powers of M/P_+ . At leading twist:



$$\Phi(x, \mathbf{k}_\perp, S) = \frac{1}{2} \left[f_1 \gamma_- - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M} \gamma_- \right.$$

$$\left. + g_{1L} S_L \gamma_5 \gamma_- + g_{1T} \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} \gamma_5 \gamma_- \right.$$

$$\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$$

$$\left. + h_{1T} i \sigma_{-\nu} \gamma_5 S_T^\nu + h_{1L}^\perp i \sigma_{-\nu} \gamma_5 S_L \frac{k_\perp^\nu}{M} \right.$$

$$\left. + h_{1T}^\perp \frac{\mathbf{k}_\perp \cdot \mathbf{S}_T}{M} i \sigma_{-\nu} \gamma_5 \frac{k_\perp^\nu}{M} - h_{1L}^\perp \sigma_{-\nu} \frac{k_\perp^\nu}{M} \right]$$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \equiv \Phi^{[\gamma_+]} \rightarrow \text{2 TMDPDFs for unpol. parton}$$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+ \gamma_5] \equiv \Phi^{[\gamma_+ \gamma_5]} \rightarrow \text{2 TMDPDFs for long. pol. parton}$$

$$\frac{1}{2} \text{Tr}[\Phi i \sigma_{+i} \gamma_5] \equiv \Phi^{[i \sigma_{+i} \gamma_5]} \rightarrow \text{4 TMDPDFs for transv. pol. parton along } i$$

Notations:

$$t_{1X}^{(\perp)}(x, \mathbf{k}_\perp^2) \quad \text{waited by } k_\perp^i$$

leading twist

$X = L$ longitudinally polarized hadron
 $X = T$ transversely polarized hadron

$t = f$ unpolarized parton


$t = g$ longitudinally polarized parton

$t = h$ transversely polarized parton

survive upon $\int d\mathbf{k}_\perp \rightarrow$ collinear PDF




The TMD PDF table



TMD PDFs ($x, \mathbf{k}_\perp; Q^2$) at leading twist for a spin-1/2 hadron (Nucleon)

polarizations
nucleon



quark



Mulders & Tangerman, N.P. **B461** (96)
Boer & Mulders, P.R. **D57** (98)


		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow \ominus - \downarrow \ominus$
	L		$g_1 = \odot \rightarrow - \ominus \rightarrow$	$h_{1L}^\perp = \uparrow \rightarrow - \downarrow \rightarrow$
	T	$f_{1T}^\perp = \uparrow \odot - \downarrow \ominus$	$g_{1T} = \uparrow \rightarrow - \downarrow \rightarrow$	$h_1 = \uparrow \uparrow \ominus - \downarrow \uparrow \ominus$ $h_{1T}^\perp = \uparrow \rightarrow \uparrow \ominus - \downarrow \rightarrow \uparrow \ominus$

Each entry has a nice probabilistic interpretation

N.B. Using appropriate projections $\Phi^{[\Gamma]}$ one can extract also subleading-twist TMD PDFs, but no probabilistic interpretation




The TMD PDF table



TMD PDFs $(x, \mathbf{k}_\perp; Q^2)$ at leading twist for a spin-1/2 hadron (Nucleon)

polarizations
nucleon



quark



Mulders & Tangerman, N.P. **B461** (96)
Boer & Mulders, P.R. **D57** (98)

		Quark polarization			nomenclature
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)	
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow \ominus - \downarrow \ominus$	no-name Boer-Mulders
	L		$g_1 = \odot \rightarrow - \ominus \rightarrow$	$h_{1L}^\perp = \uparrow \rightarrow - \downarrow \rightarrow$	helicity Kotzinian-Mulders
	T	$f_{1T}^\perp = \uparrow \odot - \downarrow \ominus$	$g_{1T} = \uparrow \rightarrow - \downarrow \rightarrow$	$h_1 = \uparrow \uparrow \ominus - \downarrow \uparrow \ominus$ $h_{1T}^\perp = \uparrow \rightarrow \ominus - \downarrow \rightarrow \ominus$	transversity pretzelosity

Sivers


worm gear

Each entry has a nice probabilistic interpretation

N.B. Using appropriate projections $\Phi^{[\Gamma]}$ one can extract also subleading-twist TMD PDFs, but no probabilistic interpretation



The TMD FF table



TMD FFs ($z, P_{\perp}; Q^2$) at leading twist (and $S_h \leq 1/2$)

polarizations
hadron

quark



Mulders & Tangerman, N.P. **B461** (96)
Boer & Mulders, P.R. **D57** (98)

		Quark polarization			nomenclature	
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)		
Nucleon Polarization	U	D_1		H_1^{\perp} -	no-name	Collins
	L		G_{1L} -	H_{1L}^{\perp} -
	T	D_{1T}^{\perp} -	G_{1T} -	H_1 - H_{1T}^{\perp} -

polarising FF


...

Each entry has a nice probabilistic interpretation

N.B. Using appropriate projections $\Phi^{[\Gamma]}$ one can extract also subleading-twist TMD FFs, but no probabilistic interpretation




The TMD PDF table



TMD PDFs ($x, \mathbf{k}_\perp; Q^2$) at leading twist for a spin-1/2 hadron (Nucleon)

polarizations
nucleon



quark



Mulders & Tangerman, N.P. **B461** (96)
Boer & Mulders, P.R. **D57** (98)

		Quark polarization			nomenclature
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)	
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow \ominus - \downarrow \ominus$	no-name Boer-Mulders
	L		$g_1 = \odot \rightarrow - \ominus \rightarrow$	$h_{1L}^\perp = \uparrow \rightarrow - \downarrow \rightarrow$	helicity Kotzinian-Mulders
	T	$f_{1T}^\perp = \uparrow \odot - \downarrow \ominus$	$g_{1T} = \uparrow \rightarrow - \downarrow \rightarrow$	$h_1 = \uparrow \uparrow \ominus - \downarrow \uparrow \ominus$ $h_{1T}^\perp = \uparrow \rightarrow \uparrow \ominus - \downarrow \rightarrow \uparrow \ominus$	transversity pretzelosity

Sivers

worm gear

Each entry has a nice probabilistic interpretation

N.B. Using appropriate projections $\Phi^{[\Gamma]}$ one can extract also subleading-twist TMD PDFs, but no probabilistic interpretation





The unpolarized TMD PDF



polarizations

quark   

nucleon

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$

$f_1^q(x, \mathbf{k}_\perp^2)$ probability density of finding a quark q with “longitudinal” (along “+” LC direction) fraction x of nucleon momentum, and transverse momentum \mathbf{k}_\perp



The Sivers TMD PDF




polarizations

quark




nucleon



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \rightarrow f_1 - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M}$$




$$\mathbf{S}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$



The Sivers TMD PDF



polarizations
nucleon





quark




		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$

Bacchetta et al.,
P.L. **B827** (22) 136961,
arXiv:2004.14278



$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \rightarrow f_1 - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M}$$



$$\mathbf{S}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$

Sivers effect: how the momentum distribution of quarks is distorted by the transverse polarization of parent nucleon (“spin-orbit” correlation)

Sivers $f_{1T}^\perp \rightarrow$ indirect access to quark orbital angular momentum

Burkardt, P.R. **D66** (2002) 114005;
N.P. **A735** (2004) 185

Bacchetta & Radici, P.R.L. **107** (2011) 212001
Ji et al., N.P. **B652** (2003) 383



The Boer-Mulders TMD PDF



polarizations

quark • •→ •↑

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$


nucleon

•

•→

•↑

$$\frac{1}{2} \text{Tr}[\Phi i\sigma_{+i}\gamma_5] \rightarrow \dots + h_1^\perp \frac{(\mathbf{k}_\perp \times \mathbf{s}_T) \cdot \hat{\mathbf{P}}}{M}$$



$$\mathbf{s}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$

Boer-Mulders effect: “spin-orbit” correlation at partonic level




Forbidden combinations



polarizations

nucleon



quark



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$?	$h_1^\perp = \uparrow - \downarrow$
	L	?	$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$

Why?




Forbidden combinations



polarizations

nucleon



quark • •→ •↑

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$? *	$h_1^\perp = \uparrow - \downarrow$
	L	?	$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \searrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \uparrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \searrow$

Why?
prohibited by
parity invariance

$$\mathbf{S}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$

$$\mathbf{S}_L \cdot \mathbf{k}_\perp \times \mathbf{P} = 0$$

not enough
vectors for f_{1L}^\perp !

* similarly for “swapped” combination



The chiral-odd TMD PDFs



polarizations

quark

nucleon

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \text{dot}$		$h_1^\perp = \text{dot with up arrow} - \text{dot with down arrow}$
	L		$g_1 = \text{dot with right arrow} - \text{dot with left arrow}$	$h_{1L}^\perp = \text{dot with up and right arrow} - \text{dot with up and left arrow}$
	T	$f_{1T}^\perp = \text{dot with up arrow} - \text{dot with down arrow}$	$g_{1T} = \text{dot with right arrow} - \text{dot with left arrow}$	$h_1 = \text{dot with up arrow} - \text{dot with down arrow}$ $h_{1T}^\perp = \text{dot with up and right arrow} - \text{dot with up and left arrow}$

all TMD PDFs belonging to right column involve transverse polarization of quarks, hence they are “chiral-odd” and are suppressed in perturbative QCD as m_q/Q .

Similarly to transversity h_1 , they can appear in the cross section at leading twist if paired to another chiral-odd structure. For SIDIS, they must be paired to a chiral-odd TMD FF.



Transversity




polarizations

quark



nucleon




		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \searrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$

- transversity is the prototype of chiral-odd structures
- the only chiral-odd structure that survives in collinear kinematics
- only way to determine the tensor charge $\delta^q(Q^2) = \int_0^1 dx h_1^{q-\bar{q}}(x, Q^2)$

Transversity properties

both defined in
Infinite Mom. Frame



boosted Nucleon

$$g_1 = \text{[Diagram 1]} - \text{[Diagram 2]}$$


helicity

$$h_1 = \text{[Diagram 3]} - \text{[Diagram 4]}$$


transversity

Transversity properties


both defined in
Infinite Mom. Frame



boosted Nucleon



helicity



transversity

=

Non-relativistic theory:
boosts & rotations commute




Differences


=> info on relativistic motion of quarks

Transversity properties


both defined in
Infinite Mom. Frame



boosted Nucleon




helicity



transversity

=


Non-relativistic theory:
boosts & rotations commute



Differences

=> info on relativistic motion of quarks


In a spin-1/2 hadron,
no transversity of gluons



singlet and
non-singlet
evolution

only
non-singlet
evolution


In a spin-1 hadron, gluon transversity possible
because transverse tensor polariz. => $\Delta\lambda=2$
but $h_{1,TT}^g \equiv h_1^g$ is only a TMD and T-odd




Jaffe & Manohar (1989), Artru & Mekhfi (1990), Bacchetta & Mulders (2000)

Transversity properties


both defined in
Infinite Mom. Frame



boosted Nucleon




helicity



transversity

=



Non-relativistic theory:
boosts & rotations commute



Differences

=> info on relativistic motion of quarks


In a spin-1/2 hadron,
no transversity of gluons



singlet and
non-singlet
evolution

only
non-singlet
evolution


In a spin-1 hadron, gluon transversity possible
because transverse tensor polariz. => $\Delta\lambda=2$
but $h_{1,TT}^g \equiv h_1^g$ is only a TMD and T-odd




Jaffe & Manohar (1989), Artru & Mekhfi (1990), Bacchetta & Mulders (2000)

Soffer bound: $|h_1| \leq \frac{1}{2}(f_1 + g_1)$ for any (x, Q^2)

Transversity properties



helicity



transversity

charges connected to hadronic matrix elements of local operators (calculable on lattice)

$$\langle P, S_L | \bar{q} \gamma^\mu \gamma_5 q | P, S_L \rangle = S_L P^\mu g_A^q$$

axial current \Leftrightarrow axial charge

$$= S_L P^\mu \int_0^1 dx g_1^{q+\bar{q}}(x, Q^2)$$

connected to C-even structure


$$\langle P, S | \bar{q} \sigma^{\mu\nu} q | P, S \rangle = P^{[\mu} S^{\nu]} \delta^q(Q^2)$$

tensor current \Leftrightarrow tensor charge


$$= P^{[\mu} S^{\nu]} \int_0^1 dx h_1^{q-\bar{q}}(x, Q^2)$$

connected to C-odd structure

Transversity properties



helicity



transversity

charges connected to hadronic matrix elements of local operators (calculable on lattice)

$$\langle P, S_L | \bar{q} \gamma^\mu \gamma_5 q | P, S_L \rangle = S_L P^\mu g_A^q$$

axial current \Leftrightarrow axial charge

$$= S_L P^\mu \int_0^1 dx g_1^{q+\bar{q}}(x, Q^2)$$

connected to C-even structure

$$\langle P, S | \bar{q} \sigma^{\mu\nu} q | P, S \rangle = P^{[\mu} S^{\nu]} \delta^q(Q^2)$$

tensor current \Leftrightarrow tensor charge

$$= P^{[\mu} S^{\nu]} \int_0^1 dx h_1^{q-\bar{q}}(x, Q^2)$$

connected to C-odd structure

anomalous dim. $\Delta\gamma^{(1)} = 0$


$\Rightarrow g_A^q$ is constant

anomalous dim. $\delta\gamma^{(1)} = -C_F/2$

$\Rightarrow \delta^q$ scales with Q^2


$$C_F = \frac{N_c^2 - 1}{2N_c}$$

Transversity properties



helicity

\neq



transversity

charges connected to hadronic matrix elements of local operators (calculable on lattice)

$$\langle P, S_L | \bar{q} \gamma^\mu \gamma_5 q | P, S_L \rangle = S_L P^\mu g_A^q$$

axial current \Leftrightarrow axial charge

$$= S_L P^\mu \int_0^1 dx g_1^{q+\bar{q}}(x, Q^2)$$

connected to C-even structure

$$\langle P, S | \bar{q} \sigma^{\mu\nu} q | P, S \rangle = P^{[\mu} S^{\nu]} \delta^q(Q^2)$$

tensor current \Leftrightarrow tensor charge

$$= P^{[\mu} S^{\nu]} \int_0^1 dx h_1^{q-\bar{q}}(x, Q^2)$$

connected to C-odd structure

anomalous dim. $\Delta\gamma^{(1)} = 0$

$\Rightarrow g_A^q$ is constant

anomalous dim. $\delta\gamma^{(1)} = -C_F/2$

$\Rightarrow \delta^q$ scales with Q^2


$$C_F = \frac{N_c^2 - 1}{2N_c}$$

helicity and transversity are very different !

Potential for BSM discovery ?

Tensor (and chiral-odd) structures do not appear in the Standard Model Lagrangian at tree level.


Is it a possible low-energy footprint of BSM physics at higher scale ?



Potential for BSM discovery ?

Tensor (and chiral-odd) structures do not appear in the Standard Model Lagrangian at tree level.

Is it a possible low-energy footprint of BSM physics at higher scale ?



neutron β -decay

$n \rightarrow p e^- \bar{\nu}_e$

$$\mathcal{L}_{\text{SM}} \sim G_F V_{ud} \bar{e} \gamma^\mu (1 - \gamma_5) \nu_e \bar{p} \gamma_\mu (1 - \gamma_5) n$$


$$+ \mathcal{L}_{\text{eff}} \sim G_F V_{ud} g_T \varepsilon_T \bar{e} \sigma^{\mu\nu} \nu_e \bar{p} \sigma_{\mu\nu} n$$

precision \Rightarrow BSM scale $\frac{M_W^2}{M_{\text{BSM}}^2} \approx g_T \varepsilon_T$ \swarrow BSM coupling ?
 \nwarrow $g_T = \delta u - \delta d$ isovector tensor charge

Potential for BSM discovery ?

Tensor (and chiral-odd) structures do not appear in the Standard Model Lagrangian at tree level.

Is it a possible low-energy footprint of BSM physics at higher scale ?



neutron β -decay

$$n \rightarrow p e^- \bar{\nu}_e$$

$$\mathcal{L}_{\text{SM}} \sim G_F V_{ud} \bar{e} \gamma^\mu (1 - \gamma_5) \nu_e \bar{p} \gamma_\mu (1 - \gamma_5) n$$

$$+ \mathcal{L}_{\text{eff}} \sim G_F V_{ud} g_T \varepsilon_T \bar{e} \sigma^{\mu\nu} \nu_e \bar{p} \sigma_{\mu\nu} n$$

precision \Rightarrow BSM scale $\frac{M_W^2}{M_{\text{BSM}}^2} \approx g_T \varepsilon_T$ \swarrow BSM coupling ?

$g_T = \delta u - \delta d$ \swarrow isovector tensor charge

SMEFT with strong CP violation

permanent Electric Dipole Mom.

$$\mathcal{L}_{\text{SMEFT}} \rightarrow \sum_{f=u,d,s,c} d_f \bar{\psi}_f \sigma_{\mu\nu} \gamma_5 \psi_f F^{\mu\nu} \quad ?$$

\swarrow quark EDM

neutron EDM $d_n = \delta u d_u + \delta d d_d + \delta s d_s + \dots$

exp. data + **tensor charge** \Rightarrow constrain amount of CP violation



The Sivers TMD PDF




polarizations

quark




nucleon




		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$

$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \rightarrow f_1 - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M}$$



$$\mathbf{S}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$



$$\mathbf{S}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$

$$\frac{1}{2} \text{Tr}[\Phi i\sigma_{+i}\gamma_5] \rightarrow \dots + h_1^\perp \frac{(\mathbf{k}_\perp \times \mathbf{s}_T) \cdot \hat{\mathbf{P}}}{M}$$



Sivers and Boer-Mulders TMD PDFs vanish without gauge link U


$$\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi, 0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$$

$$U_{[a,b]} = \mathcal{P} \exp \left[-ig \int_a^b d\eta_\mu A^\mu(\eta) \right]$$

They are generated by interference of different channels.

(for example, f_{1T}^\perp can be reproduced by interference of model LC wave functions with different orbital angular momentum)

Gauge link U represents the residual color interactions that generate the necessary phase difference for the interference. As such, time reversal puts no constraints on these structures.



Sivers and Boer-Mulders TMD PDFs are conventionally named “T-odd” TMD PDFs

Boer & Mulders, P.R. D57 (98)




The gauge link



$$\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi,0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$$

TMD factorisation for SIDIS process suggests a trick similar to collinear framework case:

$$\begin{aligned} \langle P, S | \bar{\psi}(\xi) U_{[\xi,0]} \psi(0) | P, S \rangle &= \langle P, S | \bar{\psi}(\xi) U_{[\xi, \infty_-]} U_{[\infty_-, \infty_T]} U_{[\infty_T, \infty_-]} U_{[\infty_-, 0]} \psi(0) | P, S \rangle \\ &= \langle P, S | \{ \bar{\psi}(\xi) \} \{ \psi(0) \} | P, S \rangle \end{aligned}$$





The gauge link




$$\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi,0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$$

TMD factorisation for SIDIS process suggests a trick similar to collinear framework case:

$$\begin{aligned} \langle P, S | \bar{\psi}(\xi) U_{[\xi,0]} \psi(0) | P, S \rangle &= \langle P, S | \bar{\psi}(\xi) U_{[\xi, \infty_-]} U_{[\infty_-, \infty_T]} U_{[\infty_T, \infty_-]} U_{[\infty_-, 0]} \psi(0) | P, S \rangle \\ &= \langle P, S | \{ \bar{\psi}(\xi) \} \{ \psi(0) \} | P, S \rangle \end{aligned}$$

In Drell-Yan process, TMD factorisation gives the following path for gauge link:






The gauge link




$$\Phi(x, \mathbf{k}_\perp, S) = \int \frac{d\xi_- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P, S | \bar{\psi}(\xi) U_{[\xi, 0]} \psi(0) | P, S \rangle_{\xi_+ = 0}$$

TMD factorisation for SIDIS process suggests a trick similar to collinear framework case:

$$\begin{aligned} \langle P, S | \bar{\psi}(\xi) U_{[\xi, 0]} \psi(0) | P, S \rangle &= \langle P, S | \bar{\psi}(\xi) U_{[\xi, \infty\xi_-]} U_{[\infty\xi_-, \infty T]} U_{[\infty T, \infty_-]} U_{[\infty_-, 0]} \psi(0) | P, S \rangle \\ &= \langle P, S | \{ \bar{\psi}(\xi) \} \{ \psi(0) \} | P, S \rangle \end{aligned}$$



In Drell-Yan process, TMD factorisation gives the following path for gauge link:



Notations: gauge link $U_{[+]}$ for SIDIS; $U_{[-]}$ for Drell-Yan

Important result: T-even TMD PDF $_{[+]}$ = TMD PDF $_{[-]}$
T-odd TMD PDF $_{[+]}$ = -TMD PDF $_{[-]}$

← breaking universality!
(but in a calculable way)



Process dependence



Sivers

$$f_{1T}^{\perp[+]} = -f_{1T}^{\perp[-]}$$

Boer-Mulders

$$h_1^{\perp[+]} = -h_1^{\perp[-]}$$

SIDIS

Drell-Yan

Prediction of QCD based on interplay between time-reversal and (color) gauge symmetry



Intense experimental work to test this prediction

Intuition: in SIDIS, gauge link $U_{[+]}$ describes color final-state interactions between

struck parton and spectators

in Drell-Yan, gauge link $U_{[-]}$ describes color initial-state interactions between


struck parton and spectators





Gluon TMDs are phenomenologically unknown.
Why ?

- gluons carry no electric charge \rightarrow in SIDIS they appear only at higher orders
- gluons carries “two color charges” \rightarrow in general, difficult to neutralise them all
- in hadronic collisions, gluons appear at tree level, but :
 - factorisation theorem available only for Drell-Yan processes
 - for $H_1+H_2 \rightarrow h+X$ no factor. th. but also no counterexample disproving it
- useful processes under study:



Boer et al., P.R.L. **108** (12) 032002
 den Dunnen et al., P.R.L. **112** (14) 212001
 Mukherjee & Rajesh, arXiv:1609.05596
 Boer et al., arXiv:1605.07934
 Godbole et al., arXiv:1703.01991
 D'Alesio et al., arXiv:1705.04169
 Rajesh et al., arXiv:1802.10359
 Zheng et al., arXiv:1805.05290
 Bacchetta et al., arXiv:1809.02056
 D'Alesio et al., arXiv:1908.00446
 D'Alesio et al., arXiv:1910.09640

gluon TMDs

- First classification given in

Mulders & Rodrigues,
P.R. D63 (01) 094021, arXiv:hep-ph/0009343

- Factorization, evolution & universality
 studied in

Ji et al., JHEP 07 (05) 020, arXiv:hep-ph/0503015
Buffing et al., P.R. D88 (13) 054027, arXiv:1306.5897
Boer & Van Dunnen, N.P. B886 (14) 421, arXiv:1404.6753
Echevarria et al., JHEP 07 (15) 158 [E: 05 (17) 073], arXiv:1502.05354

polarization
 gluon • ↻ ↑

nucleon

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	f_1^g	×	$h_1^{\perp,g}$
	L	×	g_1^g	$h_{1L}^{\perp,g}$
	T	$f_{1T}^{\perp,g}$	g_{1T}^g	$h_1^g, h_{1T}^{\perp,g}$

T-odd TMDs

gluon TMDs

- First classification given in

Mulders & Rodrigues, P.R. D63 (01) 094021, arXiv:hep-ph/0009343

- Factorization, evolution & universality studied in

Ji et al., JHEP 07 (05) 020, arXiv:hep-ph/0503015
Buffing et al., P.R. D88 (13) 054027, arXiv:1306.5897
Boer & Van Dunnen, N.P. B886 (14) 421, arXiv:1404.6753
Echevarria et al., JHEP 07 (15) 158 [E: 05 (17) 073], arXiv:1502.05354


polarization gluon • ↻ ↑

nucleon


		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	f_1^g	✗	$h_1^{\perp,g}$
	L	✗	g_1^g	$h_{1L}^{\perp,g}$
	T	$f_{1T}^{\perp,g}$	g_{1T}^g	$h_1^g, h_{1T}^{\perp,g}$

T-odd TMDs


gluons carry “two color charges” → intricate non-universality




two-jets SIDIS



color structure [+,+]



gluon fusion to Higgs [-,-]



T-even: $f_1^{[+,+]} = f_1^{[-,-]}$
 T-odd: $f_{1T}^{\perp[+,+]} = -f_{1T}^{\perp[-,-]}$

gluon TMDs

- First classification given in

Mulders & Rodrigues, P.R. D63 (01) 094021, arXiv:hep-ph/0009343

- Factorization, evolution & universality studied in

Ji et al., JHEP 07 (05) 020, arXiv:hep-ph/0503015
Buffing et al., P.R. D88 (13) 054027, arXiv:1306.5897
Boer & Van Dunnen, N.P. B886 (14) 421, arXiv:1404.6753
Echevarria et al., JHEP 07 (15) 158 [E: 05 (17) 073], arXiv:1502.05354


polarization gluon • ↻ ↑

nucleon


		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	f_1^g	✗	$h_1^{\perp,g}$
	L	✗	g_1^g	$h_{1L}^{\perp,g}$
	T	$f_{1T}^{\perp,g}$	g_{1T}^g	$h_1^g, h_{1T}^{\perp,g}$

T-odd TMDs


gluons carry “two color charges” → intricate non-universality




two-jets SIDIS



color structure [+,+]



gluon fusion to Higgs [-,-]



T-even: $f_1^{[+,+]} = f_1^{[-,-]}$

T-odd: $f_{1T}^{\perp [+,+]} = -f_{1T}^{\perp [-,-]}$

WW-type TMDs

gluon TMDs

- First classification given in

Mulders & Rodrigues, P.R. D63 (01) 094021, arXiv:hep-ph/0009343

- Factorization, evolution & universality studied in


Ji et al., JHEP 07 (05) 020, arXiv:hep-ph/0503015
Buffing et al., P.R. D88 (13) 054027, arXiv:1306.5897
Boer & Van Dunnen, N.P. B886 (14) 421, arXiv:1404.6753
Echevarria et al., JHEP 07 (15) 158 [E: 05 (17) 073], arXiv:1502.05354

polarization gluon • ↻ ↑


		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	f_1^g	✗	$h_1^{\perp,g}$
	L	✗	g_1^g	$h_{1L}^{\perp,g}$
	T	$f_{1T}^{\perp,g}$	g_{1T}^g	$h_1^g, h_{1T}^{\perp,g}$

T-odd TMDs


gluons carry “two color charges” → intricate non-universality




two-jets SIDIS



color structure [+,+]




gluon fusion to Higgs [-,-]




T-even: $f_1^{[+,+]} = f_1^{[-,-]}$
 T-odd: $f_{1T}^{\perp[+,+]} = -f_{1T}^{\perp[-,-]}$


WW-type TMDs



$p p \rightarrow \gamma^* + \text{jet}$



[+,-]



[-,+]

T-even: $f_1^{[+,-]} = f_1^{[-,+]}$
 T-odd: $f_{1T}^{\perp[+,-]} = -f_{1T}^{\perp[-,+]}$

gluon TMDs

- First classification given in

Mulders & Rodrigues, P.R. D63 (01) 094021, arXiv:hep-ph/0009343

- Factorization, evolution & universality studied in

Ji et al., JHEP 07 (05) 020, arXiv:hep-ph/0503015
Buffing et al., P.R. D88 (13) 054027, arXiv:1306.5897
Boer & Van Dunnen, N.P. B886 (14) 421, arXiv:1404.6753
Echevarria et al., JHEP 07 (15) 158 [E: 05 (17) 073], arXiv:1502.05354


polarization gluon • ↻ ↑

nucleon


		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	f_1^g	✗	$h_1^{\perp,g}$
	L	✗	g_1^g	$h_{1L}^{\perp,g}$
	T	$f_{1T}^{\perp,g}$	g_{1T}^g	$h_1^g, h_{1T}^{\perp,g}$

T-odd TMDs


gluons carry “two color charges” → intricate non-universality




two-jets SIDIS



color structure [+,+]




gluon fusion to Higgs [-,-]




T-even: $f_1^{[+,+]} = f_1^{[-,-]}$

T-odd: $f_{1T}^{\perp[+,+]} = -f_{1T}^{\perp[-,-]}$


WW-type TMDs



$p p \rightarrow \gamma^* + \text{jet}$



[+,-]



[-,+]

T-even: $f_1^{[+,-]} = f_1^{[-,+]}$

T-odd: $f_{1T}^{\perp[+,-]} = -f_{1T}^{\perp[-,+]}$

dipole-type TMDs

gluon TMDs

many papers exploring useful channels at colliders to extract WW and dipole gluon TMDs.

Handy pocket list:

Boer, talk at IWHSS 2020

(see also recent review on quarkonium physics)

Boer et al., arXiv:2409.03691

$f_1^g^{[+,+]}$	$pp \rightarrow \gamma J/\psi X$	LHC
	$pp \rightarrow \gamma \Upsilon X$	LHC
$f_1^g^{[+,-]}$	$pp \rightarrow \gamma \text{jet} X$	LHC & RHIC
$h_1^{\perp g [+,+]}$	$e p \rightarrow e' Q \bar{Q} X$	EIC
	$e p \rightarrow e' \text{jet jet} X$	EIC
	$pp \rightarrow \eta_{c,b} X$	LHC & NICA
	$pp \rightarrow H X$	LHC
$h_1^{\perp g [+,-]}$	$pp \rightarrow \gamma^* \text{jet} X$	LHC & RHIC
$f_{1T}^{\perp g [+,+]}$	$e p^\uparrow \rightarrow e' Q \bar{Q} X$	EIC
	$e p^\uparrow \rightarrow e' \text{jet jet} X$	EIC
$f_{1T}^{\perp g [-,-]}$	$p^\uparrow p \rightarrow \gamma \gamma X$	RHIC
$f_{1T}^{\perp g [+,-]}$	$p^\uparrow A \rightarrow \gamma^{(*)} \text{jet} X$	RHIC
	$p^\uparrow A \rightarrow h X \ (x_F < 0)$	RHIC & NICA

gluon TMDs

many papers exploring useful channels at colliders to extract WW and dipole gluon TMDs.

Handy pocket list:

Boer, talk at IWHSS 2020

(see also recent review on quarkonium physics)

Boer et al., arXiv:2409.03691

$f_1^g[+,+]$	$pp \rightarrow \gamma J/\psi X$	LHC
	$pp \rightarrow \gamma \Upsilon X$	LHC
$f_1^g[+,-]$	$pp \rightarrow \gamma \text{jet} X$	LHC & RHIC
$h_1^{\perp g}[+,+]$	$ep \rightarrow e' Q \bar{Q} X$	EIC
	$ep \rightarrow e' \text{jet jet} X$	EIC
	$pp \rightarrow \eta_{c,b} X$	LHC & NICA
	$pp \rightarrow H X$	LHC
$h_1^{\perp g}[+,-]$	$pp \rightarrow \gamma^* \text{jet} X$	LHC & RHIC
$f_{1T}^{\perp g}[+,+]$	$ep^\uparrow \rightarrow e' Q \bar{Q} X$	EIC
	$ep^\uparrow \rightarrow e' \text{jet jet} X$	EIC
$f_{1T}^{\perp g}[-,-]$	$p^\uparrow p \rightarrow \gamma \gamma X$	RHIC
$f_{1T}^{\perp g}[+,-]$	$p^\uparrow A \rightarrow \gamma^{(*)} \text{jet} X$	RHIC
	$p^\uparrow A \rightarrow h X (x_F < 0)$	RHIC & NICA

- TMD factorization $\xrightarrow{\text{small } x}$ UGD k_t factorization

WW $f_1^g[+,+]$ \longrightarrow # density of gluons in CGC

dipole $f_1^g[+,-]$ \longrightarrow Fourier Transform of color-dipole cross section in CGC

Dominguez et al., P.R.L. 106 (11) 022301, arXiv:1009.2141

Dominguez et al., P.R. D83 (11) 105005, arXiv:1101.0715

gluon TMDs

many papers exploring useful channels at colliders to extract WW and dipole gluon TMDs.

Handy pocket list:

Boer, talk at IWHSS 2020

(see also recent review on quarkonium physics)

Boer et al., arXiv:2409.03691

$f_1^g[+,+]$	$pp \rightarrow \gamma J/\psi X$	LHC
	$pp \rightarrow \gamma \Upsilon X$	LHC
$f_1^g[+,-]$	$pp \rightarrow \gamma \text{jet} X$	LHC & RHIC
$h_1^{\perp g}[+,+]$	$e p \rightarrow e' Q \bar{Q} X$	EIC
	$e p \rightarrow e' \text{jet jet} X$	EIC
	$pp \rightarrow \eta_{c,b} X$	LHC & NICA
	$pp \rightarrow H X$	LHC
$h_1^{\perp g}[+,-]$	$pp \rightarrow \gamma^* \text{jet} X$	LHC & RHIC
$f_{1T}^{\perp g}[+,+]$	$e p^\uparrow \rightarrow e' Q \bar{Q} X$	EIC
	$e p^\uparrow \rightarrow e' \text{jet jet} X$	EIC
$f_{1T}^{\perp g}[-,-]$	$p^\uparrow p \rightarrow \gamma \gamma X$	RHIC
$f_{1T}^{\perp g}[+,-]$	$p^\uparrow A \rightarrow \gamma^{(*)} \text{jet} X$	RHIC
	$p^\uparrow A \rightarrow h X (x_F < 0)$	RHIC & NICA

- TMD factorization $\xrightarrow{\text{small } x}$ UGD k_t factorization

WW $f_1^g[+,+]$ \longrightarrow # density of gluons in CGC

dipole $f_1^g[+,-]$ \longrightarrow Fourier Transform of color-dipole cross section in CGC

Dominguez et al., P.R.L. 106 (11) 022301, arXiv:1009.2141

Dominguez et al., P.R. D83 (11) 105005, arXiv:1101.0715

- small-x limit of T-odd gluon TMDs:

WW $f_{1T}^\perp, h_1, h_{1T}^\perp \rightarrow 0$

spin-dependent T-odd part of dipole amplitude
describes the colorless C-odd t-channel 3-gluon exchange

dipole $x f_{1T}^\perp = x h_1 = x h_{1T}^\perp \rightarrow -\frac{k_T^2 N_c}{4\pi\alpha_s} O_{1T}^\perp(x, k_T^2)$ spin Odderon

Boer et al., P.R.L. 116 (16) 122001, arXiv:1511.03485

gluon TMDs : only models

- Available experimental information on gluon TMDs is scarce.
- Very few attempts of phenomenological studies:

*Lansberg et al., P.L. **B784** (18) 217 [E: P.L. **B791** (19) 420], arXiv:1710.01684
D'Alesio et al., P.R. **D96** (17) 036011, arXiv:1705.04169
D'Alesio et al., P.R. **D99** (19) 036013, arXiv:1811.02970
D'Alesio et al., P.R. **D102** (20) 094011, arXiv:2007.03353*

- Many models on the market (list of references too long).



Let me advertise our one, **for first time providing systematically all T-even and T-odd gluon TMDs at leading twist:**

*Bacchetta et al., E.P.J.C **80** (20) 733, arXiv:2005.02288* T-even

*Bacchetta et al., E.P.J.C **84** (24) 576, arXiv:2402.17556* T-odd




spectator model of gluon TMDs

- Nucleon = gluon + spectator on-shell spin-1/2 particle




- T-odd generated by gluon-spectator FSI via 1 gluon-exchange
- Spectator mass takes continuous range of values through a parametric spectral function
- Parameters fixed by reproducing collinear gluon PDFs f_1 and g_1 from NNPDF3.0

Bacchetta et al., *E.P.J.C* **80** (20) 733, arXiv:2005.02288
 Bacchetta et al., *E.P.J.C* **84** (24) 576, arXiv:2402.17556





More on factorisation \rightarrow evolution



inclusive DIS: QCD corrections generate soft and collinear divergences

sum of real and virtual diagrams cancel soft divergences


collinear divergences reabsorbed in collinear PDFs

factorisation scale μ determines what is perturbative (calculable) from what is non perturbative (inside PDFs)

\rightarrow scale dependence given by DGLAP evolution eq.'s



More on factorisation \rightarrow evolution




inclusive DIS: QCD corrections generate soft and collinear divergences

sum of real and virtual diagrams cancel soft divergences

collinear divergences reabsorbed in collinear PDFs

factorisation scale μ determines what is perturbative (calculable) from what is non perturbative (inside PDFs)

\rightarrow scale dependence given by DGLAP evolution eq.'s




SIDIS: soft divergences do not cancel anymore

new class of light-cone (rapidity) divergences

need to introduce a **soft factor** convoluted with TMD PDFs and FFs



More on factorisation → evolution




inclusive DIS: QCD corrections generate soft and collinear divergences

sum of real and virtual diagrams cancel soft divergences

collinear divergences reabsorbed in collinear PDFs

factorisation scale μ determines what is perturbative (calculable) from what is non perturbative (inside PDFs)

→ scale dependence given by DGLAP evolution eq.'s



SIDIS: soft divergences do not cancel anymore

new class of light-cone (rapidity) divergences

need to introduce a **soft factor** convoluted with TMD PDFs and FFs

need to introduce a new **“rapidity scale”** ζ that regulates the rapidity divergences and splits the soft factor content between TMD PDFs and FFs → new scale dependence

$$\text{DGLAP eq.'s} \quad \frac{d \log \text{TMD}}{d \log \mu} = \gamma_D(\mu, \zeta)$$

$$\text{CSS eq.'s} \quad \frac{d \log \text{TMD}}{d \log \sqrt{\zeta}} = K(\mu)$$



TMDs in position space



TMD evolution from initial (μ_0, ζ_0) scales is better studied in position space $b_T (\leftrightarrow P_{hT})$

In fact, by Fourier transforming the complicate convolution between internal transverse momenta gets broken

$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{C} \left[\text{Tr} \left[\Phi(x_B, \mathbf{k}_\perp, S) \gamma^\mu \Delta(z_h, \mathbf{P}_\perp) \gamma^\nu \right] \right]$$
$$\mathcal{C} [\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]$$
$$\int d^2\mathbf{b}_T e^{-i\mathbf{b}_T \cdot \mathbf{P}_{hT}} \dots \int d\mathbf{P}_\perp e^{i\mathbf{b}_T \cdot \mathbf{P}_\perp} \dots \int d\mathbf{k}_\perp e^{i\mathbf{b}_T \cdot \mathbf{k}_\perp} \dots$$



TMDs in position space



TMD evolution from initial (μ_0, ζ_0) scales is better studied in position space $b_T (\leftrightarrow P_{hT})$

In fact, by Fourier transforming the complicate convolution between internal transverse momenta gets broken

$$2MW^{\mu\nu}(q, P, S, P_h) = 2z_h \mathcal{C} \left[\text{Tr} \left[\Phi(x_B, \mathbf{k}_\perp, S) \gamma^\mu \Delta(z_h, \mathbf{P}_\perp) \gamma^\nu \right] \right]$$



$$\mathcal{C}[\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots] \\ \int d^2\mathbf{b}_T e^{-i\mathbf{b}_T \cdot \mathbf{P}_{hT}} \dots \int d\mathbf{P}_\perp e^{i\mathbf{b}_T \cdot \mathbf{P}_\perp} \dots \int d\mathbf{k}_\perp e^{i\mathbf{b}_T \cdot \mathbf{k}_\perp} \dots$$


$$\frac{d\sigma}{dx dz dq_T dQ} \sim \mathcal{H}^{\text{SIDIS}}(Q^2) \frac{1}{2\pi} \int_0^\infty db_T b_T J_0(b_T, q_T) \tilde{f}_1^q(x, b_T^2; Q^2) \tilde{D}_1^{q \rightarrow h}(z, b_T^2; Q^2)$$

hard part

TMDPDF

TMDFF

$$q_T^2 = \frac{P_{hT}^2}{z^2} \ll Q^2$$






More on factorisation \rightarrow evolution



For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid

$$f_1^q(x, b_T^2; \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_0, \zeta_0) \right] f_1^q(x, b_T^2; \mu_0, \zeta_0)$$






For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid

$$f_1^q(x, b_T^2; \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_0, \zeta_0) \right] f_1^q(x, b_T^2; \mu_0, \zeta_0)$$

DGLAP+CSS eqs. ↓

$$\exp \left[\int_{\mu_0}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_0) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right]$$





More on factorisation → evolution



For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid


$$f_1^q(x, b_T^2; \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_0, \zeta_0) \right]$$

DGLAP+CSS eqs. ↓
$$\exp \left[\int_{\mu_0}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_0) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right]$$

$$f_1^q(x, b_T^2; \mu_0, \zeta_0)$$

OPE on PDFs ↓
$$= \sum_i [C_{q \rightarrow i}(x, b_T^2; \mu_0, \zeta_0) \otimes f_1^i(x, \mu_0)]$$

small b_T (large k_T) from perturbative splitting





More on factorisation → evolution




For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid

$$f_1^q(x, b_T^2; \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_0, \zeta_0) \right] f_1^q(x, b_T^2; \mu_0, \zeta_0)$$

DGLAP+CSS eqs. ↓
↓ OPE on PDFs

$$\exp \left[\int_{\mu_0}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_0) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right] = \sum_i [C_{q \rightarrow i}(x, b_T^2; \mu_0, \zeta_0) \otimes f_1^i(x, \mu_0)]$$

For large b_T perturbation theory breaks down; need to find a suitable function that smoothly connects the two regions





More on factorisation → evolution



For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid

$$f_1^q(x, b_T^2; \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_0, \zeta_0) \right] f_1^q(x, b_T^2; \mu_0, \zeta_0)$$

DGLAP+CSS eqs. ↓

$$\exp \left[\int_{\mu_0}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_0) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right]$$

OPE on PDFs ↓


$$= \sum_i [C_{q \rightarrow i}(x, b_T^2; \mu_0, \zeta_0) \otimes f_1^i(x, \mu_0)]$$

For large b_T perturbation theory breaks down; need to find a suitable function that smoothly connects the two regions

$\overline{\text{MS}}$ factorization scheme suggests the following scale:

$$\mu_0 = \sqrt{\zeta_0} = \mu_b = \frac{2e^{-\gamma_E}}{b^*(b_T)}$$

$$b_{min} = \frac{2e^{-\gamma_E}}{Q} \leq b^*(b_T) \leq b_{max} = 2e^{-\gamma_E}$$





More on factorisation → evolution



For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid

$$f_1^q(x, b_T^2; \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_0, \zeta_0) \right] f_1^q(x, b_T^2; \mu_0, \zeta_0)$$

DGLAP+CSS eqs. ↓
OPE on PDFs ↓


$$\exp \left[\int_{\mu_0}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_0) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right] = \sum_i [C_{q \rightarrow i}(x, b_T^2; \mu_0, \zeta_0) \otimes f_1^i(x, \mu_0)]$$



$$f_1^q(x, b_T^2; \mu, \zeta) = \frac{f_1^q(x, b_T^2; \mu, \zeta)}{f_1^q(x, b^*(b_T); \mu, \zeta)} f_1^q(x, b^*(b_T); \mu, \zeta)$$

$$\tilde{f}_{NP}(x, b_T^2; Q_0^2)$$

Q_0 = scale at which the nonperturbative term is parametrised





More on factorisation → evolution



For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid


$$f_1^q(x, b^*(b_T); \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_b, \zeta_0) \right] f_1^q(x, b^*(b_T); \mu_b, \zeta_0) \times F_{NP}(b_T; Q_0^2)$$

$$\begin{aligned} & \text{DGLAP+CSS eqs.} \downarrow \\ & \exp \left[\int_{\mu_b}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_b) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right] \\ & \text{OPE on PDFs} \downarrow \\ & = \sum_i [C_{q \rightarrow i}(x, b^*(b_T); \mu_b, \zeta_0) \otimes f_1^i(x, \mu_b)] \end{aligned}$$

$$\mu_b = \frac{2e^{-\gamma_E}}{b_*(b_T)}$$

$$K \rightarrow K + g_{NP}(b_T)$$

conventional choice: $\mu = \sqrt{\zeta} = Q$ $\mu_0 = \sqrt{\zeta_0} = \mu_b = \frac{2e^{-\gamma_E}}{b_*(b_T)}$





For $b_T \ll 1/\Lambda_{QCD}$ perturbation theory is valid

$$f_1^q(x, b^*(b_T); \mu, \zeta) = \text{Evo} \left[(\mu, \zeta) \leftarrow (\mu_b, \zeta_0) \right] f_1^q(x, b^*(b_T); \mu_b, \zeta_0) \quad \times \quad F_{NP}(b_T; Q_0^2)$$

$$\begin{array}{l} \text{DGLAP+CSS eqs.} \downarrow \\ \exp \left[\int_{\mu_b}^{\mu} \frac{d\mu'}{\mu'} \gamma_D(\mu, \zeta) + K(\mu_b) \log \frac{\sqrt{\zeta}}{\sqrt{\zeta_0}} \right] \\ \text{OPE on PDFs} \swarrow \\ = \sum_i [C_{q \rightarrow i}(x, b^*(b_T); \mu_b, \zeta_0) \otimes f_1^i(x, \mu_b)] \end{array}$$

$$\mu_b = \frac{2e^{-\gamma_E}}{b^*(b_T)}$$

$$K \rightarrow K + g_{NP}(b_T)$$

Final formula

$$f_1^q(x, b^*; Q^2) = \exp \left[\int_{\mu_b}^Q \frac{d\mu'}{\mu'} \gamma_D(Q) + K(\mu_b) \log \left(\frac{Q}{\mu_b} \right) + g_{NP}(b_T) \log \left(\frac{Q}{Q_0} \right) \right] \sum_i [C_{q \rightarrow i} \otimes f_1^i](x, b^*, \mu_b) F_{NP}(b_T, Q_0^2)$$

Collins, Soper, Sterman, N.P. B250 (85)
Collins, "Foundations of Perturbative QCD" (2011)
Rogers and Aybat, P.R. D83 (11)



others schemes possible:

Laenen, Sterman Vogelsang, P.R.L. 84 (00)

Bozzi et al., N.P. B737 (06)

Echevarria et al., E.P.J. C73 (13) ...

CSS evolution formula for TMD

$$f_1^q(x, b^*; Q^2) = \exp \left[\int_{\mu_b}^Q \frac{d\mu'}{\mu'} \gamma_D(Q) + K(\mu_b) \log \left(\frac{Q}{\mu_b} \right) + g_{NP}(b_T) \log \left(\frac{Q}{Q_0} \right) \right] \sum_i [C_{q \rightarrow i} \otimes f_1^i](x, b^*, \mu_b) F_{NP}(b_T, Q_0^2)$$
$$\mu_b = \frac{2e^{-\gamma_E}}{b^*(b_T)}$$



others schemes possible:

Laenen, Sterman Vogelsang, P.R.L. 84 (00)

Bozzi et al., N.P. B737 (06)

Echevarria et al., E.P.J. C73 (13) ...

CSS evolution formula for TMD

$$f_1^q(x, b^*; Q^2) = \exp \left[\int_{\mu_b}^Q \frac{d\mu'}{\mu'} \gamma_D(Q) + K(\mu_b) \log \left(\frac{Q}{\mu_b} \right) + g_{NP}(b_T) \log \left(\frac{Q}{Q_0} \right) \right] \sum_i [C_{q \rightarrow i} \otimes f_1^i](x, b^*, \mu_b, F_{NP}(b_T, Q_0^2))$$

$$\mu_b = \frac{2e^{-\gamma_E}}{b^*(b_T)}$$

arbitrariness of nonperturbative components

- choice of $b^*(b_T)$ functional form
- choice of $g_{NP}(b_T)$ functional form
- choice of $F_{NP}(b_T, Q_0)$ functional form

each one affects evolution: how k_{\perp} -distribution changes with scale

→ source of theoretical bias/uncertainty

need to be constrained by experimental data with large lever arm in Q^2

EIC is the suitable machine for that



Quality parameters of TMD extraction



$$\frac{d\sigma}{dx dz dq_T dQ} \sim \mathcal{H}^{\text{SIDIS}}(Q^2) \frac{1}{2\pi} \int_0^\infty db_T b_T J_0(b_T, q_T) \tilde{f}_1^q(x, b^*(b_T); Q^2) \tilde{D}_1^{q \rightarrow h}(z, b^*(b_T); Q^2)$$

$$f_1^q(x, b^*; Q^2) = \exp \left[\int_{\mu_b}^Q \frac{d\mu'}{\mu'} \gamma_D(Q) + K(\mu_b) \log \left(\frac{Q}{\mu_b} \right) + g_{NP}(b_T) \log \left(\frac{Q}{Q_0} \right) \right] \sum_i [C_{q \rightarrow i} \otimes f_1^i](x, b^*, \mu_b) F_{NP}(b_T, Q^2)$$

$\gamma_D = \gamma_F - \gamma_K \log(\sqrt{\zeta}/\mu) \quad \frac{dK}{d \log \mu} = -\gamma_K \quad \text{cusp anomalous dimension}$



Quality parameters of TMD extraction



$$\frac{d\sigma}{dx dz dq_T dQ} \sim \mathcal{H}^{\text{SIDIS}}(Q^2) \frac{1}{2\pi} \int_0^\infty db_T b_T J_0(b_T, q_T) \tilde{f}_1^q(x, b^*(b_T); Q^2) \tilde{D}_1^{q \rightarrow h}(z, b^*(b_T); Q^2)$$

$$f_1^q(x, b^*; Q^2) = \exp \left[\int_{\mu_b}^Q \frac{d\mu'}{\mu'} \gamma_D(Q) + K(\mu_b) \log \left(\frac{Q}{\mu_b} \right) + g_{NP}(b_T) \log \left(\frac{Q}{Q_0} \right) \right] \sum_i [C_{q \rightarrow i} \otimes f_1^i](x, b^*, \mu_b) F_{NP}(b_T, Q_0^2)$$


$$\gamma_D = \gamma_F - \gamma_K \log(\sqrt{\zeta}/\mu) \quad \frac{dK}{d \log \mu} = -\gamma_K \quad \text{cusp anomalous dimension}$$

perturbative accuracy	α_S^n					
	\mathcal{H} and C	K and γ_F	γ_K	PDF and α_S evol.	FF	
LL	0	-	1	-	-	
NLL	0	1	2	LO	LO	
NLL'	1	1	2	NLO	NLO	
NNLL	1	2	3	NLO	NLO	
NNLL'	2	2	3	NNLO	NNLO	
N ³ LL(-)	2	3	4	NNLO	NLO	
N ³ LL	2	3	4	NNLO	NNLO	

nonperturbative accuracy: quality of the fit from χ^2 value e number of data points



- **Where to find TMDs**






link TMD \leftrightarrow structure functions



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \searrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \uparrow - \downarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \searrow$






link TMD \leftrightarrow structure functions



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
on	u	D_1		H_1^\perp -

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 =$		$h_1^\perp =$ -
	L		$g_1 =$ -	$h_{1L}^\perp =$ -
	T	$f_{1T}^\perp =$ -	$g_{1T} =$ -	$h_1 =$ - $h_{1T}^\perp =$ -



each structure function \sim

$$F \sim d\hat{\sigma}(Q^2) \mathcal{E}[\text{TMDPDF}(x, \mathbf{k}_\perp^2), \text{TMDFF}(z, \mathbf{P}_\perp^2)]$$

$$\mathcal{E}[\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]$$



link TMD \leftrightarrow structure functions



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
on	U	D_1		H_1^\perp -

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	f_1		h_1^\perp -
	L		g_1 -	h_{1L}^\perp -
	T	f_{1T}^\perp -	g_{1T} -	h_1 - h_{1T}^\perp -

target polariz. $\frac{d\sigma}{dx dy dz d\phi_h dP_{hT}^2} \sim$

$$\begin{aligned}
 & \text{○} \quad A(y) F_U + B(y) \cos 2\phi_h F_U^{\cos 2\phi_h} \\
 & + C(y) F_{LL} + B(y) \sin 2\phi_h F_L^{\sin 2\phi_h} \\
 & + A(y) \sin(\phi_h - \phi_S) F_T^{\sin(\phi_h - \phi_S)} \\
 & + B(y) \sin(\phi_h + \phi_S) F_T^{\sin(\phi_h + \phi_S)} \\
 & + B(y) \sin(3\phi_h - \phi_S) F_T^{\sin(3\phi_h - \phi_S)} \\
 & + C(y) \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)}
 \end{aligned}$$

each structure function \sim

$$\begin{aligned}
 F & \sim d\hat{\sigma}(Q^2) \mathcal{E}[\text{TMDPDF}(x, \mathbf{k}_\perp^2), \text{TMDFF}(z, \mathbf{P}_\perp^2)] \\
 \mathcal{E}[\dots] & = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]
 \end{aligned}$$



link TMD \leftrightarrow structure functions



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
on	U	D_1		H_1^\perp -

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 =$		$h_1^\perp =$ -
	L		$g_1 =$ -	$h_{1L}^\perp =$ -
	T	$f_{1T}^\perp =$ -	$g_{1T} =$ -	$h_1 =$ - $h_{1T}^\perp =$ -

target polariz.

$$\frac{d\sigma}{dx dy dz d\phi_h dP_{hT}^2} \sim$$

$$A(y) F_U + B(y) \cos 2\phi_h F_U^{\cos 2\phi_h}$$

$$\text{pink circle} \rightarrow + C(y) F_{LL} + B(y) \sin 2\phi_h F_L^{\sin 2\phi_h}$$

$$+ A(y) \sin(\phi_h - \phi_S) F_T^{\sin(\phi_h - \phi_S)}$$

$$+ B(y) \sin(\phi_h + \phi_S) F_T^{\sin(\phi_h + \phi_S)}$$

$$+ B(y) \sin(3\phi_h - \phi_S) F_T^{\sin(3\phi_h - \phi_S)}$$

$$+ C(y) \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)}$$

each structure function \sim

$$F \sim d\hat{\sigma}(Q^2) \mathcal{E}[\text{TMDPDF}(x, \mathbf{k}_\perp^2), \text{TMDFF}(z, \mathbf{P}_\perp^2)]$$

$$\mathcal{E}[\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]$$



link TMD ↔ structure functions



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
on	U	D_1		H_1^\perp -

		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 =$		$h_1^\perp =$ -
	L		$g_1 =$ -	$h_{1L}^\perp =$ -
	T	$f_{1T}^\perp =$ -	$g_{1T} =$ -	$h_1 =$ - $h_{1T}^\perp =$ -

target polariz.

$$\frac{d\sigma}{dx dy dz d\phi_h dP_{hT}^2} \sim$$

$$\begin{aligned}
& A(y) F_U + B(y) \cos 2\phi_h F_U^{\cos 2\phi_h} \\
& + C(y) F_{LL} + B(y) \sin 2\phi_h F_L^{\sin 2\phi_h} \\
& + A(y) \sin(\phi_h - \phi_S) F_T^{\sin(\phi_h - \phi_S)} \\
& + B(y) \sin(\phi_h + \phi_S) F_T^{\sin(\phi_h + \phi_S)} \\
& + B(y) \sin(3\phi_h - \phi_S) F_T^{\sin(3\phi_h - \phi_S)} \\
& + C(y) \cos(\phi_h - \phi_S) F_{LT}^{\cos(\phi_h - \phi_S)}
\end{aligned}$$




each structure function ~

$$F \sim d\hat{\sigma}(Q^2) \mathcal{E}[\text{TMDPDF}(x, \mathbf{k}_\perp^2), \text{TMDFF}(z, \mathbf{P}_\perp^2)]$$

$$\mathcal{E}[\dots] = \int d\mathbf{P}_\perp d\mathbf{k}_\perp \delta^{(2)}(z\mathbf{k}_\perp + \mathbf{P}_\perp - \mathbf{P}_{hT}) [\dots]$$



TMDs with jets: SIDIS



$$\mathbf{q}_T = \mathbf{p}_{Te} + \mathbf{p}_{Tj} \ll \mathbf{p}_T = \frac{\mathbf{p}_{Te} - \mathbf{p}_{Tj}}{2}$$

$$F_{UU} \sim H(Q) J(p_T R, Q) \{f_1(x, q_T, Q)\}$$

hard jet "dressed" TMD

similarly for other $F_{..}$

"familiar" expression

$$\frac{d\sigma}{dy_j d\mathbf{p}_T d\mathbf{q}_T} = F_{UU} +$$


$$\lambda_e S_L F_{LL}^{g_{1L}}$$

$$+ S_T \sin(\phi_j - \phi_S) F_{UT}^{f_{1T}^\perp \sin(\phi_j - \phi_S)} + \lambda_e S_T \cos(\phi_j - \phi_S) F_{LT}^{g_{1T}^\perp \cos(\phi_j - \phi_S)}$$

Kang et al., arXiv:2106.15624



TMDs with jets: SIDIS



$$\mathbf{q}_T = \mathbf{p}_{Te} + \mathbf{p}_{Tj} \ll \mathbf{p}_T = \frac{\mathbf{p}_{Te} - \mathbf{p}_{Tj}}{2}$$

$$F_{UU} \sim H(Q) \text{TMDJFF}(z_h, p_T R, Q) \{f_1(x, q_T, Q)\}$$

hard jet "dressed" TMD

similarly for other $F_{..}$


"familiar" expression

$$\begin{aligned} \frac{d\sigma}{dy_j d\mathbf{p}_T d\mathbf{q}_T} = & f_1^{\perp} \mathcal{D}_1^{\perp} h_1^{\perp} \mathcal{H}_1^{\perp} g_{1L} \mathcal{D}_1^{\perp} \\ & F_{UU} + \cos(\phi_j - \phi_h) F_{UU}^{\cos(\phi_j - \phi_h)} + \lambda_e S_L F_{LL} \\ & + S_L \sin(\phi_j - \phi_h) F_{UL}^{\sin(\phi_j - \phi_h)} \\ & + S_T \sin(\phi_j - \phi_S) F_{UT}^{\sin(\phi_j - \phi_S)} + \lambda_e S_T \cos(\phi_j - \phi_S) F_{LT}^{\cos(\phi_j - \phi_S)} \\ & + S_T \sin(\phi_h - \phi_S) F_{UT}^{\sin(\phi_h - \phi_S)} + S_T \sin(2\phi_j - \phi_h - \phi_S) F_{UT}^{\sin(2\phi_j - \phi_h - \phi_S)} \end{aligned}$$


Kang et al., arXiv:2106.15624



TMDs with jets: hybrid factorisation



SIDIS



hybrid scheme:

- TMD framework for TMD **fragmentation**
- collinear framework for PDF

Factorization theorem for $j_T \ll Q$
universality for TMD **fragmentation**

*Kang, Liu, Ringer, Xing, JHEP **1711** (17), arXiv:1705.08443*

*Kang, Prokudin, Ringer, Yuan, P.L. **B774** (17), arXiv:1707.00913*



Outline



- Pause



- Phenomenology of TMDs




The unpolarized TMD PDF



polarizations

nucleon



quark



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \searrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \searrow$

$f_1^q(x, \mathbf{k}_\perp^2)$ probability density of finding a quark q with “longitudinal” (along “+” LC direction) fraction x of nucleon momentum, and transverse momentum \mathbf{k}_\perp

Most recent extractions of unpolarized TMD f_1

SIDIS

	Accuracy	HERMES	COMPASS	DY	W / Z production	N of points	χ^2/N_{points}
PV 2017 arXiv:1703.10157	NLL	✓	✓	✓	✓	8059	1.5
SV 2017 arXiv:1706.01473	NNLL'	✗	✗	✓	✓	309	1.23
BSV 2019 arXiv:1902.08474	NNLL'	✗	✗	✓	✓	457	1.17
SV 2019 arXiv:1912.06532	N ³ LL(-)	✓	✓	✓	✓	1039	1.06
PV 2019 arXiv:1912.07550	N ³ LL	✗	✗	✓	✓	353	1.07
SV19 + flavor dep. arXiv:2201.07114	N ³ LL	✗	✗	✓	✓	309	<1.08>
MAPTMD 2022 arXiv:2206.07598	N ³ LL(-)	✓	✓	✓	✓	2031	1.06
ART23 arXiv:2305.07473	N ⁴ LL	✗	✗	✓	✓	627	0.96
MAPTMD 2024 arXiv:2405.13833	N ³ LL	✓	✓	✓	✓	2031	1.08
MAPNN 2025 arXiv:2502.04166	N ³ LL	✗	✗	✓	✓	482	0.97

first use of Neural Networks

Most recent extractions of unpolarized TMD f_1

SIDIS

	Accuracy	HERMES	COMPASS	DY	W / Z production	N of points	χ^2/N_{points}
PV 2017 arXiv:1703.10157	NLL	✓	✓	✓	✓	8059	1.5
SV 2017 arXiv:1706.01473	NNLL'	✗	✗	✓	✓	309	1.23
BSV 2019 arXiv:1902.08474	NNLL'	✗	✗	✓	✓	457	1.17
SV 2019 arXiv:1912.06532	N ³ LL(-)	✓	✓	✓	✓	1039	1.06
PV 2019 arXiv:1912.07550	N ³ LL	✗	✗	✓	✓	353	1.07
SV19 + flavor dep. arXiv:2201.07114	N ³ LL	✗	✗	✓	✓	309	<1.08>
MAPTMD 2022 arXiv:2206.07598	N ³ LL(-)	✓	✓	✓	✓	2031	1.06
ART23 arXiv:2305.07473	N ⁴ LL	✗	✗	✓	✓	627	0.96
MAPTMD 2024 arXiv:2405.13833	N ³ LL	✓	✓	✓	✓	2031	1.08
MAPNN 2025 arXiv:2502.04166	N ³ LL	✗	✗	✓	✓	482	0.97

increasing accuracy & precision

Most recent extractions of unpolarized TMD f_1

SIDIS

	Accuracy	HERMES	COMPASS	DY	W / Z production	N of points	χ^2/N_{points}
 PV 2017 arXiv:1703.10157	NLL	✓	✓	✓	✓	8059	1.5
SV 2017 arXiv:1706.01473	NNLL'	✗	✗	✓	✓	309	1.23
BSV 2019 arXiv:1902.08474	NNLL'	✗	✗	✓	✓	457	1.17
 SV 2019 arXiv:1912.06532	N ³ LL(-)	✓	✓	✓	✓	1039	1.06
PV 2019 arXiv:1912.07550	N ³ LL	✗	✗	✓	✓	353	1.07
SV19 + flavor dep. arXiv:2201.07114	N ³ LL	✗	✗	✓	✓	309	<1.08>
 MAPTMD 2022 arXiv:2206.07598	N ³ LL(-)	✓	✓	✓	✓	2031	1.06
ART23 arXiv:2305.07473	N ⁴ LL	✗	✗	✓	✓	627	0.96
 MAPTMD 2024 arXiv:2405.13833	N ³ LL	✓	✓	✓	✓	2031	1.08
 MAPNN 2025 arXiv:2502.04166	N ³ LL	✗	✗	✓	✓	482	0.97

only four global fits

increasing accuracy & precision

Most recent extractions of unpolarized TMD f_1

SIDIS


	Accuracy	HERMES	COMPASS	DY	W / Z production	N of points	χ^2/N_{points}
PV 2017 arXiv:1703.10157	NLL	✓	✓	✓	✓	8059	1.5
SV 2017 arXiv:1706.01473	NNLL'	✗	✗	✓	✓	309	1.23
BSV 2019 arXiv:1902.08474	NNLL'	✗	✗	✓	✓	457	1.17
SV 2019 arXiv:1912.06532	N ³ LL(-)	✓	✓	✓	✓	1039	1.06
PV 2019 arXiv:1912.07550	N ³ LL	✗	✗	✓	✓	353	1.07
SV19 + flavor dep. arXiv:2201.07114	N ³ LL	✗	✗	✓	✓	309	<1.08>
MAPTMD 2022 arXiv:2206.07598	N ³ LL(-)	✓	✓	✓	✓	2031	1.06
ART23 arXiv:2305.07473	N ⁴ LL	✗	✗	✓	✓	627	0.96
MAPTMD 2024 arXiv:2405.13833	N ³ LL	✓	✓	✓	✓	2031	1.08
MAPNN 2025 arXiv:2502.04166	N ³ LL	✗	✗	✓	✓	482	0.97

MAPTMD24 : introduce **flavor sensitivity of k_T -dependence**

The MAPTMD24 data sets

N_{data} after cuts

Drell Yan	{	233	fixed target
		251	collider
		1547	SIDIS
		<hr/>	
		2031 data points	



kinematic cuts

$$\langle Q \rangle > 1.4 \text{ GeV}$$

$$0.2 < z < 0.7$$

Drell-Yan


$$q_T < 0.2 Q$$

SIDIS

$$P_{hT} < \min \left[\min [0.2 Q, 0.5 Qz] + 0.3 \text{ GeV}, zQ \right]$$

“Normalized” MAPTMD24 TMD PDF

$$\frac{f_1(x, k_T; Q)}{f_1(x, 0; Q)}$$




th. error band =
68% of all replicas

- very different k_T behavior
- it changes with x
- potential impact on the extraction of W mass parameter from collider data

Bacchetta et al., P.L. **B788** (19) 542, arXiv:1807.02101
Bozzi & Signori, Adv.HighEn.Phys. **2019** (19) 2526897, arXiv:1901.01162

Data-driven nonperturbative TMD




at given (x, Q^2) ,
different slopes for different z

at given z ,
different slopes for different x

if we model nonperturbative TMDs
with Gaussians, we need
 x - and z -dependent widths!

MAPTMD22: validity of TMD region?



validity of TMD factorization seems to extend well beyond $P_{hT}/z \ll Q$!

Collins-Soper evolution kernel


universal flavor-independent
drives evolution in rapidity ζ

$$K(b_T, \mu_{b_*}) = K(b_*, \mu_{b_*}) + g_K(b_T)$$

perturbative
(computed)

non-perturbative
(fitted)


[Bermudez Martinez, Vladimirov, arXiv:2206.01105](#)



TMD phenomenology





Lattice QCD

[Avkhadiev, Shanahan, Wagman, Zhao, arXiv:2307.12359](#)




Bacchetta, ePIC 2025 general meeting

The EIC impact at $x=0.01$



$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.01$$

MAPTMD24	2031		
EIC	# pts.	lumi [fb ⁻¹]	
5x41	1273	2.85	
10x100	1611	51.3	
18x275	1648	10	







L. Rossi, Ph.D. Thesis


The EIC impact with 10x100 at $x=0.01$

$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.01$$

	MAPTMD24	2031	# pts.	lumi [fb ⁻¹]
EIC			1611	51.3
10x100				



(simulation campaign of May 2024)





courtesy L. Rossi

The EIC impact with 10x100 at $x=0.01$

$$\frac{\text{TMD}q - \langle \text{TMD}q \rangle}{\langle \text{TMD}q \rangle} \quad x=0.01$$

	MAPTMD24	2031	# pts.	lumi [fb ⁻¹]
EIC			1611	5
10x100				


(early Science conditions)



courtesy L. Rossi

Early Science Conditions

Proposal for EIC Science Program in the First Years



Electron-Ion Collider


ePIC Collaboration Meeting, January 2025

E.C. Aschenauer & R. Ent

17

Early Science Conditions

Proposal for EIC Science Program in the First Years



Electron-Ion Collider

ePIC Collaboration Meeting, January 2025

E.C. Aschenauer & R. Ent

17

Early Science Conditions

ep Luminosity for Phase-1

High Divergence	Lumi per Fill (5 h)	Lumi per Year	Low Divergence	Lumi per Fill (5 h)	Lumi per Year
5 GeV e x 250 GeV p	9.26 pb ⁻¹	6.48 fb ⁻¹	5 GeV e x 250 GeV p	6.81 pb ⁻¹	4.78 fb ⁻¹
10 GeV e x 250 GeV p	13.12 pb ⁻¹	9.18 fb ⁻¹	10 GeV e x 250 GeV p	8.8 pb ⁻¹	6.19 fb ⁻¹
5 GeV e x 130 GeV p	6.3 pb ⁻¹	4.36 fb ⁻¹	5 GeV e x 130 GeV p	5.8 pb ⁻¹	4.1 fb ⁻¹
10 GeV e x 130 GeV p	7.6 pb ⁻¹	5.33 fb ⁻¹	10 GeV e x 130 GeV p	7.1 pb ⁻¹	4.95 fb ⁻¹


Compare to HERA integrated luminosity 1992 – 2007: 0.6 fb⁻¹

Remember:

high divergence: higher lumi, but reduced acceptance for low forward particle p_T^{\min}

low divergence: lower lumi, but increased acceptance for low forward particle p_T^{\min}

→ important for exclusive processes




Electron-Ion Collider

ePIC Collaboration Meeting, January 2025

E.C. Aschenauer & R. Ent


EIC impact in Early Science Conditions



For each (x, Q^2) bin:

- from MAPTMD24, max. uncertainty of $f_{1q}(x, k_T; Q)$ over all k_T and all flavors q
- including EIC pseudodata, color code indicates the flavor with max. reduction in uncertainty over all k_T

EIC impact in Early Science Conditions



For each (x, Q^2) bin:

- from MAPTMD24, max. uncertainty of $f_{1q}(x, k_T; Q)$ over all k_T and all flavors q
- including EIC pseudodata, color code indicates the flavor with max. reduction in uncertainty over all k_T

The EIC impact with 10x130 at $x=0.16$

MAPTMD24 2031

EIC

pts.

lumi [fb⁻¹]

10x130


~1620

5

(early Science conditions)

$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle}$$

$x=0.16, Q=1.77 \text{ GeV}$




courtesy L. Rossi



The Sivers TMD PDF



polarizations
nucleon





quark




		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$
	L		$g_1 = \rightarrow - \leftarrow$	$h_{1L}^\perp = \nearrow - \nwarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$	$g_{1T} = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ $h_{1T}^\perp = \nearrow - \nwarrow$

Bacchetta et al.,
P.L. **B827** (22) 136961,
arXiv:2004.14278



$$\frac{1}{2} \text{Tr}[\Phi \gamma_+] \rightarrow f_1 - f_{1T}^\perp \frac{(\mathbf{k}_\perp \times \mathbf{S}_T) \cdot \hat{\mathbf{P}}}{M}$$



$$\mathbf{S}_T \cdot \mathbf{k}_\perp \times \mathbf{P}$$

Sivers effect: how the momentum distribution of quarks is distorted by the transverse polarization of parent nucleon (“spin-orbit” correlation)

Sivers $f_{1T}^\perp \rightarrow$ indirect access to quark orbital angular momentum

Burkardt, P.R. **D66** (2002) 114005;
N.P. **A735** (2004) 185
Bacchetta & Radici, P.R.L. **107** (2011) 212001
Ji et al., N.P. **B652** (2003) 383


Most recent Sivers extractions

	Framework	SIDIS	DY	W/Z production	forward EM jet	e+e-	N. of points	χ^2/N	
JAM 2020 arXiv:2002.08384	generalized parton model	✓	✓	✓	✗	✓	517	1.04	
PV 2020 arXiv:2004.14278	LO+NLL	✓	✓	✓	✗	✗	125	1.08	
EKT 2020 arXiv:2009.10710	NLO+N ² LL	✓	✓	✓	✗	✗	226/452	0.99 / 1.45	SIDIS / +STAR
BPV 2020 arXiv:2012.05135 arXiv:2103.03270	ζ prescription	✓	✓	✓	✗	✗	76	0.88	
TO-CA 2021 arXiv:2101.03955	generalized parton model	✓	✗	✗	✓	✗	238	$1.05^{+0.03}_{-0.01}$	SIDIS + reweighting
JAM 2022 arXiv:2205.00999	generalized parton model	✓	✓	✓	✗	✗	255	1.10	+ A_N^π data
Fernando-Keller arXiv:2304.14328	generalized parton model	✓	✓	✗	✗	✗	732	1.66	first using Neural Networks

lower accuracy and less data w.r.t. unpolarized TMD

Most recent Sivers extractions


first k_T -moment $f_{1T}^{\perp(1)}(x)$




all parametrizations are in fair agreement for x -dependence of valence flavors

k_T -dependence is still much unconstrained

sea-quarks $\sim O(10^{-3})$ smaller, large errors
 \Rightarrow impact of EIC




Bacchetta et al., P.L. **B827** (22) 136961 arXiv:2004.14278



Sivers extraction using Neural Networks


first k_T -moment $f_{1T}^{\perp(1)}(x)$



Fernando & Keller, *P.R.D***108** (23) 054007 arXiv:2304.14328

but limited analysis: - parton model \Rightarrow no TMD evolution
- no consistent knowledge of unpolarized TMD in denominator of spin asymmetry


Sign change puzzle



π - $p \uparrow$ Drell-Yan



spin asymmetry
 $A_T \sim f_{1,\pi} \otimes f_{1T,p}^\perp$

Aghasyan et al., P.R.L. **119** (17) 112002



sign change

no sign change




p - $p \uparrow \rightarrow W+X$

p - $p \uparrow \rightarrow Z^0+X$

$$A_N \sim f_{1,p} \otimes f_{1T,p}^\perp$$

Adamczyk et al., P.R.L. **116** (16) 132301

still not enough to confirm sign change ?






Transversity



polarizations

nucleon




quark



		Quark polarization		
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow \ominus - \downarrow \ominus$
	L		$g_1 = \rightarrow \ominus - \leftarrow \ominus$	$h_{1L}^\perp = \nearrow \ominus - \nwarrow \ominus$
	T	$f_{1T}^\perp = \uparrow \odot - \downarrow \odot$	$g_{1T} = \uparrow \rightarrow \ominus - \uparrow \leftarrow \ominus$	$h_1 = \uparrow \uparrow \ominus - \uparrow \downarrow \ominus$
				$h_{1T}^\perp = \nearrow \uparrow \ominus - \nwarrow \uparrow \ominus$

- transversity is the prototype of chiral-odd structures
- the only chiral-odd structure that survives in collinear kinematics
- only way to determine the tensor charge $\delta^q(Q^2) = \int_0^1 dx h_1^{q-\bar{q}}(x, Q^2)$

Analyzers of transversity at leading twist



Collins effect


$$\mathbf{S}_T \cdot \mathbf{k} \times \mathbf{P}_{hT}$$

Collins, N.P. **B396** (93) 161

$$\propto h_1(x, k_\perp) \otimes H_1^\perp(z, P_\perp)$$

TMD framework

SIDIS



di-hadron mechanism

$$\mathbf{S}_T \cdot \mathbf{P}_2 \times \mathbf{P}_1 = \mathbf{S}_T \cdot \mathbf{P}_h \times \mathbf{R}_T$$


Collins et al., N.P. **B420** (94)

$$\propto h_1(x) H_1^{\perp 4}(z, R_T^2 \propto M_{h_1 h_2}^2)$$

collinear framework

SIDIS

pp^\uparrow



hadron-in-jet Collins effect

$$j_T^2 \ll Q^2 = (P_T^{jet})^2$$


Yuan, P.R.L. **100** (08)

$$\propto h_1(x) [C(z, \mu) \otimes H_1^\perp(z_h, j_T, P_T^{jet} r)]$$

hybrid framework

SIDIS

pp^\uparrow



Λ spin transfer

Jaffe, P.R. **D54** (96)


$$\propto h_1(x) H_1(z)$$

collinear framework

SIDIS

pp^\uparrow

Analyzers of transversity at leading twist



Collins effect


$$\mathbf{S}_T \cdot \mathbf{k} \times \mathbf{P}_{hT}$$

Collins, N.P. **B396** (93) 161

$$\propto h_1(x, k_\perp) \otimes H_1^\perp(z, P_\perp)$$

TMD framework

SIDIS



di-hadron mechanism

$$\mathbf{S}_T \cdot \mathbf{P}_2 \times \mathbf{P}_1 = \mathbf{S}_T \cdot \mathbf{P}_h \times \mathbf{R}_T$$


Collins et al., N.P. **B420** (94)

$$\propto h_1(x) H_1^{\perp 4}(z, R_T^2 \propto M_{h_1 h_2}^2)$$

collinear framework

SIDIS

$p p^\uparrow$




mechanisms used so far
to extract transversity from data

$$H_1^\perp(z_h, j_T, P_T^{\text{jet}, r})$$

SIDIS

$p p^\uparrow$



Lambda spin transfer


Jaffe, P.R. **D54** (96)

$$\propto h_1(x) H_1(z)$$

collinear framework

SIDIS

$p p^\uparrow$



single-polarised Drell-Yan

Boer, P.R. **D60** (99)

$$\propto h_1^\perp(x_1, k_{\perp 1}) \otimes h_1(x_2, k_{\perp 2})$$

TMD framework


πp^\uparrow

Most recent extractions

Collins effect	Framework	e+e-	SIDIS	Drell-Yan A_N	Lattice
Anselmino 2015 P.R. D92 (15) 114023	parton model	✓	✓	✗	✗
Kang et al. 2016 P.R. D93 (16) 014009	TMD / CSS	✓	✓	✗	✗
Lin et al. 2018 P.R.L. 120 (18) 152502	parton model	✗	✓	✗	✓ g_T
D'Alesio et al. 2020 (CA) P.L. B803 (20) 135347	parton model	✓	✓	✗	✗
JAM3D-20 P.R. D102 (20) 054002	parton model	✓	✓	✓	✗
JAM3D-22 P.R. D106 (22) 034014	parton model	✓	✓	✓	✓ g_T
Bogliione et al. 2024 (TO) P.L. B854 (24) 138712	parton model	✓	✓	✓ reweighting	✗

Dihadron mechanism	e+e- unpol. $d\sigma^0$	e+e- asymmetry	SIDIS	p-p collisions	Lattice
Radici & Bacchetta 2018 P.R.L. 120 (18) 192001	PYTHIA (separately)	✓ (separately)	✓	✓	✗
Benel et al. 2020 E.P.J. C80 (20) 5	PYTHIA (separately)	✓ (separately)	✓	✗	✗
JAMDIFF 2024 P.R.L. 132 (24) 091901	✓	✓	✓	✓	✓ $\delta u, \delta d$

Transversity




D. Pitonyak, QCD Evolution 24

* JAM3D includes $\bar{u} = -\bar{d}$ w.r.t. JAM22

consistency of phenomenological extractions from a variety of exp. data with different approaches
(provided that no LQCD points are included in the fit)

Pheno - lattice : tensor charge

$$g_T = \delta u - \delta d$$





adapted from C. Alexandrou, QCD Evolution 24

green $N_f=2+1+1$

open symbols = no continuum extrapolation

yellow $N_f=2+1$


tension between pheno and lattice ?



adapted from D. Pitonyak, QCD Evolution 24

Pheno - lattice : tensor charge

$$g_T = \delta u - \delta d$$




adapted from C. Alexandrou, QCD Evolution 24

green $N_f=2+1+1$

open symbols = no continuum extrapolation


yellow $N_f=2+1$

tension between pheno and lattice ?



Including lattice data,
JAM finds **compatibility**,
still under discussion...



Experiment	N_{dat}	χ^2_{red}	
		With LQCD	No LQCD
Belle (cross section) [63]	1094	1.01	1.01
Belle (Artru-Collins) [92]	183	0.74	0.73
HERMES [94]	12	1.13	1.10
COMPASS (p) [95]	26	1.24	0.75
COMPASS (D) [95]	26	0.78	0.76
STAR (2015) [96]	24	1.47	1.67
STAR (2018) [64]	106	1.20	1.04
ETMC δu [28]	1	0.71	...
ETMC δd [28]	1	1.02	...
PNDME δu [25]	1	8.68	...
PNDME δd [25]	1	8.04	...
Total $\chi^2_{\text{red}} (N_{\text{dat}})$		1.01 (1475)	0.98 (1471)



adapted from D. Pitonyak, QCD Evolution 24

Pheno - lattice : tensor charge

$$g_T = \delta u - \delta d$$



green $N_f=2+1+1$


open symbols = no continuum extrapolation

yellow $N_f=2+1$

tension between pheno and lattice ?

adapted from C. Alexandrou, QCD Evolution 24

Including lattice data,
JAM finds **compatibility**,
still under discussion...



But most data
insensitive to tensor charge

For data sensitive to δu , δd

$$\chi^2 = 203 \rightarrow 239$$

$$\chi^2/N_{dat} = 1.02 \rightarrow 1.21$$

Experiment	N_{dat}	χ^2_{red}	
		With LQCD	No LQCD
Belle (cross section) [63]	1094	1.01	1.01
Belle (Artru-Collins) [92]	183	0.74	0.73
HERMES [94]	12	1.13	1.10
COMPASS (p) [95]	26	1.24	0.75
COMPASS (D) [95]	26	0.78	0.76
STAR (2015) [96]	24	1.47	1.67
STAR (2018) [64]	106	1.20	1.04
ETMC δu [28]	1	0.71	...
ETMC δd [28]	1	1.02	...
PNDME δu [25]	1	8.68	...
PNDME δd [25]	1	0.04	...
Total χ^2_{red} (N_{dat})		1.01 (1475)	0.98 (1471)


Future

New data already available:


- Compass SIDIS spin asymmetry on deuteron target with Collins effect & di-hadron mechanism S. Asatryan, DIS 2024 COMPASS Alexeev *et al.*, arXiv:2401.00309
- updated Hermes SIDIS spin asymmetry Airapetian *et al.*, JHEP **12** (20) 010
$$p^\uparrow + p \rightarrow \Lambda^\uparrow + X$$
- Compass π - p^\uparrow Drell-Yan A_T asymmetry Alexeev *et al.*, arXiv:2312.17379
- STAR asymmetry in $p^\uparrow + p \rightarrow \text{jet} + \pi^\pm + X$ hadron-in-jet Collins effect X. Chu, DIS 2024
- STAR asymmetry in $p^\uparrow + p \rightarrow \Lambda^\uparrow + X$ Λ spin transfer STAR, P.R. D**109** (24) 012004
- STAR asymmetry in $p^\uparrow + p \rightarrow \pi^+ \pi^- + X$ di-hadron mechanism B. Surrow, DIS 2024




EIC impact on tensor charge



Collins effect



Abdul-Khalek *et al.*
(EIC Yellow Report),
N.P. **A1026** (22) 122447




di-hadron mechanism

- 1) ETMC '19 *Alexandrou et al., arXiv:1909.00485*
- 2) Mainz '19 *Harris et al., PR. D100 (19) 034513*
- 3) LHPC '19 *Hasan et al., PR. D99 (19) 114505*
- 4) JLQCD '18 *Yamanaka et al., PR. D98 (18) 054516*
- 5) PNDME '18 *Gupta et al., PR. D98 (18) 034503*
- 6) ETMC '17 *Alexandrou et al., PR. D95 (17) 114514; (E) PR. D96 (17) 099906*
- 7) RQCD '14 *Bali et al., PR. D91 (15) 054501*
- 8) LHPC '12 *Green et al., PR. D86 (12) 114509*




EIC impact on tensor charge



Collins effect




 proton + ^3He



Abdul-Khalek *et al.*
(EIC Yellow Report),
N.P. **A1026** (22) 122447

expected precision close to
(or higher than) lattice




di-hadron mechanism

- 1) ETMC '19 *Alexandrou et al., arXiv:1909.00485*
- 2) Mainz '19 *Harris et al., PR. D100 (19) 034513*
- 3) LHPC '19 *Hasan et al., PR. D99 (19) 114505*
- 4) JLQCD '18 *Yamanaka et al., PR. D98 (18) 054516*
- 5) PNDME '18 *Gupta et al., PR. D98 (18) 034503*
- 6) ETMC '17 *Alexandrou et al., PR. D95 (17) 114514; (E) PR. D96 (17) 099906*
- 7) RQCD '14 *Bali et al., PR. D91 (15) 054501*
- 8) LHPC '12 *Green et al., PR. D86 (12) 114509*

Hadron-in-jet Collins effect

$$p^\uparrow + p \rightarrow \text{jet} + \pi^\pm + X$$



STAR 2010-11 $\sqrt{s} = 500 \text{ GeV}$




D'Alesio et al., P.L. **B773** (17) 300



no evolution




Kang et al., P.L. **B774** (17) 635



TMD evolution




M. Grosse-Perdekamp, Transversity 2022



STAR 2012-15 $\sqrt{s} = 200 \text{ GeV}$

Future: EIC impact

hadron-in-jet Collins effect




electron - hadron-in-jet azimuthal correlations

$$|p_T^e + p_T^{\text{jet}}| \ll |p_T^e - p_T^{\text{jet}}|/2 \Rightarrow \text{factorization theorem}$$




theory uncertainty bands from



Arratia *et al.*, P.R. D102 (20) 074015




Summary



TMD PDFs ($x, k_{\perp}; Q^2$) at leading twist for a spin-1/2 hadron (Nucleon)

polarizations
nucleon



quark



		Quark polarization			nomenclature
		Unpolarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)	
Nucleon Polarization	U	$f_1 = \odot$		$h_1^{\perp} = \uparrow \ominus - \downarrow \ominus$	no-name Boer-Mulders
	L		$g_1 = \odot \rightarrow - \ominus \rightarrow$	$h_{1L}^{\perp} = \uparrow \rightarrow - \downarrow \rightarrow$	helicity Kotzinian-Mulders
	T	$f_{1T}^{\perp} = \uparrow \odot - \downarrow \ominus$	$g_{1T} = \uparrow \ominus - \downarrow \ominus$	$h_1 = \uparrow \uparrow - \downarrow \uparrow$ $h_{1T}^{\perp} = \uparrow \rightarrow - \downarrow \rightarrow$	transversity pretzelosity

Sivers

worm gear

- very good knowledge of x-dependence of f_1 and g_1
- good knowledge of k_T -dependence of f_1
- fair knowledge of x-dependence of h_1 and k_T -moments of f_{1T}^{\perp}
- some hints about all others



List of latest extractions



Unpol. TMD	MAP 22 arXiv:2206.07598 , ART23 2305.07473 , MAP24 arXiv:2405.13833
Helicity	arXiv:2409.08110 , MAP24 , arXiv:2409.18078
Transversity	arXiv:1505.05589 , arXiv:1612.06413 , arXiv:2205.00999
Sivers	MAP20 arXiv:2004.14278 , arXiv:2009.10710 , arXiv:2103.03270 , arXiv:2205.00999 , arXiv:2304.14328
Boer-Mulders	arXiv:2004.02117 , arXiv:2407.06277
Worm-gear g1T	arXiv:2110.10253 , arXiv:2210.07268
Worm-gear h1L	
Pretzelosity	arXiv:1411.0580

not mentioned pion TMDs, TMD fragmentation functions, nuclear TMDs

The Nanga Parbat fitting framework



Nanga Parbat: a TMD fitting framework

Nanga Parbat is a fitting framework aimed at the determination of the non-perturbative component of TMD distributions.

Download

You can obtain NangaParbat directly from the github repository:


<https://github.com/MapCollaboration/NangaParbat>



The Artemide fitting framework

<https://teorica.fis.ucm.es/artemide/>

arTeMiDe



News



- 12 Dec 2019:** Version 2.02 released (+manual update).
 - 23 Feb 2019:** Version 1.4 released (+manual update).
 - 21 Jan 2019:** Artemide now has a [repository](#).
- [Archive of older links/news.](#)

Articles, presentations & supplementary materials



- [Extra pictures for the paper arXiv:1902.08474](#)
 - [Seminar of A.Vladimirov in Pavia 2018 on TMD evolution.](#)
 - [Link to the text in Inspire.](#)
- [Archive of older links/news.](#)

Download



[Recent version/release can be found in repository.](#)

About us & Contacts



If you have found mistakes, or have suggestions/questions, please, contact us.

Some extra materials can be found on [Alexey's web-page](#)

Alexey Vladimirov Alexey.Vladimirov@physik.uni-regensburg.de

Ignazio Scimemi ignazios@fis.ucm.es

The TMDLib and TMDPlotter tools

<https://tmdlib.hepforge.org/>

TMD plotter — Density as a function of k_t



Home

TMD PDF

Luminosity

New PDFs

Publications

HEP Links

Parameters

X-axis: min = max = GeV log lin


Y-axis: min = max = log lin

ratio: min = max = log lin

Curves

1. x

$\mu =$ GeV $x =$





Outline



- Backup

The EIC impact at $x=0.001$

$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.001$$

MAPTMD24 2031






EIC # pts. lumi [fb^{-1}]

5x41 1273 2.85

10x100 1611 51.3

18x275 1648 10

(simulation campaign of May 2024)








The EIC impact with 10x100 at $x=0.001$

$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.001$$

	MAPTMD24	2031	# pts.	lumi [fb ⁻¹]
EIC			1611	51.3
10x100				



(simulation campaign of May 2024)



The EIC impact with 10x100 at $x=0.001$






$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.001$$

MAPTMD24 2031

EIC # pts. lumi [fb⁻¹]

10x100 1611 5

(early Science conditions)



courtesy L. Rossi

The EIC impact at $x=0.1$

$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.1$$

$x=0.1$






MAPTMD24 2031

EIC # pts. lumi [fb⁻¹]

5x41 1273 2.85

10x100 1611 51.3

18x275 1648 10




The EIC impact with 10x100 at $x=0.1$

$$\frac{\text{TMD}_q - \langle \text{TMD}_q \rangle}{\langle \text{TMD}_q \rangle} \quad x=0.1$$

MAPTMD24	2031	
EIC	# pts.	lumi [fb⁻¹]
10x100	1611	51.3

(simulation campaign of May 2024)





The EIC impact with 10x100 at $x=0.1$

$$\frac{\text{TMD}q - \langle \text{TMD}q \rangle}{\langle \text{TMD}q \rangle} \quad x=0.1$$

MAPTMD24	2031	
EIC	# pts.	lumi [fb⁻¹]
10x100	1611	5


(early Science conditions)




courtesy L. Rossi

“Normalized” MAPTMD24 TMD FF


$$\frac{D_1(z, P_T; Q)}{D_1(z, 0; Q)}$$



Correlation matrix



Average transverse momenta



clusters = 68% of all replicas