

BabaYaga and its theoretical accuracy

Carlo M. Carloni Calame

INFN, Sezione di Pavia

Working Group on Radiative corrections and generators for low energy hadronic cross section and luminosity

based on [hep-ph/0607181](#) (accepted by **NPB**)

in collaboration with G. Balossini, G. Montagna, O. Nicosini,
F. Piccinini

- BabaYaga web site

<http://www.pv.infn.it/hepcomplex/babayaga.html>

- Theoretical framework of the new **BabaYaga** (BabaYaga@NLO)
- Estimate of the Bhabha theoretical accuracy
 - comparison with **independent generators**
 - comparison with **two loop calculations**
 - **vacuum polarization** uncertainties

★ **BabaYaga@NLO** for $e^+e^- \rightarrow \gamma\gamma$

NEW!

- Conclusions

The original BabaYaga (v. 3.5)

- it is a MCEG for $e^+e^- \rightarrow e^+e^-, \gamma\gamma, \mu^+\mu^-, \pi^+\pi^-$ at flavour factories, developed for luminosity measurement

C.M.C.C. et al., **NPB** 584 (2000)

C.M.C.C., **PLB** 520 (2001)

- the QED RC corrections were included with an (original) QED Parton Shower (PS), allowing for
 - ① fully exclusive multi-photon generation (up to ∞ photons)
 - ② natural inclusion of $\mathcal{O}(\alpha)$ and higher order QED photonic corrections in **leading-log (LL) approximation**
- theoretical error due to **missing $\mathcal{O}(\alpha)$ non-log terms**, not naturally reproduced by the PS.
Estimated accuracies:
 - **0.5%** for Bhabha
 - **$\simeq \mathcal{O}(1\%)$** for $\gamma\gamma$ and $\mu^+\mu^-$

PS and exact $\mathcal{O}(\alpha)$ (**NLO**) matrix elements must be combined and matched. **How?**

- $d\sigma_{LL}^\infty = \Pi(Q^2, \varepsilon) \sum_{n=0}^{\infty} \frac{1}{n!} |\mathcal{M}_{n,LL}|^2 d\Phi_n$
- $d\sigma_{LL}^\alpha = [1 + C_{\alpha,LL}] |\mathcal{M}_0|^2 d\Phi_0 + |\mathcal{M}_{1,LL}|^2 d\Phi_1 \equiv d\sigma_{SV}(\varepsilon) + d\sigma_H(\varepsilon)$
- $d\sigma_{exact}^\alpha = [1 + C_\alpha] |\mathcal{M}_0|^2 d\Phi_0 + |\mathcal{M}_1|^2 d\Phi_1$
- $F_{SV} = 1 + (C_\alpha - C_{\alpha,LL}) \quad F_H = 1 + \frac{|\mathcal{M}_1|^2 - |\mathcal{M}_{1,LL}|^2}{|\mathcal{M}_{1,LL}|^2}$
- $d\sigma_{exact}^\alpha \stackrel{\text{at } \mathcal{O}(\alpha)}{=} F_{SV} (1 + C_{\alpha,LL}) |\mathcal{M}_0|^2 d\Phi_0 + F_H |\mathcal{M}_{1,LL}|^2 d\Phi_1$

$$d\sigma_{matched}^\infty = F_{SV} \Pi(Q^2, \varepsilon) \sum_{n=0}^{\infty} \frac{1}{n!} \left(\prod_{i=0}^n F_{H,i} \right) |\mathcal{M}_{n,LL}|^2 d\Phi_n$$

Contents of the *matched* formula

- F_{SV} and $F_{H,i}$ are infrared safe and account for missing $\mathcal{O}(\alpha)$ non-logs, **avoiding double counting of LL**
- $[\sigma_{matched}^\infty]_{\mathcal{O}(\alpha)} = \sigma_{exact}^\alpha$
- resummation of higher orders LL contributions preserved
- **the cross section is still fully differential in the momenta of the final state particles (e^+ , e^- and $n\gamma$)**
- as a by-product, **part of photonic $\alpha^2 L$ included by means of terms of the type $F_{SV | H,i} \times LL$**

G. Montagna et al., **PLB** 385 (1996)

- **the error is shifted to $\mathcal{O}(\alpha^2)$ (NNLO, 2 loop) not infrared terms: very naively and roughly (for photonic corrections)**

$$\frac{1}{2}\alpha^2 L \equiv \frac{1}{2}\alpha^2 \log \frac{s}{m^2} \sim 0.5 \times 10^{-4}$$

Vacuum Polarization


- $\alpha \rightarrow \alpha(q^2) \equiv \frac{\alpha}{1 - \Delta\alpha(q^2)}$ $\Delta\alpha = \Delta\alpha_{e,\mu,\tau,\text{top}} + \Delta\alpha_{\text{had}}^{(5)}$
- $\Delta\alpha_{\text{had}}^{(5)}$ is a **non-perturbative** contribution. Evaluated with **HADR5N** by F. Jegerlehner. **It returns also an error.**
 - S. Eidelman and F. Jegerlehner, **Z. Phys. C** 67 (1995)
 - F. Jegerlehner, **NPB Proc. Supp.** 131 (2004)
- VP included both **in lowest order** and **(at best) in one-loop** diagrams \Rightarrow part of the 2 loop factorizable corrections are included
- the effect of VP is $\sim 2\%$
- Z exchange included at lowest order.
Its effect is $\mathcal{O}(0.1\%)$ @ 10 GeV

Estimate of the theoretical accuracy

- switching off VP, tuned comparisons with independent calculations/approaches (Labspv, Bhwide)
 - ★ $\Delta\sigma/\sigma < 0.03\%$ on cross sections
 - ★ up-to-0.5% differences between BabaYaga and Bhwide in distribution tails
- comparison with existing perturbative 2-loop calculations
 - ★ currently available
 1. Penin: complete virtual 2-loop photonic corrections (for $Q^2 \gg m_e^2$) plus real radiation in the soft limit
 2. Bonciani et al.: virtual $N_F = 1$ [only electron in the loops] fermionic contributions plus real radiation in the soft limit
 - ★ the photonic and $N_F = 1$ $\mathcal{O}(\alpha^2)$ content of the S+V part in the BabaYaga matched formula can be easily extracted. The terms to be directly compared to 1. and 2. can be read out!
 - ★ the impact of the missing $\mathcal{O}(\alpha^2)$ S+V corrections can be quantified within realistic setup

Light-pair corrections (real & virtual)

- They contribute at $\mathcal{O}(\alpha^2)$, VPC (part of 2-loop $N_F = 1$) and RPC **largely cancel**. **Not included in BabaYaga**.





- To estimate the impact, VPC evaluated as in Jadach et al. ('97); Kniehl ('90); Burgers ('85); Barbieri et al. ('72); RPC evaluated in soft approximation as in Arbuzov et al. ('97)
- the correction does **not exceed 0.05%** in LABS¹ and VLABS² at 1 and 10 GeV (see hep-ph/0607181)

¹ $20^\circ < \vartheta_\pm < 160^\circ$

² $55^\circ < \vartheta_\pm < 125^\circ$

Differences from Penin & Bonciani et al.

- diff. between **Penin** and **Bonciani et al.** and the corresponding BabaYaga content, as $f(\varepsilon)$ and $g(\log(m_e))$. E.g. LABS at 1 GeV



★ differences are **infrared safe**

★ $\delta\sigma(\text{phot.})/\sigma_0 \propto \alpha^2 L$ $\delta\sigma(N_F = 1)/\sigma_0 \propto \alpha^2 L^2$

★ Numerically, in LABS and VLABS,

$$\delta\sigma(\text{phot.}) + \delta\sigma(N_F = 1) < \mathbf{0.015\%} \times \sigma_0$$

$\Delta\alpha_{\text{had}}^{(5)}$ and other $\mathcal{O}(\alpha^2)$ uncertainties

- $\Delta\alpha_{\text{had}}^{(5)}$ is affected by an error, **returned by HADR5N**
 - ★ the error induced on Bhabha cross section is **negligible around the Φ and $< 0.05\%$ at 10 GeV**
 - ★ it is larger (**0.5%**) around J/Ψ resonances, becoming here a limiting factor
 - ★ we wonder why the $\Delta\alpha_{\text{had}}^{(5)}$ error around the Υ is so small. . .
- the 1-loop virtual corrections to the 1-photon real emission are not completely known for Bhabha (even if feasible)
 - ★ relying on the LEP experience and being the error at **the $\alpha^2 L$ level**, the missing corrections are **$\leq 0.05\%$**
- the double real bremsstrahlung contribution is in principle approximated
 - ★ observed **really negligible differences** with the exact matrix elements, calculated with the ALPHA (Caravaglios and Moretti ('95)) algorithm/routine

Summary of theoretical errors

- for **Bhabha cross section**, within realistic setup for luminometry, the theoretical errors of **the new BabaYaga** are summarized

$ \delta^{err} $ (%)	(a)	(b)	(c)	(d)
$ \delta_{VP}^{err} $	0.01	0.00	0.02	0.04
$ \delta_{pairs}^{err} $	0.02	0.03	0.03	0.04
$ \delta_{H,H}^{err} $	0.00	0.00	0.00	0.00
$ \delta_{phot+N_f=1}^{err} $	0.01	0.01	0.00	0.01
$ \delta_{SV,H}^{err} $	0.05	0.05	0.05	0.05
$ \delta_{total}^{err} $	0.09	0.09	0.10	0.14

Table: LABS (a) (c), VLABS (b) (d), 1.02 GeV (a) (b), 10 GeV (c) (d)

Resummation beyond α^2

- ★ with a complete 2-loop generator at hand, (leading-log) resummation beyond α^2 can be neglected?




Figure: Impact of α^2 (solid line) and resummation of higher order ($\geq \alpha^3$) (dotted) corrections on the acollinearity distribution

- ★ resummation beyond α^2 still important

$$e^+e^- \rightarrow \gamma\gamma$$

- ★ the matching is now applied also to $\gamma\gamma$, relying on the 1-loop formulae in Berends and Kleiss **NPB** 186 (1981) and Berends et al. **NPB** 202 (1981)
- ★ double counting, affecting ~ 3.5 , is avoided in the new approach
- e.g., $E_{cms} = 1$ GeV, at least 2 photons with $20^\circ < \vartheta_\gamma < 160^\circ$, $E_\gamma > 0.3$ GeV and varying the acollinearity cut


$\zeta_{\gamma\gamma}$ ($^\circ$)	σ_0 (nb)	$O(\alpha)_{LL}$	$O(\infty)_{LL}$	$O(\alpha)_{ex}$	$O(\infty)_{matched}$
5	329.8	302.5	304.0	304.4	305.6
10	329.8	314.3	314.8	316.3	316.6
15	329.8	320.2	320.4	322.2	322.2
20	329.8	323.6	323.6	325.6	325.4

- $O(\alpha)$ non-log $\simeq 0.7\%$, now included
- ★ estimated theoretical error $\leq 0.1\%$ (VP error is not present here)


$e^+e^- \rightarrow \gamma\gamma$ distributions

- photon energies

markers = $\mathcal{O}(\alpha)$, hist. = $\mathcal{O}(\infty)$



- exponentiation effect



Conclusions

- a new release of BabaYaga is available, matching exact $\mathcal{O}(\alpha)$ corrections with h.o. in a PS approach
- the matching is now applied to **Bhabha** and $e^+e^- \rightarrow \gamma\gamma$
- the **Bhabha theoretical error** is at 2-loop order and **estimated** $\leq 0.15\%$, by considering
 - ★ missing $\mathcal{O}(\alpha^2)$ corrections (pairs, photonic 2-loops, $N_F = 1, \dots$)
 - ★ vacuum polarization uncertainties
 - ★ VP induces larger errors around the charmonium resonances (J/Ψ)
- $e^+e^- \rightarrow \gamma\gamma$ is not affected by VP uncertainty, the theoretical error is **estimated** $\leq 0.1\%$