## **EUROPEAN RADIATION RESEARCH 2012**



Contribution ID: 136

Type: oral (travel award)

## Application of the FISH technique to estimate individual susceptiblity to radiation

Thursday, 18 October 2012 15:15 (15 minutes)

Radiotherapy plays an important role in the treatment of many cancers. Therefore, there are many approaches and strong needs to establish the most appropriate method that would maximize efficiency of therapy and minimize the side effects and late health risk. It should also allow rapid and reliable screening of individuals and stratifications due to lower or upper radiosensitivity for modeling and optimizing a therapeutic procedure and eliminating the highest risk.

The aim of this study was to compare variability in the individual susceptibility to the induction by challenging dose of the aberrations in chromosome 1 in prostate cancer patients (PC) in comparison with benign prostatic hyperplasia (BPH) persons.

Whole blood samples were irradiated with a challenging dose of X-rays, cultured for 3 days, harvested in a metaphase and referred to the FISH procedure using probes for chromosome 1.

Results of our studies have shown differencies in susceptibility to radiation, expressed in significantly elevated or lowered frequency of aberrations for chromosome 1 for cancer patients when compared to BPH group. After irradiation significantly higher levels of all the studied chromosome 1 aberrations, except for deletions, in the case of PC patients were revealed (% AB.C(1)  $5.09 \pm 1.44$  vs  $2.70 \pm 1.21$ ; p<0.001). Additionally, almost five times higher frequencies of acentric fragments were observed in lymphocytes of cancer patients in comparison to BPH group (on average  $2.29 \pm 0.96$  vs  $0.48 \pm 0.27$ , p<0.001). Lymphocytes of the same patients before irradiation (in vivo level) showed significantly higher levels of all studied chromosome 1 aberrations, except for translocations. To estimate individual radiosensitivities after subtracting in vivo levels, the same trends were observed as for the cells after irradiation. In the case of PC patients with the most prominent differences for frequencies of acentric fragments ( $2.27 \pm 1.16$  vs  $0.46 \pm 0.24$ ; p<0.001) in comparison to BPH group. Our results demonstrate that the frequency of acentric fragments detected as the response to a challenging treatment can be proproposed as a very efficient predictor of susceptibility to planned treatment. Limitation of analysis of acentric fragments only after challenging irradiation may greatly speed up the diagnosis.

Primary author: Dr MISZCZYK, Justyna (Institute of Nuclear Physics PAN, Kraków, Poland)

**Co-authors:** Prof. CEBULSKA-WASILEWSKA, Antonina (Institute of Nuclear Physics PAN, Kraków, Poland); Dr DOBROWOLSKA, Barbara (Department and Clinic of Urology, Collegium Medicum, The Jagiellonian University, Kraków, Poland); Prof. DOBROWOLSKI, Zygmunt (Department and Clinic of Urology, Collegium Medicum, The Jagiellonian University, Kraków, Poland)

Presenter: Dr MISZCZYK, Justyna (Institute of Nuclear Physics PAN, Kraków, Poland)

Session Classification: Awardees 1

Track Classification: Biological and Physical Dosimetry