

QFT as a set of coupled ODEs

Seminar Università di Genova / INFN

WIP w/ Gregoire Mathys, Joao Penedones, Jiaxin Qiao, Xiang Zhao



Manuel Loparco (Unito), 05/11/2025

Motivation

- QFT is fantastic in statistical physics, particle physics, cosmology
- Many phenomenologically relevant QFTs are strongly coupled

How do we formulate predictions in strongly coupled QFTs?

Method	Downsides
Conformal bootstrap	Only CFT
S-matrix bootstrap	Only 2-to-2
Hamiltonian truncation	Cutoff breaks Lorentz, many divergences
Lattice	Expensive, sign problem, no dynamics
Integrability	Only 2D and extra conserved charges
Holography	Only large N and SUSY

A framework in the continuum, with no divergences and (maybe??) cheap

A framework in the continuum, with no divergences and (maybe??) cheap

• In CFT, local observables are fixed by $\{\Delta_i, C_{ijk}\}$

$$\langle \mathcal{O}_i(x)\mathcal{O}_j(0)\rangle = \frac{\delta_{ij}}{x^{2\Delta_i}}$$
 $\mathcal{O}_i(x)\mathcal{O}_j(0) = \sum_k \frac{C_{ijk}}{x^{\Delta_i + \Delta_j - \Delta_k}} \left(\mathcal{O}_k(0) + \text{descendants}\right)$


A framework in the continuum, with no divergences and (maybe??) cheap

• In CFT, local observables are fixed by $\{\Delta_i, C_{ijk}\}$


$$\langle \mathcal{O}_i(x)\mathcal{O}_j(0)\rangle = \frac{\delta_{ij}}{x^{2\Delta_i}}$$
 $\mathcal{O}_i(x)\mathcal{O}_j(0) = \sum_k \frac{C_{ijk}}{x^{\Delta_i + \Delta_j - \Delta_k}} \left(\mathcal{O}_k(0) + \text{descendants}\right)$

- QFT in AdS defines a family of CFTs (without stress tensor)
- Observables in QFT in AdS fixed by $\{\Delta_i(\lambda),\ b_i^{\hat{\mathcal{O}}}(\lambda)\ , C_{ijk}(\lambda)\}$


The Boundary Operator Expansion (BOE)



The Boundary Operator Expansion (BOE)



The Boundary Operator Expansion (BOE)



A framework in the continuum, with no divergences and (maybe?) cheap

• In CFT, local observables are fixed by $\{\Delta_i, C_{ijk}\}$

$$\langle \mathcal{O}_i(x)\mathcal{O}_j(0)\rangle = \frac{\delta_{ij}}{x^{2\Delta_i}}$$
 $\mathcal{O}_i(x)\mathcal{O}_j(0) = \sum_k \frac{C_{ijk}}{x^{\Delta_i + \Delta_j - \Delta_k}} \left(\mathcal{O}_k(0) + \text{descendants}\right)$

- QFT in AdS defines a family of CFTs (without stress tensor)
- Observables in QFT in AdS fixed by $\{\Delta_i(\lambda),\ b_i^{\hat{\mathcal{O}}}(\lambda)\ , C_{ijk}(\lambda)\}$
- For arbitrary λ , we can compute <u>universally</u> the derivatives of the data

A framework in the continuum, with no divergences and (maybe?) cheap


• In CFT, local observables are fixed by $\{\Delta_i, C_{ijk}\}$

$$\langle \mathcal{O}_i(x)\mathcal{O}_j(0)\rangle = \frac{\delta_{ij}}{x^{2\Delta_i}}$$
 $\mathcal{O}_i(x)\mathcal{O}_j(0) = \sum_k \frac{C_{ijk}}{x^{\Delta_i + \Delta_j - \Delta_k}} \left(\mathcal{O}_k(0) + \text{descendants}\right)$

- QFT in AdS defines a family of CFTs (without stress tensor)
- Observables in QFT in AdS fixed by $\{\Delta_i(\lambda),\ b_i^{\hat{\mathcal{O}}}(\lambda)\ , C_{ijk}(\lambda)\}$
- For arbitrary λ , we can compute <u>universally</u> the derivatives of the data
- Truncated system of ODEs can in principle be solved iteratively, flowing the data to <u>strong coupling</u> and to <u>flat space</u>


What are local correlation functions in AdS?

And how do we compute them?



What is the S-matrix in QFT in flat space?

And how do we compute it?



$$S = S_{\text{QFT}_2} + \lambda \int_{\text{AdS}_2} \hat{\Phi}$$

"QFT data" fully fixes all local observables

$$\Delta_i(\lambda)\,,\quad C_{ijk}(\lambda)\,,\quad b_i^{\hat{\mathcal{O}}_a}(\lambda)$$

Taking $\lambda \to \lambda + \delta \lambda$, with PT we compute

$$\frac{\Delta_i}{d\lambda}(\lambda)$$
, $\frac{C_{ijk}}{d\lambda}(\lambda)$, $\frac{db_i^{\hat{\mathcal{O}}_a}}{d\lambda}(\lambda)$

Truncated system of ODEs can be iteratively solved

Inspiration: [Hollands, Wald, '23]

1D conformal manifolds: [Behan, '17]

Outline

Correlation functions from QFT data

Perturbation theory, once and for all


Examples and checks

Further properties of the flow equations

Observables from QFT data

Geometry

Embedding space formalism



• Embed AdS_2 in $\mathbb{R}^{1,2}$

$$X^A X^B \eta_{AB} = -R^2$$


Boundary = space of light rays


$$P^A P^B \eta_{AB} = 0, \qquad P^A \sim \lambda P^A$$


• Poincaré metric $ds^2 = \frac{\mathrm{d}z^2 + \mathrm{d}\tau^2}{z^2}$

$$P_{12} \equiv -2P_1 \cdot P_2$$
$$\Delta_{123} \equiv \Delta_1 + \Delta_2 - \Delta_3$$

At any point in the RG flow










$$P_{12} \equiv -2P_1 \cdot P_2$$

$$\Delta_{123} \equiv \Delta_1 + \Delta_2 - \Delta_3$$

At any point in the RG flow








These are the coefficients in the BOE

$$\hat{\mathcal{O}}(z,\tau) = \sum_{i} z^{\Delta_i} b_i^{\hat{\mathcal{O}}} \left(\mathcal{O}_i(\tau) + \text{descendants} \right)$$

• State-operator map: any other correlation function is built from QFT data

$$=\sum_{i,j,k}b_i^{\hat{\mathcal{O}}_1}b_j^{\hat{\mathcal{O}}_2}b_k^{\hat{\mathcal{O}}_3}C_{ijk} \, (\text{kinematic block})_{ijk}$$

• State-operator map: any other correlation function is built from QFT data



The QFT data

QFT observables are thus fully fixed by these functions

$$\Delta_i(\lambda), \quad C_{ijk}(\lambda), \quad b_i^{\hat{\mathcal{O}}_a}(\lambda)$$

• For boundary observables, we only need

$$\Delta_i(\lambda), \quad C_{ijk}(\lambda), \quad b_i^{\hat{\Phi}}(\lambda)$$

ODEs for these functions come by solving perturbation theory once and for all

Perturbation theory, once and for all Shifting data

Change the coupling infinitesimally

$$S = S_{\text{QFT}_2} + (\lambda + \delta \lambda) \int_X \hat{\Phi}$$

At leading order, the only effect is

$$\delta \Delta_{i}(\lambda) = \delta \lambda \ \Delta'_{i}(\lambda)$$

$$\delta C_{ijk}(\lambda) = \delta \lambda \ C'_{ijk}(\lambda)$$

$$\delta b_{\hat{\mathcal{O}}_{i}}(\lambda) = \delta \lambda \ b'_{\hat{\mathcal{O}}_{i}}(\lambda)$$

2-pt function

$$S = S_{\text{QFT}_2} + (\lambda + \delta \lambda) \int_X \hat{\Phi}$$

$$\delta \Delta_i(\lambda) = \delta \lambda \ \Delta'_i(\lambda)$$

Perturbation theory, once and for all 2-pt function

$S = S_{\text{QFT}_2} + (\lambda + \delta \lambda) \int_{\mathbf{v}} \hat{\Phi}$

$$\delta\Delta_i(\lambda) = \delta\lambda \ \Delta_i'(\lambda)$$

At the same time,

$$\langle \mathcal{O}(P_1)\mathcal{O}(P_2)\rangle_{(\lambda+\delta\lambda)} = \langle \mathcal{O}(P_1)\mathcal{O}(P_2)\rangle_{(\lambda)} - \delta\lambda \int_X \langle \mathcal{O}(P_1)\mathcal{O}(P_2)\hat{\Phi}(X)\rangle_{(\lambda)}$$


Perturbation theory, once and for all 2-pt function

$$S = S_{\text{QFT}_2} + (\lambda + \delta \lambda) \int_{X} \hat{\Phi}$$

$$\delta \Delta_i(\lambda) = \delta \lambda \ \Delta'_i(\lambda)$$

At the same time,

$$\langle \mathcal{O}(P_1)\mathcal{O}(P_2)\rangle_{(\lambda+\delta\lambda)} = \langle \mathcal{O}(P_1)\mathcal{O}(P_2)\rangle_{(\lambda)} - \delta\lambda \int_X \langle \mathcal{O}(P_1)\mathcal{O}(P_2)\hat{\Phi}(X)\rangle_{(\lambda)}$$



Perturbation theory, once and for all 2-pt function

$$S = S_{\text{QFT}_2} + (\lambda + \delta \lambda) \int_X \hat{\Phi}$$

$$\delta \Delta_i(\lambda) = \delta \lambda \ \Delta'_i(\lambda)$$


At the same time,

$$\langle \mathcal{O}(P_1)\mathcal{O}(P_2)\rangle_{(\lambda+\delta\lambda)} = \langle \mathcal{O}(P_1)\mathcal{O}(P_2)\rangle_{(\lambda)} - \delta\lambda \int_X \langle \mathcal{O}(P_1)\mathcal{O}(P_2)\hat{\Phi}(X)\rangle_{(\lambda)}$$

Renormalization conditions


$$\mathcal{O}_{(\lambda+\delta\lambda)} = (1 + Z(\epsilon)) \mathcal{O}^{R}_{(\lambda+\delta\lambda)}$$

$$\langle \mathcal{O}_i^R(P_1)\mathcal{O}_j^R(P_2)\rangle_{(\lambda+\delta\lambda)} \equiv \frac{\delta_{ij}}{P_{12}^{\Delta_i+\delta\Delta_i}}$$




Perturbation theory, once and for all Conformal blocks

$$\int_{X,\epsilon} \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\Phi}(X)|0\rangle$$



Perturbation theory, once and for all Conformal blocks

$$\int_{X,\epsilon} \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\Phi}(X)|0\rangle = \int_{X,\epsilon} \sum_j \int_P \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)|\Delta_j, P\rangle \langle \Delta_j, P|\hat{\Phi}(X)|0\rangle$$




Conformal blocks

$$\int_{X,\epsilon} \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\Phi}(X)|0\rangle = \int_{X,\epsilon} \sum_j \int_P \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)|\Delta_j, P\rangle \langle \Delta_j, P|\hat{\Phi}(X)|0\rangle$$








$$C_{iij}$$

$$b_{j}^{\hat{\Phi}}$$

Conformal blocks

$$\int_{X,\epsilon} \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\Phi}(X)|0\rangle = \int_{X,\epsilon} \sum_j \int_P \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)|\Delta_j, P\rangle \langle \Delta_j, P|\hat{\Phi}(X)|0\rangle$$




$$=\frac{1}{P_{12}^{\Delta_i}}\int_{X,\epsilon}\sum_j C_{iij}\,b_j^{\hat\Phi}\,G_{\Delta_i,\Delta_j}(\chi) \qquad \text{Conformal block}$$

$$\chi \equiv \frac{P_{12}}{(P_1 \cdot X)(P_2 \cdot X)}$$

Conformal blocks

We want to carry out these integrals universally

$$\int_{X,\epsilon} \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\Phi}(X)|0\rangle = \int_{X,\epsilon} \sum_j \int_P \langle 0|\mathcal{O}_i(P_1)\mathcal{O}_i(P_2)|\Delta_j, P\rangle \langle \Delta_j, P|\hat{\Phi}(X)|0\rangle$$



$$= \frac{1}{P_{12}^{\Delta_i}} \int_{X,\epsilon} \sum_j C_{iij} b_j^{\hat{\Phi}} G_{\Delta_i,\Delta_j}(\chi)$$

$$\uparrow \qquad \qquad \chi \equiv \frac{P_{12}}{(P_1 \cdot X)(P_2 \cdot X)}$$

Can we swap?


Local blocks

- Big technical parenthesis: must use local blocks [Levine, Paulos '23], [Meineri, Penedones, Spirig '23]
- Naive conformal blocks present unphysical cuts
- Local blocks form a basis with physical analytic structure
- Endowed with an extra free parameter α
- There exists an integral transform

$$F^{(\alpha)}[G] = G^{(\alpha)}$$

$$F^{(\alpha)}[I] = I^{(\alpha)}$$


• Expanding for small $\delta \lambda$ and matching,



The flow equations

• The ODEs:


$$rac{d\Delta_i}{d\lambda} = \sum_l b_l^{\hat{\Phi}} C_{iil} \mathcal{I}^{(lpha)}(\Delta_l)$$



$$\sum_{j} b_{j}^{\hat{\Phi}} \mathcal{O}_{j}$$
 $\hat{\Phi}$
 $\sum_{l} b_{l}^{\hat{\Phi}} \mathcal{O}_{l}$

$$rac{db_i^{\hat{\Phi}}}{d\lambda} = \sum_{j,l} b_l^{\hat{\Phi}} b_j^{\hat{\Phi}} C_{ijl} \mathcal{J}_{\Delta_i}^{(lpha)}(\Delta_l,\Delta_j)$$

$$\frac{dC_{ijk}}{d\lambda} = \sum_{l,m} b_l^{\hat{\Phi}} C_{lim} C_{mjk} \mathcal{K}_{\Delta_i \Delta_j \Delta_k}^{(\alpha)} (\Delta_l, \Delta_m) + (ijk) \text{ perms.}$$



Examples and Checks

ExampleScalar GFF

$$S = -\frac{1}{2} \int \sqrt{g} d^2x \left(\partial_{\mu} \hat{\phi} \partial^{\mu} \hat{\phi} + m^2 \hat{\phi}^2 \right)$$

• In this example $\;\hat{\Phi}=\hat{\phi}^2\;\; {\rm and} \;\;\; \lambda=\frac{m^2}{2}\;\; .$

ExampleScalar GFF

$$S = -\frac{1}{2} \int \sqrt{g} d^2x \left(\partial_{\mu} \hat{\phi} \partial^{\mu} \hat{\phi} + m^2 \hat{\phi}^2 \right)$$

- In this example $\;\hat{\Phi}=\hat{\phi}^2\;\; {\rm and} \;\;\; \lambda=\frac{m^2}{2}$.
- The theory is solvable. It has, for example, primaries ϕ and $[\phi^2]_n$

$$\Delta_{\phi}(m^2) = \frac{1}{2} \pm \sqrt{\frac{1}{4} + m^2} \qquad \Delta_{[\phi^2]_n}(m^2) = 2\Delta_{\phi}(m^2) + 2n$$

ExampleScalar GFF

$$S = -\frac{1}{2} \int \sqrt{g} d^2x \left(\partial_{\mu} \hat{\phi} \partial^{\mu} \hat{\phi} + m^2 \hat{\phi}^2 \right)$$

- In this example $\; \hat{\Phi} = \hat{\phi}^2 \; \; \text{and} \; \; \; \lambda = \frac{m^2}{2} \; \; \; .$
- The theory is solvable. It has, for example, primaries ϕ and $[\phi^2]_n$

$$\Delta_{\phi}(m^2) = \frac{1}{2} \pm \sqrt{\frac{1}{4} + m^2} \qquad \Delta_{[\phi^2]_n}(m^2) = 2\Delta_{\phi}(m^2) + 2n$$

$$b_{\phi}^{\hat{\phi}}(m^2) = \sqrt{\frac{\Gamma(\Delta)}{2\sqrt{\pi}\Gamma(\Delta + \frac{1}{2})}} \qquad b_{[\phi^2]_n}^{\hat{\phi}^2}(m^2) = \frac{\Gamma(\Delta)}{\sqrt{2\pi}\Gamma(\Delta + \frac{1}{2})} \sqrt{\frac{(\frac{1}{2})_n(\Delta)_n^3(2\Delta - \frac{1}{2})_n}{n!\left(\Delta - \frac{1}{4}\right)_n(2\Delta)_n\left(\Delta + \frac{1}{4}\right)_n\left(\Delta + \frac{1}{2}\right)_n}}$$

Example

Scalar GFF

$$\frac{1}{2\Delta_{\phi} - 1} = \frac{d\Delta_{\phi}}{dm^2} = \frac{1}{2} \sum_{n=0}^{\infty} b_{[\phi^2]_n}^{\hat{\phi}^2} C_{[\phi^2]_n \phi \phi} \mathcal{I}^{(\alpha)} (2\Delta_{\phi} + 2n)$$

Example

Scalar GFF

From boundary expansion of $\langle \hat{\phi}^2 \hat{\phi}^2 \rangle$


$$\frac{1}{2\Delta_{\phi}-1}=\frac{d\Delta_{\phi}}{dm^2}=\frac{1}{2}\sum_{n=0}^{\infty}b_{[\phi^2]_n}^{\hat{\phi}^2}C_{[\phi^2]_n\phi\phi}\mathcal{I}^{(\alpha)}(2\Delta_{\phi}+2n)$$
 From harmonic analysis on $\langle\phi\phi\phi\phi\rangle$

[Simmons-Duffin, Kravchuk, Karateev, '19]

$$\mathcal{I}^{(\alpha)}(\Delta) = \frac{\pi\Gamma(\Delta + \frac{1}{2})\Gamma(2\alpha - 1)}{2^{2\alpha - 2}(\Delta - 1)\Gamma(\frac{\Delta}{2})\Gamma(\frac{\Delta}{2} + 1)\Gamma(\frac{\Delta - 1}{2} + \alpha)\Gamma(\alpha - \frac{\Delta}{2})}$$

Flow of scaling dimensions $\frac{d\Delta_{\phi}}{dm^2} = \frac{1}{2} \sum_{n=0}^{n_{\text{max}}} b_{[\phi^2]_n}^{\hat{\phi}^2} C_{[\phi^2]_n \phi \phi} \mathcal{I}^{(\alpha)}(2\Delta_{\phi} + 2n)$

$$\frac{d\Delta_{\phi}}{dm^2} = \frac{1}{2} \sum_{n=0}^{n_{\text{max}}} b_{[\phi^2]_n}^{\hat{\phi}^2} C_{[\phi^2]_n \phi \phi} \mathcal{I}^{(\alpha)} (2\Delta_{\phi} + 2n)$$




Sum over normal blocks converges only if

$$\Delta_i < \frac{3}{2}$$

Flow of scaling dimensions

$$\frac{d\Delta_{\phi}}{dm^2} = \frac{1}{2} \sum_{n=0}^{n_{\text{max}}} b_{[\phi^2]_n}^{\hat{\phi}^2} C_{[\phi^2]_n \phi \phi} \mathcal{I}^{(\alpha)} (2\Delta_{\phi} + 2n)$$



Universal error estimate:

$$\Delta_{\max}^{2(\Delta_i - \alpha) + \Delta_{\hat{\Phi}}}$$


From harmonic analysis on $\langle \phi \phi \phi \phi \phi \phi \phi \phi \rangle$

$$\frac{db_{\phi^2}^{\hat{\phi}^2}}{dm^2} = \frac{1}{2} \sum_{l,j}^{\infty} b_{[\phi^2]_l}^{\hat{\phi}^2} b_{[\phi^2]_j}^{\hat{\phi}^2} C_{\phi^2[\phi^2]_l[\phi^2]_j} \mathcal{J}_{2\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, 2\Delta_{\phi} + 2j)$$


From harmonic analysis on $\langle \phi \phi \phi \phi \phi \phi \rangle$ $\frac{db_{\phi^2}^{\hat{\phi}^2}}{dm^2} = \frac{1}{2} \sum_{l,j}^{\infty} b_{[\phi^2]_l}^{\hat{\phi}^2} b_{[\phi^2]_j}^{\hat{\phi}^2} C_{\phi^2[\phi^2]_l[\phi^2]_j} \mathcal{J}_{2\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, 2\Delta_{\phi} + 2j)$ $\frac{db_{\phi^2}^{\hat{\phi}^2}}{dm^2} = \frac{1}{2} \sum_{l,j}^{\infty} b_{[\phi^2]_l}^{\hat{\phi}^2} b_{[\phi^2]_j}^{\hat{\phi}^2} C_{\phi^2[\phi^2]_l[\phi^2]_j} \mathcal{J}_{2\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, 2\Delta_{\phi} + 2j)$ $\frac{db_{\phi^2}^{\hat{\phi}^2}}{dm^2} = \frac{2\sqrt{\pi}\Gamma\left(\frac{\Delta_{1}+1}{2}\right)\Gamma\left(\Delta_{2}+\frac{1}{2}\right)\Gamma\left(\alpha-\frac{1}{2}\right)\Gamma\left(\alpha-\frac{\Delta_{23}}{2}\right)}{\left(\Delta_{1}-1\right)\Delta_{22}\Gamma\left(\frac{\Delta_{123}}{2}\right)\Gamma\left(\alpha-\frac{\Delta_{1}}{2}\right)\Gamma\left(\alpha+\frac{\Delta_{132}-1}{2}\right)\Gamma\left(\frac{\Delta_{2}+\Delta_{3}+1}{2}\right)}$

$$\mathcal{J}_{\Delta_{3}}^{(\alpha)}(\Delta_{1}, \Delta_{2}) = \frac{2\sqrt{\pi}\Gamma\left(\frac{\Delta_{1}+1}{2}\right)\Gamma\left(\Delta_{2}+\frac{1}{2}\right)\Gamma\left(\alpha-\frac{1}{2}\right)\Gamma\left(\alpha-\frac{\Delta_{23}}{2}\right)}{(\Delta_{1}-1)\Delta_{23}\Gamma\left(\frac{\Delta_{123}}{2}\right)\Gamma\left(\alpha-\frac{\Delta_{1}}{2}\right)\Gamma\left(\alpha+\frac{\Delta_{132}-1}{2}\right)\Gamma\left(\frac{\Delta_{2}+\Delta_{3}+1}{2}\right)} \times {}_{3}F_{2}\left(\begin{array}{c} \frac{\Delta_{132}}{2}, & \frac{\Delta_{1}-1}{2}, \\ \Delta_{1}+\frac{1}{2}, & \alpha-\frac{\Delta_{23}}{2}+\frac{\Delta_{1}-1}{2} \end{array}; 1\right)$$

$$\frac{db_{\phi^2}^{\hat{\phi}^2}}{dm^2} = \frac{1}{2} \sum_{j=0}^{j_{\text{max}}} \sum_{l=0}^{l_{\text{max}}} b_{[\phi^2]_l}^{\hat{\phi}^2} b_{[\phi^2]_j}^{\hat{\phi}^2} C_{\phi^2[\phi^2]_l[\phi^2]_j} \mathcal{J}_{2\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, 2\Delta_{\phi} + 2j)$$



$$\frac{db_{\phi^2}^{\hat{\phi}^2}}{dm^2} = \frac{1}{2} \sum_{j=0}^{j_{\text{max}}} \sum_{l=0}^{l_{\text{max}}} b_{[\phi^2]_l}^{\hat{\phi}^2} b_{[\phi^2]_j}^{\hat{\phi}^2} C_{\phi^2[\phi^2]_l[\phi^2]_j} \mathcal{J}_{2\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, 2\Delta_{\phi} + 2j)$$



 $\Delta_{\phi} = 2.1$


$$0 = \frac{dC_{\phi\phi1}}{dm^{2}} = \sum_{l=0}^{\infty} b_{[\phi^{2}]_{l}}^{\hat{\phi}^{2}} C_{\phi\phi[\phi^{2}]_{l}} \left(\mathcal{K}_{\Delta_{\phi},\Delta_{\phi},0}^{(\alpha)}(2\Delta_{\phi}+2l,\Delta_{\phi}) + \mathcal{K}_{\Delta_{\phi},0,\Delta_{\phi}}^{(\alpha)}(2\Delta_{\phi}+2l,\Delta_{\phi}) + \mathcal{K}_{0,\Delta_{\phi},\Delta_{\phi}}^{(\alpha)}(2\Delta_{\phi}+2l,2\Delta_{\phi}+2l) \right)$$

$$0 = \frac{dC_{\phi\phi1}}{dm^{2}} = \sum_{l=0}^{\infty} b_{[\phi^{2}]_{l}}^{\hat{\phi}^{2}} C_{\phi\phi[\phi^{2}]_{l}} \Big(\mathcal{K}_{\Delta_{\phi},\Delta_{\phi},0}^{(\alpha)} (2\Delta_{\phi} + 2l, \Delta_{\phi}) + \mathcal{K}_{\Delta_{\phi},0,\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, \Delta_{\phi}) + \mathcal{K}_{0,\Delta_{\phi},\Delta_{\phi}}^{(\alpha)} (2\Delta_{\phi} + 2l, 2\Delta_{\phi} + 2l) \Big)$$

This is the numerical bottleneck for now:

$$\mathcal{K}_{\Delta_i \Delta_j \Delta_k}^{(\alpha)}(\Delta_l, \Delta_m) = \int \frac{dc_1 dc_2 dc_3}{(2\pi i)^3} f_{\Delta_i \Delta_j \Delta_k}^{\Delta_l, \Delta_m}(c_1, c_2, c_3)$$

$$0 = \frac{dC_{\phi\phi1}}{dm^2} = \sum_{l=0}^{\infty} b_{[\phi^2]_l}^{\hat{\phi}^2} C_{\phi\phi[\phi^2]_l} \left(\mathcal{K}_{\Delta_{\phi},\Delta_{\phi},0}^{(\alpha)}(2\Delta_{\phi} + 2l, \Delta_{\phi}) + \mathcal{K}_{\Delta_{\phi},0,\Delta_{\phi}}^{(\alpha)}(2\Delta_{\phi} + 2l, \Delta_{\phi}) + \mathcal{K}_{0,\Delta_{\phi},\Delta_{\phi}}^{(\alpha)}(2\Delta_{\phi} + 2l, \Delta_{\phi}) + \mathcal{K}_{0,\Delta_{\phi}}^{(\alpha)}(2\Delta_{\phi} + 2l, \Delta_{\phi})$$



Further properties of the flow equations

The flat space limit and the S-matrix

- The coupling λ is in units of R
- Flow can be seen as happening from heavily curved AdS to flat space
- In principle, spectrum and S-matrix can be extracted

$$\frac{m_i}{m_0} = \lim_{\lambda \to \infty} \frac{\Delta_i(\lambda)}{\Delta_0(\lambda)} \qquad S(s) \sim \lim_{\lambda \to \infty} \sum_{\mathcal{O}_{\Delta}} f(\Delta) C_{\phi\phi\mathcal{O}_{\Delta}}^2(\lambda) \delta\left(\frac{\sqrt{s}}{m} - \frac{\Delta(\lambda)}{\Delta_{\phi}(\lambda)}\right)$$


[Meineri, Penedones, Spirig, '23]

[Paulos, Penedones, Toledo, van Rees, Vieira, '16]

More: [van Rees, Zhao, '22], [Polchinski, '99], [Cordova, He, Paulos '22]

Level repulsion

- Eigenvalues of hamiltonians which depend on continuous parameters don't cross
- Proven in RMT and finite dimensional QM
- Expected to hold generically in QFT
- We can prove it with our equations



$$0 < \Delta_1 - \Delta_2 \ll 1$$
:

$$\frac{d^2(\Delta_1 - \Delta_2)}{d\lambda^2} = \frac{c}{(\Delta_1 - \Delta_2)^3}$$

Mergers and annihilations of theories

Expected when operators hit marginality

[Kaplan et al 2009, Gorbenko et al 2018]

From the first flow equation,

$$\Delta_* \gtrsim 1$$
, $\frac{d\Delta_i}{d\lambda} \sim \frac{c}{\Delta_* - 1}$

$$\Delta_i(\lambda) \sim \Delta_i(\lambda_c) \pm \sqrt{2c(\lambda - \lambda_c)}$$
 $c = \frac{C_{ii*}^2 b_1^{\Phi}}{C_{***}}$


Mergers and annihilations of theories

- Expected when operators hit marginality
 - [Kaplan et al 2009, Gorbenko et al 2018]
- From the first flow equation,

$$\Delta_* \gtrsim 1$$
, $\frac{d\Delta_i}{d\lambda} \sim \frac{c}{\Delta_* - 1}$


$$\Delta_i(\lambda) \sim \Delta_i(\lambda_c) \pm \sqrt{2c(\lambda - \lambda_c)}$$
 $c = \frac{C_{ii*}^2 b_1^{\hat{\Phi}}}{C_{***}}$

Example in free massive theory:



Free bulk and interacting boundary

- There exists a family of 1D CFTs called *Long Range Ising*
- Critical point of Ising model with decaying interactions
- It is equivalent to the following construction
 - Take free massive scalar in AdS2
 - Turn on $\,\phi^4\,$ only on the boundary
 - Flow to IR fixed point on boundary
- Fits in our flow equations framework



Conclusion

Conclusion Sumary

- QFT in AdS can be phrased in terms of $\Delta_i(\lambda)$, $C_{ijk}(\lambda)$, $b_i^{\hat{\mathcal{O}}}(\lambda)$
- Their evolution under RG flows is governed by coupled ODEs
- No more divergences to deal with
- Fully non-perturbative
- Nontrivial checks for scalar and fermion GFF
- They know a lot about QFT (level repulsion, merger-annihilation, etc..)
- What next?

Conclusion Short term wishes

- Efficiency
- Solve simple interacting cases (ex: LRI family)
- More concise form? No $b_i^{\hat{\Phi}}$, no λ

Conclusion

Longer term wishes

- Generalization to multiple deformations (maybe trivial)
- Ising Field Theory and flows between minimal models [Lauria, Milam, van Rees '24]
- O(N) nonlinear sigma model and O(N) GN model [Copetti, Di Pietro, Ji, Komatsu '24]
- Generalization to higher dimensions
 - •
- YM₄ [Ciccone, De Cesare, Di Pietro, Serone '24]
- QCD₄!

What is QFT?

- This framework inherits all axioms from CFT (except bdy stress tensor)
- Existence of bulk stress tensor implies [Meineri, Penedones, Spirig '23]

$$c^{\text{UV}} = \sum_{i} \frac{24\Gamma(2\Delta_{i})}{4^{\Delta_{i}}\Gamma^{2}(\Delta_{i}+2)} \left(b_{i}^{\hat{\Theta}}\right)^{2} \qquad \Delta_{i} = \sum_{j} b_{j}^{\hat{\Theta}} C_{iij} \kappa(\Delta_{j}, \alpha)$$


Possibly exciting for axiomatic (and computable) approach to QFT

Thank you

Backup

Local blocks


Analytic structure



[Meineri, Penedones, Spirig '23]

Local blocks

The dispersive integral



 $\lfloor \chi'$

Consider the following trivial rewriting

$$\langle \mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\mathcal{O}}(X)\rangle = \frac{1}{P_{12}^{\Delta_i}}f(\chi) \qquad f(\chi) = \chi^{\alpha} \oint_{\chi} \frac{d\chi'}{2\pi i} \frac{1}{\chi' - \chi} \frac{f(\chi')}{\chi'^{\alpha}}$$

Imposing a physical analytic structure for the correlator,

$$f(\chi) = \chi^{\alpha} \int_{-\infty}^{0} \frac{d\chi'}{2\pi i} \frac{1}{\chi' - \chi} \operatorname{Disc}_{\chi' \le 0} \left(\frac{f(\chi')}{\chi'^{\alpha}} \right)$$

Now plug in $f(\chi) = \sum_j b_{\hat{\mathcal{O}}j} C_{iij} G_{\Delta_i,\Delta_j}(\chi)$. Swapping (valid for $\alpha > \Delta_i + \frac{\Delta_{\hat{\mathcal{O}}}}{2}$)

$$\langle \mathcal{O}_i(P_1)\mathcal{O}_i(P_2)\hat{\mathcal{O}}(X)\rangle = \frac{1}{P_{12}^{\Delta_i}} \sum_j b_{\hat{\mathcal{O}}_j} C_{iij} G_{\Delta_i, \Delta_j}^{(\alpha)}(\chi) \qquad G_{\Delta_i, \Delta_j}^{(\alpha)}(\chi) \equiv \chi^{\alpha} \int_{-\infty}^0 \frac{d\chi'}{2\pi i} \frac{1}{\chi' - \chi} \operatorname{Disc}_{\chi' \leq 0} \left(\frac{G_{\Delta_i, \Delta_j}(\chi')}{\chi'^{\alpha}} \right)$$

Renormalization condition

• Explicitly, we choose

$$\langle \mathcal{O}_i^R(P_1)\mathcal{O}_j^R(P_2)\rangle_{(\lambda+\delta\lambda)} \equiv \frac{\delta_{ij}}{P_{12}^{\Delta_i+\delta\Delta_i}}$$

$$\langle \mathcal{O}_i^R(P_1)\mathcal{O}_j^R(P_2)\mathcal{O}_k^R(P_3)\rangle_{(\lambda+\delta\lambda)} \equiv \frac{C_{ijk} + \delta C_{ijk}}{P_{12}^{\Delta_{ijk} + \delta\Delta_{ijk}} P_{23}^{\Delta_{jki} + \delta\Delta_{jki}} P_{13}^{\Delta_{ikj} + \delta\Delta_{ikj}}}$$

$$\mathcal{O}_i(P)\Big|_{(\lambda+\delta\lambda)} = \left(\delta_i^j + \delta Z_i^j \sum_{n=0}^{\infty} a_n D_P^n\right) \mathcal{O}_j^R(P)\Big|_{(\lambda+\delta\lambda)}$$