



Attosecond x-ray free-electron lasers

**Agostino Marinelli
SLAC National Accelerator Laboratory
and Stanford University**

Acknowledgements

XLEAP/XLEAP-II Collaboration

SLAC

A. Marinelli, J. Cryan, et al.

ANL

A. Zholents, J. Xu, M. Qian et al.

UCLA

S. Carbajo et al.

Hard X-ray Attosecond Pulses

University of Tokyo/SACLA

I. Inoue

SLAC

A. Marinelli, R. Robles, A. Halavanau

Attosecond Campaign (PIs)

LCLS - Leads

James Cryan

Agostino Marinelli

SLAC - Pulse

Taran Driver

Phil Bucksbaum

Imperial College

Vitali Averbukh

Jon Marangos

UAM

Fernando Martin

Antonio Picon

Alicia Palacios

OSU

Lou Dimauro

LBNL

Oliver Gessner

Daniel Slaughter

Thorsten Weber

Argonne NL

Gilles Doumy

Linda Young

KSU

Daniel Rolles

Artem Rudenko

Univ. Conn.

Nora Berrah

LSU

Ken Lopata

Tohoku Univ.

Kiyoshi Ueda

PSI

Christoph Bostedt

Andre Al-Haddad

Funding:

DOE Basic Energy Sciences
Accelerator and Detector
Research Program

SLAC LDRD and PD programs
Panofsky and Siemann
Fellowships



U.S. DEPARTMENT OF
ENERGY

Office of
Science

Basic Energy
Sciences

Attosecond Core Group at LCLS

Principal Investigators




A. Marinelli



J. Cryan



Current Team



Z. Zhang



D. Cesar



N. Sudar



V. Guo



P. Franz



R. Robles




K. Larsen



E. Thierstein



T. Driver



J. Wang



E. Isele



S. O'Tool

Alumni




Z. Guo



A. Wang



J. O'Neal



Siqi Li



J. Duris




S. Beauvarlet
U. Conn

Laser Group




R. Lemons

X-ray Free-Electron Lasers




Free-Electron Laser





$$\lambda_r = \lambda_u \left(\frac{1}{\beta_z} - 1 \right)$$

The X-ray Free-Electron Laser



X-ray Free-Electron Lasers







Attosecond Electron Dynamics

Attosecond science =
Study of electron dynamics


First step in chemistry



Radiation-matter interaction



Multi-body quantum mechanics




Attosecond Electron Dynamics

First step in chemistry


Attosec
Study of e

$$\hat{H}_{\text{mol}} \Psi(\mathbf{r}\sigma, \mathbf{R}\Sigma, t) = i\hbar \frac{\partial}{\partial t} \Psi(\mathbf{r}\sigma, \mathbf{R}\Sigma, t)$$



$$\begin{aligned} \hat{H}_{\text{mol}} &= \hat{T}_{\text{nuc}} + \hat{V}_{\text{nuc-nuc}} + \hat{T}_{\text{el}} + \hat{V}_{\text{el-el}} + \hat{V}_{\text{nuc-el}} \\ &= \sum_{\alpha=1}^N \frac{\hat{P}_{\alpha}^2}{2M_{\alpha}} + \frac{1}{2} \sum_{\alpha,\beta}^{N,N} \frac{Z_{\alpha}Z_{\beta}e^2}{|\mathbf{R}_{\alpha} - \mathbf{R}_{\beta}|} \\ &\quad + \sum_{i=1}^n \frac{\hat{p}_i^2}{2m} + \frac{1}{2} \sum_{i,j}^{n,n} \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|} \\ &\quad - \sum_{\alpha=1}^N \sum_{i=1}^n \frac{Z_{\alpha}e^2}{|\mathbf{r}_i - \mathbf{R}_{\alpha}|} \end{aligned}$$



Radiation



mechanics




Attosecond Electron Dynamics

First step in chemistry


Attosec
Study of e

$$\hat{H}_{\text{mol}} \Psi(\mathbf{r}\sigma, \mathbf{R}\Sigma, t) = i\hbar \frac{\partial}{\partial t} \Psi(\mathbf{r}\sigma, \mathbf{R}\Sigma, t)$$



$$\begin{aligned} \hat{H}_{\text{mol}} &= \hat{T}_{\text{nuc}} + \hat{V}_{\text{nuc-nuc}} + \hat{T}_{\text{el}} + \hat{V}_{\text{el-el}} + \hat{V}_{\text{nuc-el}} \\ &= \sum_{\alpha=1}^N \frac{\hat{P}_{\alpha}^2}{2M_{\alpha}} + \frac{1}{2} \sum_{\alpha,\beta}^{N,N} \frac{Z_{\alpha}Z_{\beta}e^2}{|\mathbf{R}_{\alpha} - \mathbf{R}_{\beta}|} \\ &\quad + \sum_{i=1}^n \frac{\hat{p}_i^2}{2m} + \frac{1}{2} \sum_{i,j}^{n,n} \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|} \\ &\quad - \sum_{\alpha=1}^N \sum_{i=1}^n \frac{Z_{\alpha}e^2}{|\mathbf{r}_i - \mathbf{R}_{\alpha}|} \end{aligned}$$




Radiation



mechanics



Attosecond X-ray FELs



Three Orders of Magnitude Improvement in Time Resolution

Time Resolution

Early Days
2010

Time sorting
2013-14

2-Color FELs
2013-16

Attosecond FELs
2020-ongoing


1 ps

100 fs


10 fs

1 fs

0.1 fs




J. M. Glownia., et al *Optics express*
18.17 (2010): 17620-17630.




T. Driver et al. arXiv:2411.01700



SASE Spikes




Attosecond FELs



Attosecond FELs




Attosecond FELs




E-Beam Shaping

Self-modulation




J. Duris, S. Li et al. *Nat. Photonics* 14.1 (2020): 30-36.

Laser modulation




J. Duris et al. *Phys. Rev. Lett.* 126, 104802 (2021)

Seeded microbunching instability




Zhang, Zhen, et al. *New Journal of Physics* 22.8 (2020): 083030.
S. Li et al. *APL* 125.19 (2024).

Non-linear compression



S. Huang et al. *PRL* 119, 154801 (2017)
E. Prat et al. *APL Photonics* 8, 111302 (2023)
Yan et al. *Nat. Photonics* 18.12 (2024): 1293-1298.

Typical Results




900 eV


$\langle \text{BW} \rangle = 7.1 \text{ eV}$

$\langle E_p \rangle = 10 \text{ uJ}$

Typical Results




900 eV
<BW> = 7.1 eV
<Ep> = 10 μ J




10 keV, unpublished
<BW> = 30 eV
<Ep> = 20 μ J


Typical Results



900 eV
<BW> = 7.1 eV
<Ep> = 10 uJ




10 keV, unpublished
<BW> = 30 eV
<Ep> = 20 uJ




*K. Li et al. Communications
Physics* **volume 5**, 191 (2022)

THE PULSE LENGTH HAS NO UPPER LIMIT!
THE PULSE LENGTH HAS NO UPPER LIMIT!
THE PULSE LENGTH HAS NO UPPER LIMIT!
THE PULSE LENGTH HAS NO UPPER LIMIT!
THE PULSE LENGTH HAS NO UPPER LIMIT!
THE PULSE LENGTH HAS NO UPPER LIMIT!
THE PULSE LENGTH HAS NO UPPER LIMIT!



Time-Domain Measurements

Time-Domain Measurements



Original concept

P. Eckle et al. *Nature Physics* 4.7 (2008): 565-570.

Application to femtosecond XFELs

N. Hartmann



Nature Photonics 12.4 (2018): 215-220.

Application to attosecond XFELs:



S. Li et al. *Optics Express* 26 4531 (2018)

J. Duris, S. Li et al. *Nat. Photonics* 14.1 (2020): 30-36.



Time-Domain Measurements





Time-Domain Measurements




Time-Domain Measurements






Time-Domain Measurements



Raw Data



Raw Data




J. Duris, S. Li et al. *Nature Photonics* 14.1 (2020): 30-36.

S. Li et al. *Optics Express* Vol. 26, [Issue 4](#), pp. 4531-4547 (2018)


First Experimental Evidence of Sub-Fs Hard X-ray Pulses

I. Inoue, R. Robles,
A. Halavanau ... A. Marinelli
arXiv preprint arXiv:2506.07968


Under Review, Nature Photonics



Comparison with 3-D Model of ASE
AND Literature value of ASE threshold
Pulse Duration \sim 100-400 as FWHM



Attosecond Autocorrelation Measurement



Yanwen Sun et al. arXiv:2503.00000




130 as FWHM
40-50 uJ pulse energy
30 eV Bandwidth

Advanced Attosecond Modes

TW-scale Pulses


P. Franz et al. *Nat. Photonics* 18.7 (2024): 698-703



$$\frac{d^2 b}{dx^2} - b + 3b|b|^2 = 0$$



R. Robles et al. *Phys. Rev. Research* 6, 033158

Attosecond Pulse Pairs



Guo, Zhaoheng, et al. *Nat. Photon.* (2024): 1-7.

Pulse shaping



R. Robles et al. *Phys. Rev. Lett.* **134**, 115001 (2025)

Soliton-Like Superradiance

The solution of eqs. (28) does not depend on z_1 and δ , but only on the initial conditions in $y = 0$: $A_1(0) = \langle \exp(-i\theta_{10}) \rangle = b_0$, $p_{1y}(0) = 0$. It follows that $|A|^2 \propto z_1^2 \propto \rho^2$; since $\rho \propto N^{1/3}$ and $|E|^2 \propto \rho N |A|^2$, then $|E|^2 \propto N \rho^3 \propto N^2$, and the peak intensity scales as the square of the electron current; the width of the pulse scales as $z_1^{-1/2}$, i.e. as the inverse of the square root of the peak field. Furthermore, the factor $\exp(i\delta z_1)$ moves the signal to the resonance: in fact, from eq. (2), $E \exp[ik(z - ct)] \propto A \exp[ik(z - ct) - i\delta z_1] \propto A_1 \exp[ik(z - ct) - i\delta z_2]$, where $z_1 + z_2 = \bar{z}$ has been used. From eqs. (5, 7) it follows that $\delta z_2 = (k_s - k)(ct - z)$, so that $E \propto \exp[ik_s(z - ct)]$.

Setting $x \equiv y^2$, eq. (28c) reads

$$(d/dx)(xA_1) = \langle \exp(-i\theta_1) \rangle \equiv \bar{b}_1,$$

so that by integration

$$A_1 = \frac{1}{x} \int_0^x dx' \bar{b}_1(x') \equiv \bar{b}_1, \quad (29)$$

where \bar{b}_1 is just the average of the bunching parameter with respect to x . Eq. (29) is a generalization of the adiabatic expression $A_1 = b_1$ of ref. [4].

In fig. 2 the intensity $|A_1|^2$, from the numerical integration of eqs. (28), is plotted vs y (dashed line); the result is compared with the output (continuous line) of the numerical code [5], integrating the system (9) in the short-pulse limit (see the captions for the parameters): $|A|^2$ is divided by z_1^2 and plotted vs $y = \sqrt{z_1}(\bar{z} - z_1)$, for $z_1 = 1/K$ (at the pulse head). The agreement turns out to be very satisfactory for nearly the first three peaks of the intensity, demonstrating the validity of the similarity transformation (27). It is interesting that the first pulse of the intensity in fig. 2 is very well described by the expression

$$\begin{aligned} |A_1|^2 &= \frac{|b_0|^2}{3\pi} \frac{\exp[3\sqrt{3}(y/2)^{2/3}]}{\left\{ y + (|b_0|^2/12\pi) \exp[3\sqrt{3}(y/2)^{2/3}] \right\}^2} \\ &= \frac{1}{y} \operatorname{Sech}^2 \left[\frac{3\sqrt{3}}{2} \left(\frac{y}{2} \right)^{2/3} + \log \left(\frac{|b_0|}{2\sqrt{3\pi y}} \right) \right]. \quad (30) \end{aligned}$$

In the next section we will discuss the validity of the similarity equations (28) also for the long-pulse case.

Superradiant self-similar solution:


$$P \sim z^2$$

$$\Delta t \sim z^{1/2}$$


In practice pulse shrinking effect saturates due to diffraction...

Robles et al. (2024)


Superradiance




Superradiance




Superradiance




Superradiance




Superradiance




Superradiance




Superradiance




Superradiance




Superradiance



Superradiance







Superradiance




Cascaded Amplification





P. Franz




TW-Scale Pulses



Is This Superradiance?



R. Robles




Paris Franz et al. "Nature Photonics" 18.7 (2024): 698-703.

R. Robles et al. Phys. Rev. Research 6, 033158 (2024)

See also results from FERMI: Mirian et al. Nature Photonics 15.7 (2021): 523-529


Attosecond Science at LCLS

Electronic Coherence in Auger-Meitner Decay




S. Li, T. Driver et al. *Science* 375.6578 (2022): 285-290.

Photoemission delays




T. Driver et al. *Nature* 632 (8026), 762-767

Non-linear methods






J. O'Neal et al. *Phys. Rev. Lett.* 125.7 (2020): 073203

Pump/Probe Experiments




Z. Guo et al. *Nature Photonics* (2024): 1-7.

Attosecond Pump/Probe Setup



Pump/Probe Demonstration




$t = 0.00$ fs




iso = 0.010

Pump-Probe Scheme


Pump < 280 eV




Pump-Probe Scheme



Pump-Probe Scheme




Pump-Probe Scheme




Pump/Probe Demonstration

T. Driver et al. arXiv:2411.01700



Analysis:


T. Driver, Z. Guo E. Isele
(SLAC/Stanford)
D. Tuthill (OSU)




Simulations:

M. Ruberti and
V. Averbukh (Imperial College)

Attosecond Science in the LCLS-II Era



LCLS-II




- Superconducting RF accelerator in CW mode
- Up to 1 MHz repetition rate


Currently operating in soft x-ray region.
Ongoing HE upgrade will extend to hard x-rays (2027)

LCLS-II

FEL Pulse Energy vs. Photon Energy




Demonstrated up 93 kHz




energy jitter: $1e-4$,
peak current jitter: 4%,
arrival time jitter: 20 fs.

Attosecond Pulses at LCLS-II




Angular Streaking Measurements

Demonstrated at 33 kHz
User operation at 16 kHz




Recent Results at LCLS-II

Impulsive ionization in para-aminophenol




LCLS: 1-2 hours




LCLS-II: ~1-2 minutes

Stimulated Impulsive Raman Scattering




J. O'Neal et al. *PRL*.
125.7 (2020): 073203

LCLS: 5 days for 1 data set (no time delay scan)




LCLS-II: 1 day, 18 data sets (pump/probe scan)
(Preliminary analysis)


Looking Way Into the Future...



Looking Way Into the Future...




Looking Way Into the Future...





Plasma-Based Attosecond X-ray Pulses

SIMULATION




Plasma-Based Attosecond X-ray Pulses




Single-Period Shaping (Sims at 10-30 nm)


Circular polarization switching



Linear polarization switching



Extreme Chirp




Non-SVEA simulations
Jenny Morgan (SLAC)

Stay Tuned...




Summary


Attosecond or single spike pulses are now a reality at most XFEL facilities!



LCLS-II is up and running.
Attosecond pulses at high rep-rate.



Hard X-ray Attosecond Spectroscopy and Metrology



Path to Single-Cycle Waveform Shaping with Plasma Wakefield Acceleration

