

# X17 Searches at the MAGIX Spectrometer Setup at MESA.

**Contribution** on behalf of the **MAGIX** collaboration

Sebastian Stengel (sestenge@uni-mainz.de)

Institute for Nuclear Physics, Johannes Gutenberg University Mainz, Germany

„X17 What if?“ Workshop, Frascati, Italy, March 20th, 2026

<https://agenda.infn.it/event/49565/>

# X17 - to peak or not to peak?

PRL 116, 042501 (2016) PHYSICAL REVIEW LETTERS week ending 29 JANUARY 2016

## Observation of Anomalous Internal Pair Creation in $^8\text{Be}$ : A Possible Indication of a Light, Neutral Boson


A. J. Krasznahorkay,<sup>\*</sup> M. Csatlós, L. Csige, Z. Gácsi, J. Gulyás, M. Hunyadi, I. Kuti, B. M. Nyakó, L. Stuhl, J. Timár, T. G. Tornyai, and Zs. Vajta  
Institute for Nuclear Research, Hungarian Academy of Sciences (MTA Atomki), P.O. Box 51, H-4001 Debrecen, Hungary

T. J. Ketel  
Nikhef National Institute for Subatomic Physics, Science Park 105, 1098 XG Amsterdam, Netherlands

A. Krasznahorkay  
CERN, CH-1211 Geneva 23, Switzerland and Institute for Nuclear Research, Hungarian Academy of Sciences (MTA Atomki), P.O. Box 51, H-4001 Debrecen, Hungary  
(Received 7 April 2015; published 26 January 2016)

Electron-positron angular correlations were measured for the isovector magnetic dipole 17.6 MeV ( $J^\pi = 1^+, T = 1$ ) state  $\rightarrow$  ground state ( $J^\pi = 0^+, T = 0$ ) and the isoscalar magnetic dipole 18.15 MeV ( $J^\pi = 1^+, T = 0$ ) state  $\rightarrow$  ground state transitions in  $^8\text{Be}$ . Significant enhancement relative to the internal pair creation was observed at large angles in the angular correlation for the isoscalar transition with a confidence level of  $> 5\sigma$ . This observation could possibly be due to nuclear reaction interference effects or might indicate that, in an intermediate step, a neutral isoscalar particle with a mass of  $16.70 \pm 0.35(\text{stat}) \pm 0.5(\text{syst}) \text{ MeV}/c^2$  and  $J^\pi = 1^+$  was created.

DOI: 10.1103/PhysRevLett.116.042501



- **ATOMKI (2016):** Peak-like anomaly in the angular correlation of the  $e^+e^-$  pairs produced in the decays of excited states of  $^8\text{Be}$  ( $6.8\sigma$ )

PHYSICAL REVIEW C 104, 044003 (2021)

## New anomaly observed in $^4\text{He}$ supports the existence of the hypothetical X17 particle


A. J. Krasznahorkay,<sup>1,\*</sup> M. Csatlós,<sup>1</sup> L. Csige,<sup>1</sup> J. Gulyás,<sup>1</sup> A. Krasznahorkay,<sup>1,†</sup> B. M. Nyakó,<sup>1</sup> I. Rajta,<sup>1</sup> J. Timár,<sup>1</sup> I. Vajda,<sup>1</sup> and N. J. Sas<sup>2</sup>

<sup>1</sup>Institute for Nuclear Research (ATOMKI), P.O. Box 51, H-4001 Debrecen, Hungary  
<sup>2</sup>University of Debrecen, 4010 Debrecen, PO Box 105, Hungary

(Received 27 October 2019; revised 30 June 2021; accepted 6 October 2021; published 18 October 2021)

Angular correlation spectra of  $e^+e^-$  pairs produced in the  $^3\text{H}(p, e^+e^-)^4\text{He}$  nuclear reaction have been studied at  $E_p = 510, 610,$  and  $900 \text{ keV}$  proton energies. The main features of the spectra can be understood by taking into account the internal and external pair creations following the proton capture by  $^3\text{H}$ . However, these processes cannot account for an observed peak around  $115^\circ$  in the angular correlation spectra. This anomalous excess of  $e^+e^-$  pairs can be described by the creation and subsequent decay of a light particle during the direct capture process. The derived mass of the particle is  $m_{X17}c^2 = 16.94 \pm 0.12(\text{stat}) \pm 0.21(\text{syst}) \text{ MeV}$ . According to the mass this is likely the same X17 particle, which we recently suggested [Phys. Rev. Lett. 116, 042501 (2016)] for describing the anomaly observed in the decay of  $^8\text{Be}$ .

DOI: 10.1103/PhysRevC.104.044003



- **ATOMKI (2021):** Observation of a similar peak-like anomaly in  $^4\text{He}$  ( $\geq 6.6\sigma$ )

PHYSICAL REVIEW C 106, L061601 (2022)

## New anomaly observed in $^{12}\text{C}$ supports the existence and the vector character of the hypothetical X17 boson

A. J. Krasznahorkay,<sup>\*</sup> A. Krasznahorkay,<sup>†</sup> M. Begala, M. Csatlós, L. Csige, J. Gulyás, A. Krakó, J. Timár, I. Rajta, and I. Vajda


Institute for Nuclear Research (ATOMKI), P.O. Box 51, H-4001 Debrecen, Hungary

N. J. Sas  
University of Debrecen, Faculty of Science and Technology, Physics Institute, 4010 Debrecen, PO Box 105, Hungary

(Received 5 November 2022; accepted 5 December 2022; published 12 December 2022)

Employing the  $^{11}\text{B}(p, \gamma)^{12}\text{C}$  nuclear reaction, the angular correlation of  $e^+e^-$  pairs was investigated in the angular range of  $40^\circ \leq \Theta \leq 175^\circ$  for five different proton energies between  $E_p = 1.50$ – $2.5 \text{ MeV}$ . At small angles ( $\Theta \leq 120^\circ$ ), the results can be well interpreted by the internal pair creation process of electromagnetic radiations with  $E1$  and  $M1$  multipoles and by the external pair creation in the target backing. However, at angles greater than  $120^\circ$ , additional count excesses and anomalies were observed, which could be well accounted for by the existence of the previously suggested hypothetical X17 particle. Our results suggest that the X17 particle was generated mainly in  $E1$  radiation. The derived mass of the particle is  $m_{X17}c^2 = 17.03 \pm 0.11(\text{stat}) \pm 0.20(\text{syst}) \text{ MeV}$ . According to the mass, and to the derived branching ratio [ $B_r = 3.6(3) \times 10^{-6}$ ], this is likely the same X17 particle that we recently suggested for describing the anomaly observed in the decay of  $^8\text{Be}$  and  $^4\text{He}$ .

DOI: 10.1103/PhysRevC.106.L061601



- **ATOMKI (2022):** Observation of a similar peak-like anomaly in  $^{12}\text{C}$

# X17 - to peak or not to peak?

Abamyam et al., 10.1134/S1063779624700412

ISSN 1063-7796, Physics of Particles and Nuclei, 2024, Vol. 55, No. 4, pp. 868–873. © Pleiades Publishing, Ltd., 2024.

## Observation of Structures at $\sim 17$ and $\sim 38$ MeV/c<sup>2</sup> in the $\gamma\gamma$ Invariant Mass Spectrum in $d$ Cu Collisions at a Momentum of 3.8 GeV/c per Nucleon

Kh. U. Abraamyam<sup>a, b, \*</sup>, Ch. Austin<sup>c</sup>, M. I. Baznat<sup>d</sup>, K. K. Gudima<sup>e</sup>, M. A. Kozhin<sup>a</sup>, S. G. Reznikov<sup>f</sup>, and A. S. Sorin<sup>a, g</sup>

<sup>a</sup> Veksler and Baldin Laboratory of High Energy Physics, JINR, Dubna, Moscow oblast, 141980 Russia

<sup>b</sup> International Center for Advanced Studies, Yerevan, YSU, 0025 Armenia

<sup>c</sup> 33 Collins Terrace, Maryport, Cumbria CA15 8DL, England

<sup>d</sup> Institute of Applied Physics, Kishinev, MD-2028 Moldova


<sup>e</sup> Bogolubov Laboratory of Theoretical Physics, JINR, Dubna, Moscow oblast, 141980 Russia

\*e-mail: abraamyam@jinr.ru

Received November 30, 2023; revised December 10, 2023; accepted December 14, 2023

**Abstract**—The results of an analysis of the invariant mass spectra of photon pairs produced in  $d$ Cu interactions at a momentum of 3.83 GeV/c per nucleon, are presented. Signals in the form of enhanced structures at invariant masses of about 17 and 38 MeV/c<sup>2</sup> are observed. The results of testing of the observed signals, including the results of the Monte Carlo simulation are presented. The test results support the conclusion that the observed signals are the consequence of detection of the particles with masses of about 17 and 38 MeV/c<sup>2</sup> decaying into a pair of photons.

DOI: 10.1134/S1063779624700412



- JINR (2024): Peak-like anomaly in the  $\gamma\gamma$  invariant mass spectra in  $d$ +Cu collisions ( $\geq 6\sigma$ )

Tran The Anh et al., 10.3390/universe10040168

arXiv:2401.11676v2 [nucl-ex] 19 Mar 2024



## Checking the $^8\text{Be}$ anomaly with a two-arm electron positron pair spectrometer

Tran The Anh<sup>1</sup>, Tran Dinh Trong<sup>2</sup>, Áttila J. Krassnavorháy<sup>3</sup>, Áttila Krassnavorháy<sup>3</sup>, József Molnár<sup>3</sup>, Zoltán Pintye<sup>3</sup>, Nguyen Ai Viet<sup>4</sup>, Nguyen The Nghia<sup>5</sup>, Do Thi Khanh Linh<sup>6</sup>, Bui Thi Hoa<sup>6</sup>, Le Xuan Chung<sup>6</sup> and Nguyen Tuan Anh<sup>6</sup>

<sup>1</sup> VNU-University of Science, Vietnam National University, 334 Nguyen Trãi, Hanoi, Vietnam;

<sup>2</sup> Institute of Physics, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Hanoi, Vietnam;


<sup>3</sup> Institute for Nuclear Research (HUN-REN ATOMKI), P.O. Box 51, H-4001 Debrecen, Hungary

<sup>4</sup> Institute for Nuclear Science and Technology,

VINATOM, 179 Nghia Do, Hanoi, Vietnam;

<sup>5</sup> Hanoi Irradiation Center, VINATOM, Cau Dien, Hanoi, Vietnam

March 20, 2024



- VNU (2024): Repetition of the  $^8\text{Be}$  measurement  
-> Observation of a similar peak-like anomaly ( $\geq 4\sigma$ )

Eur. Phys. J. C manuscript No. (will be inserted by the editor)

## Search for the X17 particle in $^7\text{Li}(p, e^+e^-)^8\text{Be}$ processes with the MEG II detector.

The MEG II collaboration

K. Afanasyev<sup>1</sup>, A. M. Baldini<sup>2a</sup>, S. Ban<sup>3</sup>, H. Benmansour<sup>3ab</sup>, G. Boca<sup>3ab</sup>, P. W. Cattaneo<sup>3a</sup>, G. Cavoto<sup>3ab</sup>, F. Cel<sup>3ab</sup>, M. Chiappini<sup>3ab</sup>, A. Corvaglia<sup>3a</sup>, G. Dal Maso<sup>3a</sup>, A. De Bari<sup>3a</sup>, M. De Gerone<sup>3a</sup>, L. Ferrari Barusso<sup>3ab</sup>, M. Francesconi<sup>3a</sup>, L. Galli<sup>3a</sup>, G. Gobbi<sup>3a</sup>, E. Gotti<sup>3a</sup>, L. Gerolamo<sup>3a</sup>, F. Grancagnolo<sup>3a</sup>, E. G. Grandoni<sup>3a</sup>, M. Grassi<sup>3a</sup>, D. N. Grigoriev<sup>3a,3b</sup>, M. Hildebrandt<sup>3a</sup>, F. Ignatov<sup>3a</sup>, K. Ikeda<sup>3a</sup>, T. Iwamoto<sup>3a</sup>, S. Karpov<sup>3a,3b</sup>, P.-R. Kettle<sup>3a</sup>, N. Khomutov<sup>3a</sup>, A. Kolesnikov<sup>3a</sup>, N. Kravchuk<sup>3a</sup>, V. Krylov<sup>3a</sup>, N. Kuchimskiy<sup>3a</sup>, E. Leonetti<sup>3a</sup>, W. Li<sup>3a</sup>, V. Malyshev<sup>3a</sup>, A. Matsumita<sup>3a</sup>, M. Musco<sup>3a</sup>, S. Mihara<sup>3a</sup>, W. Molzon<sup>3a</sup>, T. Mori<sup>3a</sup>, D. Nicolò<sup>3a</sup>, H. Nishiguchi<sup>3a</sup>, A. Ochi<sup>3a</sup>, W. Ostani<sup>3a</sup>, A. Oya<sup>3a</sup>, D. Palo<sup>3a</sup>, M. Panarero<sup>3a</sup>, A. Papa<sup>3a</sup>, V. Pattinacci<sup>3a</sup>, A. Popov<sup>3a,3b</sup>, F. Renga<sup>3a</sup>, S. Ritt<sup>3a</sup>, M. Rossella<sup>3a</sup>, A. Rozhdrevitsky<sup>3a</sup>, S. Scarpellini<sup>3a</sup>, F. Schwendtmann<sup>3a</sup>, G. Signorelli<sup>3a</sup>, M. Takahashi<sup>3a</sup>, Y. Uchiyama<sup>3a</sup>, A. Venturini<sup>3a</sup>, B. Vitelli<sup>3a</sup>, C. Voena<sup>3a</sup>, K. Yamamoto<sup>3a</sup>, R. Yokota<sup>3a</sup>, T. Yonemoto<sup>3a</sup>

<sup>1</sup>Institute for Nuclear Research, 141980 Dubna, Russia

<sup>2a</sup>INFN Sezione di Pisa, Dipartimento di Fisica dell'Università, Largo B. Pontecorvo 3, 56127 Pisa, Italy

<sup>2b</sup>ICPP, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>3a</sup>INFN Sezione di Genova, Dipartimento di Fisica dell'Università, Via Dodecaneso 31, 16146 Genova, Italy

<sup>3b</sup>INFN Sezione di Pavia, Dipartimento di Fisica dell'Università, Via Bassi 6, 27100 Pavia, Italy

<sup>3c</sup>INFN Sezione di Roma, Dipartimento di Fisica dell'Università "Sapienza", Piazzale A. Moro, 00185 Roma, Italy

<sup>3d</sup>INFN Sezione di Lecce, Dipartimento di Matematica e Fisica dell'Università del Salento, Via per Arnesano, 73100 Lecce, Italy

<sup>3e</sup>Paul Scherrer Institute PSI, 5232 Villigen, Switzerland

<sup>3f</sup>Institute for Particle Physics and Astrophysics, ETH Zürich, Otto-Stern-Weg 5, 8092 Zürich, Switzerland

<sup>3g</sup>INFN Sezione di Napoli, Via Cintia, 80128 Napoli, Italy

<sup>3h</sup>Bukhar Institute of Nuclear Physics of Siberian Academy of Sciences, 630090 Novosibirsk, Russia

<sup>3i</sup>Novosibirsk State Technical University, 630092 Novosibirsk, Russia

<sup>3j</sup>Novosibirsk State University, 630090 Novosibirsk, Russia


<sup>3k</sup>Osaka Lodge Laboratory, University of Liverpool, Liverpool, L69 7ZE, United Kingdom

<sup>3l</sup>University of California, Irvine, CA 92697, USA

<sup>3m</sup>KRIK, High Energy Accelerator Research Organization, 1-1 Oho, Tsukuba, Ibaraki 305-0801, Japan

<sup>3n</sup>Kobe University, 1-1 Rokko-cho, Nada-ku, Kobe, Hyogo 651-8501, Japan

Received: date / Accepted: date



- MEG II (2024): Repetition of the  $^8\text{Be}$  measurement  
-> No peak-like anomaly but results compatible within  $1.5\sigma$

The MEG II collaboration, 10.48550/arXiv:2411.07994

# X17 - to peak or not to peak?

The PADME collaboration, 10.1007/JHEP11(2025)29007

PREPARED FOR SUBMISSION TO JHEP


## Search for a new 17 MeV resonance via $e^+e^-$ annihilation with the PADME Experiment

The PADME Collaboration  
 F. Bossi<sup>a</sup> R. De Sangro<sup>c</sup> C. Di Giulio<sup>a</sup> E. Di Meo<sup>a</sup> D. Domenici<sup>a</sup> G. Finocchiaro<sup>a</sup>  
 L.G. Foggetta<sup>a</sup> M. Garattini<sup>b</sup> P. Gianotti<sup>b</sup> M. Mancini<sup>b</sup> I. Sarrà<sup>b</sup> T. Spadaro<sup>b,c</sup> C.  
 Taruggi<sup>b</sup> E. Vitucchi<sup>b</sup> K. Dimitrova<sup>d</sup> S. Ivanov<sup>d</sup> Sv. Ivanov<sup>d</sup> K. Kostova<sup>d</sup> V.  
 Kozhuharov<sup>d</sup> R. Simuzonov<sup>d</sup> F. Ferrarotto<sup>e</sup> E. Leonardi<sup>e</sup> P. Valente<sup>e</sup> E. Longo<sup>d,f</sup> G.C.  
 Organtini<sup>d</sup> M. Raggi<sup>d</sup> A. Frankenthal<sup>g</sup>

<sup>a</sup>INFN Laboratori Nazionali di Frascati, Via E. Fermi, 54 I-00044 Frascati, Italy  
<sup>b</sup>Faculty of Physics, Sofia University "St. Kl. Ohridski", 5 J. Bourchier Blvd., BG-1164 Sofia, Bulgaria  
<sup>c</sup>INFN Sezione di Roma, p.le Aldo Moro 5, I-00185 Rome, Italy  
<sup>d</sup>Physics Department, "Sapienza" Università di Roma, p.le Aldo Moro 5, I-00185 Rome, Italy  
<sup>e</sup>Department of Physics and Astronomy, University of California, Irvine, Irvine, CA 92697-1575, USA  
<sup>f</sup>INFN Laboratori Nazionali di Frascati, Via E. Fermi, 54 I-00044 Frascati, Italy  
<sup>g</sup>Department of Physics and Astronomy, University of California, Irvine, Irvine, CA 92697-1575, USA

ABSTRACT: The PADME Experiment at the Frascati DAΦNE linear accelerator has searched for a hypothetical particle with mass around 17 MeV, commonly referred to as the X17, using a positron beam incident on a fixed target. The beam energy was varied between 202 and 296 MeV, corresponding to center-of-mass energies  $\sqrt{s}$  between 16.4 and 17.4 MeV. The X17 should be produced resonantly via  $e^+e^-$  annihilation when  $\sqrt{s}$  approaches its mass, inducing an excess of events with a two-body final state over the background expectation. The beam energy spacing was fixed to less than half the expected width of the resonance's line shape. Uncertainties below 1% per  $\sqrt{s}$  point were achieved. A blind analysis has been performed. The data are consistent with the expected background in most of the explored energy range, and limits are set in previously unexplored regions of the available parameter space. The most significant deviation is found for  $\sqrt{s} \approx 16.90$  MeV, corresponding to a global significance of approximately 2 standard deviations over the null hypothesis expectation.

ARXIV EPRINT: 2505.24797



- PADME (2025): Peak-like anomaly in the invariant mass spectrum of  $e^+e^-$  annihilation ( $1.7\sigma$ )

# X17 - to peak or not to peak?

The PADME collaboration, 10.1107/JHEP11%282025%29007

PREPARED FOR SUBMISSION TO JHEP


## Search for a new 17 MeV resonance via $e^+e^-$ annihilation with the PADME Experiment

The PADME Collaboration  
F. Bossi<sup>a</sup>, R. De Sangro<sup>a</sup>, C. Di Giulio<sup>a</sup>, E. Di Meo<sup>a</sup>, D. Domenici<sup>a</sup>, G. Finocchiaro<sup>a</sup>,  
L.G. Foggetta<sup>a</sup>, M. Garattini<sup>a</sup>, P. Gianotti<sup>a</sup>, M. Mancini<sup>a</sup>, I. Sarrà<sup>a</sup>, T. Spadaro<sup>a</sup>, C.  
Taruggi<sup>a</sup>, E. Vitucchi<sup>a</sup>, K. Dimitrova<sup>b</sup>, S. Ivanov<sup>b</sup>, Sv. Ivanov<sup>b</sup>, K. Kostova<sup>b</sup>, V.  
Kozhuharov<sup>b</sup>, R. Simicic<sup>c</sup>, F. Ferrarotto<sup>d</sup>, E. Leonardi<sup>d</sup>, P. Valente<sup>d</sup>, E. Longo<sup>d</sup>, G.C.  
Organtini<sup>d</sup>, M. Raggi<sup>d</sup>, A. Frankenthal<sup>e</sup>

<sup>a</sup>INFN Laboratori Nazionali di Frascati, Via E. Fermi, 54 I-00044 Frascati, Italy  
<sup>b</sup>Faculty of Physics, Sofia University "St. Kl. Ohridski", 5 J. Bourchier Blvd., BG-1164 Sofia, Bulgaria  
<sup>c</sup>INFN Sezione di Roma, p.le Aldo Moro 5, I-00185 Rome, Italy  
<sup>d</sup>Physics Department, "Sapienza" Università di Roma, p.le Aldo Moro 5, I-00185 Rome, Italy  
<sup>e</sup>Department of Physics and Astronomy, University of California, Irvine, Irvine, CA 92697-1575, USA

ABSTRACT: The PADME Experiment at the Frascati DAΦNE linear accelerator has searched for a hypothetical particle with mass around 17 MeV, commonly referred to as the X17, using a positron beam incident on a fixed target. The beam energy was varied between 202 and 296 MeV, corresponding to center-of-mass energies  $\sqrt{s}$  between 16.4 and 17.4 MeV. The X17 should be produced resonantly via  $e^+e^-$  annihilation when  $\sqrt{s}$  approaches its mass, inducing an excess of events with a two-body final state over the background expectation. The beam energy spacing was fixed to less than half the expected width of the resonance's line shape. Uncertainties below 1% per  $\sqrt{s}$  point were achieved. A blind analysis has been performed. The data are consistent with the expected background in most of the explored energy range, and limits are set in previously unexplored regions of the available parameter space. The most significant deviation is found for  $\sqrt{s} \approx 16.90$  MeV, corresponding to a global significance of approximately 2 standard deviations over the null hypothesis expectation.



ARXIV EPRINT: 2506.24797



- PADME (2025): Peak-like anomaly in the invariant mass spectrum of  $e^+e^-$  annihilation ( $1.7\sigma$ )




**MORE DATA WOULD BE NICE!**

# X17 - to peak or not to peak?





- PADME (2025): Peak-like anomaly in the invariant mass spectrum of  $e^+e^-$  annihilation ( $1.7\sigma$ )

**MORE DATA WOULD BE NICE!**






# X17 - to peak or not to peak?



- PADME (2025): Peak-like anomaly in the invariant mass spectrum of  $e^+e^-$  annihilation ( $1.7\sigma$ )

**MORE DATA WOULD BE NICE!**



?

?

?

?

?

?

**MAGIX**

?

?

?

?

?

?

# MAGIX

**MAinz**

MESA

Multi-purpose Apparatus

Massima Accuratezza

# Greetings from Mainz.

Carnival



bistummainz.de



pixopolis.de

Printing & Johannes Gutenberg



Adobe stock



Wine








christophorus-hof.de

mainz-tourismus.com




# Greetings from MAMI.




- **MA**inzer **MI**crotron, energies  $\leq 1600 \text{ MeV}$ , currents  $\geq 100 \mu\text{A}$
- Mainz has a [long history of electron accelerators](#) (linac in the 1960s, MAMI since the 1990s)

# Dark sector searches at MAMI.



Bachelor's thesis S. Merkel



PRL 106, 251802 (2011) PHYSICAL REVIEW LETTERS week ending 24 JUNE 2011

## Search for Light Gauge Bosons of the Dark Sector at the Mainz Microtron

H. Merkel,<sup>1,4</sup> P. Achenbach,<sup>1</sup> C. Ayerbe Gayoso,<sup>1</sup> J. C. Bernauer,<sup>1,5</sup> R. Böhm,<sup>1</sup> D. Bosnar,<sup>2</sup> L. Debenjak,<sup>2</sup> A. Denig,<sup>1</sup> M. O. Distler,<sup>1</sup> A. Esser,<sup>1</sup> H. Fonvielle,<sup>1</sup> I. Frišić,<sup>2</sup> D. G. Middleton,<sup>1</sup> U. Müller,<sup>1</sup> L. Nungesser,<sup>1</sup> J. Pochodzalla,<sup>1</sup> M. Rohrbeck,<sup>1</sup> S. Sánchez Majos,<sup>1</sup> B. S. Schlimme,<sup>1</sup> M. Schoth,<sup>1</sup> S. Širca,<sup>3,5</sup> and M. Weinriefer<sup>1</sup>

(A1 Collaboration)

<sup>1</sup>Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany

<sup>2</sup>Department of Physics, University of Zagreb, HR-10002 Zagreb, Croatia

<sup>3</sup>Jozef Stefan Institute, SI-1000 Ljubljana, Slovenia

<sup>4</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, BP 10448, F-63000 Clermont-Ferrand, France

<sup>5</sup>Department of Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia

(Received 21 January 2011; published 22 June 2011)

A new exclusion limit for the electromagnetic production of a light  $U(1)$  gauge boson  $\gamma'$  decaying to  $e^+e^-$  was determined by the A1 Collaboration at the Mainz Microtron. Such light gauge bosons appear in

several extensions of the standard model of particle physics. In this Letter, the search for the signal of such a  $U(1)$  gauge boson in electron-positron pair production at the spectrometer setup of the A1 Collaboration at the Mainz Microtron is described. Exclusion limits in the mass range of  $40 \text{ MeV}/c^2$  to  $300 \text{ MeV}/c^2$ , with a sensitivity in the squared mixing parameter of as little as  $\epsilon^2 = 8 \times 10^{-7}$  are presented. A large fraction of the parameter space has been excluded where the discrepancy of the measured anomalous magnetic moment of the muon with theory might be explained by an additional  $U(1)$  gauge boson.

PRL 112, 221802 (2014) PHYSICAL REVIEW LETTERS week ending 6 JUNE 2014

## Search at the Mainz Microtron for Light Massive Gauge Bosons Relevant for the Muon $g - 2$ Anomaly

H. Merkel,<sup>1,4</sup> P. Achenbach,<sup>1</sup> C. Ayerbe Gayoso,<sup>1,4</sup> T. Beranek,<sup>1</sup> J. Beringer,<sup>2</sup> J. C. Bernauer,<sup>1,5</sup> R. Böhm,<sup>1</sup> D. Bosnar,<sup>3</sup> L. Correu,<sup>1</sup> L. Debenjak,<sup>2</sup> A. Denig,<sup>1</sup> M. O. Distler,<sup>1</sup> A. Esser,<sup>1</sup> H. Fonvielle,<sup>1</sup> I. Frišić,<sup>2</sup> M. Gómez Rodríguez de la Paz,<sup>1</sup> M. Hoek,<sup>1</sup> S. Kegeles,<sup>1</sup> Y. Kohl,<sup>1</sup> D. G. Middleton,<sup>1</sup> M. Mihovilović,<sup>1</sup> U. Müller,<sup>1</sup> L. Nungesser,<sup>1</sup> J. Pochodzalla,<sup>1</sup> M. Rohrbeck,<sup>1</sup> G. Ron,<sup>2</sup> S. Sánchez Majos,<sup>1</sup> B. S. Schlimme,<sup>1</sup> M. Schoth,<sup>1</sup> F. Schulz,<sup>1</sup> C. Sifanti,<sup>1</sup> S. Širca,<sup>2,6</sup> M. Thiel,<sup>1</sup> A. Tyukin,<sup>1</sup> A. Weber,<sup>1</sup> and M. Weinriefer<sup>1</sup>

(A1 Collaboration)

<sup>1</sup>Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany

<sup>2</sup>Jozef Stefan Institute, SI-1000 Ljubljana, Slovenia

<sup>3</sup>Department of Physics, University of Zagreb, HR-10002 Zagreb, Croatia

<sup>4</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, BP 10448, F-63000 Clermont-Ferrand, France

<sup>5</sup>Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem 91904, Israel

<sup>6</sup>Department of Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia

(Received 22 April 2014; revised manuscript received 19 May 2014; published 4 June 2014)

A massive, but light, Abelian  $U(1)$  gauge boson is a well-motivated possible signature of physics beyond the standard model of particle physics. In this Letter, the search for the signal of such a  $U(1)$  gauge boson in electron-positron pair production at the spectrometer setup of the A1 Collaboration at the Mainz Microtron is described. Exclusion limits in the mass range of  $40 \text{ MeV}/c^2$  to  $300 \text{ MeV}/c^2$ , with a sensitivity in the squared mixing parameter of as little as  $\epsilon^2 = 8 \times 10^{-7}$  are presented. A large fraction of the parameter space has been excluded where the discrepancy of the measured anomalous magnetic moment of the muon with theory might be explained by an additional  $U(1)$  gauge boson.


DOI: 10.1103/PhysRevLett.112.221802

PACS numbers: 14.70.Pw, 13.40.Ea, 25.30.Rw, 95.35.+d


Merkel et al., 10.1103/PhysRevLett.112.221802

- Searches for Light Gauge Bosons of a hypothetical dark sector in 2011 and 2014 giving exclusion limits in the mass range from  $40 \text{ MeV}/c^2$  to  $300 \text{ MeV}/c^2$

# Dark sector searches at MAMI.



Bachelor's thesis S. Merkel



PRL 106, 251802 (2011) PHYSICAL REVIEW LETTERS week ending 24 JUNE 2011

**Search for Light Gauge Bosons of the Dark Sector at the Mainz Microtron**

H. Merkel,<sup>1,4</sup> P. Achenbach,<sup>1</sup> C. Ayerbe Gayoso,<sup>1</sup> J. C. Bernauer,<sup>1,2</sup> R. Böhm,<sup>1</sup> D. Bosnar,<sup>2</sup> L. Debenjak,<sup>2</sup> A. Denig,<sup>1</sup> M. O. Distler,<sup>1</sup> A. Esser,<sup>1</sup> H. Fonvielle,<sup>1</sup> I. Frišić,<sup>2</sup> D. G. Middleton,<sup>1</sup> U. Müller,<sup>1</sup> L. Nungesser,<sup>1</sup> J. Pochodzalla,<sup>1</sup> M. Rohrbeck,<sup>1</sup> S. Sánchez Majos,<sup>1</sup> B. S. Schlimme,<sup>1</sup> M. Schoth,<sup>1</sup> S. Širca,<sup>3,5</sup> and M. Weinrieger<sup>1</sup>

(A1 Collaboration)

<sup>1</sup>Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany  
<sup>2</sup>Department of Physics, University of Zagreb, HR-10002 Zagreb, Croatia  
<sup>3</sup>Jozef Stefan Institute, SI-1000 Ljubljana, Slovenia  
<sup>4</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, BP 10448, F-63000 Clermont-Ferrand, France  
<sup>5</sup>Department of Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia

(Received 21 January 2011; published 22 June 2011)

A new exclusion limit for the electromagnetic production of a light  $U(1)$  gauge boson  $\gamma'$  decaying to  $e^+e^-$  was determined by the A1 Collaboration at the Mainz Microtron. Such light gauge bosons appear in several extensions of the standard model of particle physics. In this Letter, the search for the signal of such a  $U(1)$  gauge boson in electron-positron pair production at the spectrometer setup of the A1 Collaboration at the Mainz Microtron is described. Exclusion limits in the mass range of  $40 \text{ MeV}/c^2$  to  $300 \text{ MeV}/c^2$ , with a sensitivity in the squared mixing parameter of as little as  $\epsilon^2 = 8 \times 10^{-7}$  are presented. A large fraction of the parameter space has been excluded where the discrepancy of the measured anomalous magnetic moment of the muon with theory might be explained by an additional  $U(1)$  gauge boson.

DOI: 10.1103/PhysRevLett.106.251802 PACS numbers: 14.70.Pw, 13.40.Ea, 25.30.Rw, 95.35.+d

PRL 112, 221802 (2014) PHYSICAL REVIEW LETTERS week ending 6 JUNE 2014

**Search at the Mainz Microtron for Light Massive Gauge Bosons Relevant for the Muon  $g - 2$  Anomaly**

H. Merkel,<sup>1,4</sup> P. Achenbach,<sup>1</sup> C. Ayerbe Gayoso,<sup>1,2</sup> T. Beranek,<sup>1</sup> J. Beringer,<sup>2</sup> J. C. Bernauer,<sup>1,2</sup> R. Böhm,<sup>1</sup> D. Bosnar,<sup>3</sup> L. Correu,<sup>1</sup> L. Debenjak,<sup>2</sup> A. Denig,<sup>1</sup> M. O. Distler,<sup>1</sup> A. Esser,<sup>1</sup> H. Fonvielle,<sup>1</sup> I. Frišić,<sup>2</sup> M. Gómez Rodríguez de la Paz,<sup>1</sup> M. Hoek,<sup>1</sup> S. Kegeles,<sup>1</sup> Y. Kohl,<sup>1</sup> D. G. Middleton,<sup>1</sup> M. Mihovilović,<sup>1</sup> U. Müller,<sup>1</sup> L. Nungesser,<sup>1</sup> J. Pochodzalla,<sup>1</sup> M. Rohrbeck,<sup>1</sup> G. Ron,<sup>2</sup> S. Sánchez Majos,<sup>1</sup> B. S. Schlimme,<sup>1</sup> M. Schoth,<sup>1</sup> F. Schulz,<sup>1</sup> C. Sifanti,<sup>1</sup> S. Širca,<sup>2,6</sup> M. Thiel,<sup>1</sup> A. Tyukin,<sup>1</sup> A. Weber,<sup>1</sup> and M. Weinrieger<sup>1</sup>

(A1 Collaboration)

<sup>1</sup>Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany  
<sup>2</sup>Jozef Stefan Institute, SI-1000 Ljubljana, Slovenia  
<sup>3</sup>Department of Physics, University of Zagreb, HR-10002 Zagreb, Croatia  
<sup>4</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, BP 10448, F-63000 Clermont-Ferrand, France  
<sup>5</sup>Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem 91904, Israel  
<sup>6</sup>Department of Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia

(Received 22 April 2014; revised manuscript received 19 May 2014; published 4 June 2014)

A massive, but light, Abelian  $U(1)$  gauge boson is a well-motivated possible signature of physics beyond the standard model of particle physics. In this Letter, the search for the signal of such a  $U(1)$  gauge boson in electron-positron pair production at the spectrometer setup of the A1 Collaboration at the Mainz Microtron is described. Exclusion limits in the mass range of  $40 \text{ MeV}/c^2$  to  $300 \text{ MeV}/c^2$ , with a sensitivity in the squared mixing parameter of as little as  $\epsilon^2 = 8 \times 10^{-7}$  are presented. A large fraction of the parameter space has been excluded where the discrepancy of the measured anomalous magnetic moment of the muon with theory might be explained by an additional  $U(1)$  gauge boson.






DOI: 10.1103/PhysRevLett.112.221802 PACS numbers: 14.70.Pw, 13.40.Ea, 25.30.Rw, 95.35.+d

Merkel et al., 10.1103/PhysRevLett.112.221802

- Searches for Light Gauge Bosons of a hypothetical dark sector in 2011 and 2014 giving exclusion limits in the mass range from  $40 \text{ MeV}/c^2$  to  $300 \text{ MeV}/c^2$
- X17 range not accessible at A1@MAMI






Merkel et al., 10.1103/PhysRevLett.106.251802

# With best wishes from A4.



- The A4 experiment has been finished several years ago and its experimental halls are free for something new


# Greetings from MESA.




- The A4 experiment has been finished several years ago and its experimental halls are free for something new
- **MESA will be the next-generation electron accelerator in Mainz** with lower energies but higher intensities compared to MAMI

# The low-energy electron accelerator MESA.


- Mainz **E**nergy-**R**ecovering **S**uperconducting **A**ccelerator
- Double-sided Multi Turn **E**nergy-**R**ecovery **L**inac (ERL)
- Beam energies from *20 MeV to 155 MeV*
- Beam currents of  $\geq 1 \text{ mA}$
- Two operation modes: ERL and Extracted Beam (EB)
- Three experiments: P2, DarkMESA, and **MAGIX**




# Let's build MESA (1/7).



# Let's build MESA (2/7).




# Let's build MESA (3/7).




Extracted Beam (EB) mode - P2 experiment

# Let's build MESA (4/7).





Extracted Beam (EB) mode - MAGIX experiment

# Let's build MESA (5/7).



Energy-Recovery...



# Let's build MESA (6/7).



**Double-sided Multi Turn Energy-Recovery Linac**

Energy-Recovery Linac (ERL) mode - MAGIX experiment


# Let's build MESA (7/7).



**M**ainz **E**nergy-recovering **S**uperconducting **A**ccelerator


# The low-energy electron accelerator MESA.

- Mainz **E**nergy-**R**ecovering **S**uperconducting **A**ccelerator
- Double-sided Multi Turn **E**nergy-**R**ecovery **L**inac (ERL)
- Beam energies from *20 MeV to 155 MeV*
- Beam currents of  $\geq 1 \text{ mA}$
- Two operation modes: ERL and Extracted Beam (EB)
- Three experiments: P2, DarkMESA, and **MAGIX**



# The low-energy electron accelerator MESA.

- Mainz **E**nergy-**R**ecovering **S**uperconducting **A**ccelerator
- Double-sided Multi Turn **E**nergy-**R**ecovery **L**inac (ERL)
- Beam energies from *20 MeV to 155 MeV*
- Beam currents of  $\geq 1 \text{ mA}$
- Two operation modes: ERL and Extracted Beam (EB)
- Three experiments: P2, DarkMESA, and **MAGIX**



## P2 - Parity violating experiments


- Main focus: Measurement of the **electroweak mixing angle** with a precision of 0.15%

# The low-energy electron accelerator MESA.

- Mainz **E**nergy-**R**ecovering **S**uperconducting **A**ccelerator
- Double-sided Multi Turn **E**nergy-**R**ecovery **L**inac (ERL)
- Beam energies from *20 MeV to 155 MeV*
- Beam currents of  $\geq 1 \text{ mA}$
- Two operation modes: ERL and Extracted Beam (EB)
- Three experiments: P2, DarkMESA, and **MAGIX**

## The beam-dump experiment DarkMESA

- Direct search for **light dark matter** (LDM) particles




# The low-energy electron accelerator MESA.

- Mainz **E**nergy-**R**ecovering **S**uperconducting **A**ccelerator
- Double-sided Multi Turn **E**nergy-**R**ecovery **L**inac (ERL)
- Beam energies from *20 MeV to 155 MeV*
- Beam currents of  $\geq 1 \text{ mA}$
- Two operation modes: ERL and Extracted Beam (EB)
- Three experiments: P2, DarkMESA, and **MAGIX**

## The multi-purpose experiment MAGIX

- A compact, next-generation version of the A1 experiment at MAMI
- Mainly operated in MESA's ERL mode
- Rich physics program in nuclear, particle, and hadron physics



# MAGIX


**Gas Injection**

Gas Internal

Gas Interaction

Grande Innovazione

# The basic idea of MAGIX.




**Windowless  
gas jet target**  
and minimal material budget

+

**High-intensity  
electron beam**  
in the low-energy regime

# The basic idea of MAGIX.



**Windowless  
gas jet target**  
and minimal material budget


+

**High-intensity  
electron beam**  
in the low-energy regime



**Clean  
experimental  
environment**  
with drastically reduced particle  
interactions prior to detection

# The basic idea of MAGIX.



**Windowless  
gas jet target**  
and minimal material budget

+

**High-intensity  
electron beam**  
in the low-energy regime




**Clean  
experimental  
environment**  
with drastically reduced particle  
interactions prior to detection

+


**Competitive  
luminosities**  
in the order of  $10^{35} \text{ cm}^{-2}\text{s}^{-1}$

# The MAGIX gas jet target.


D. Bonaventura



- Developed and constructed by AG Khoukaz at WWU Münster, Germany
- Windowless, thin, point-like jet target
- Already commissioned at A1 with hydrogen...



S. Schlimme et al., 10.1016/j.nima.2021.165668



Nuclear Inst. and Methods in Physics Research, A 1013 (2021) 165668

Contents lists available at ScienceDirect

**Nuclear Inst. and Methods in Physics Research, A**

journal homepage: [www.elsevier.com/locate/nima](http://www.elsevier.com/locate/nima)

---

**Operation and characterization of a windowless gas jet target in high-intensity electron beams**

B.S. Schlimme<sup>a,\*</sup>, S. Aulenbacher<sup>a,1</sup>, P. Brand<sup>b,1,2</sup>, M. Littich<sup>a,3</sup>, Y. Wang<sup>1,3</sup>, P. Achenbach<sup>a,4,5</sup>, M. Ball<sup>6</sup>, J.C. Bernauer<sup>b,7</sup>, M. Biroth<sup>8</sup>, D. Bonaventura<sup>b</sup>, D. Bosnar<sup>9</sup>, S. Caiazza<sup>9</sup>, M. Christmann<sup>a,4</sup>, E. Cline<sup>10</sup>, A. Denig<sup>a,11</sup>, M.O. Distler<sup>12</sup>, L. Doria<sup>a,13</sup>, P. Eckert<sup>14</sup>, A. Esser<sup>15</sup>, I. Frišić<sup>16</sup>, S. Gagneur<sup>17</sup>, J. Geimer<sup>18</sup>, S. Griener<sup>19</sup>, P. Gülker<sup>20</sup>, P. Herrmann<sup>21</sup>, M. Hoek<sup>22</sup>, S. Kegel<sup>23</sup>, J. Kelsey<sup>24</sup>, P. Klag<sup>25</sup>, A. Khoukaz<sup>26</sup>, M. Kohl<sup>27</sup>, T. Kolar<sup>28,29</sup>, M. Lauß<sup>30</sup>, L. Lefsmann<sup>31</sup>, S. Lunkenheimer<sup>32</sup>, J. Marekovič<sup>33</sup>, D. Markus<sup>34</sup>, M. Mauch<sup>35</sup>, H. Merkel<sup>36</sup>, M. Mihovilović<sup>37,38</sup>, R.G. Milner<sup>39</sup>, J. Müller<sup>40</sup>, U. Müller<sup>41</sup>, T. Petrovič<sup>42</sup>, J. Pochodzalla<sup>43</sup>, J. Rausch<sup>44</sup>, J. Schlaadt<sup>45</sup>, H. Schürg<sup>46</sup>, C. Sfienti<sup>47</sup>, S. Širca<sup>48</sup>, R. Spreckels<sup>49</sup>, S. Stengel<sup>50</sup>, Y. Stöttinger<sup>51</sup>, C. Szyszka<sup>52</sup>, M. Thiel<sup>53</sup>, S. Vestrick<sup>54</sup>, C. Vidal<sup>55</sup>, for the A1 and MAGIX Collaborations


<sup>a</sup> Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany  
<sup>b</sup> Institut für Kernphysik, Westfälische Wilhelms-Universität, D-48149 Münster, Germany  
<sup>c</sup> PRISMA<sup>+</sup> Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany  
<sup>d</sup> Helmholtz-Institut Mainz, GSI Helmholtzbeschleuniger für Schwerionenforschung, Darmstadt, Johannes Gutenberg-Universität, D-55099 Mainz, Germany  
<sup>e</sup> Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität, D-53115 Bonn, Germany  
<sup>f</sup> Department of Physics, University of Zagreb, HR-10002 Zagreb, Croatia  
<sup>g</sup> Jozef Stefan Institute, SI-1000 Ljubljana, Slovenia  
<sup>h</sup> Faculty of Mathematics and Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia  
<sup>i</sup> Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA  
<sup>j</sup> MIT Bates Research and Engineering Center, Middlesex, MA, 01948, USA  
<sup>k</sup> Center for Precision in Nuclear Science, Department of Physics and Astronomy, Stony Brook University, New York 11794, USA  
<sup>l</sup> RIKEN INFL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA  
<sup>m</sup> Department of Physics, Hampton University, Hampton, Virginia 23068, USA

S. Schlimme et al., 10.1016/j.nima.2021.165668


S. Schlimme et al., 10.1016/j.nima.2021.165668

# The MAGIX gas jet target.

D. Bonaventura



- Developed and constructed by AG Khoukaz at WWU Münster, Germany
- Windowless, thin, point-like jet target
- Already commissioned at A1 with hydrogen and argon



S. Schlimme et al., 10.1016/j.nima.2021.165668

Eur. Phys. J. A (2025) 61:152  
 https://doi.org/10.1140/epja/s10050-025-01623-4

THE EUROPEAN PHYSICAL JOURNAL A

Regular Article - Experimental Physics

### Measurement of the $^{40}\text{Ar}(e,e')$ elastic scattering cross section with a novel gas-jet target

M. Littich<sup>1</sup>, L. Doria<sup>1,4</sup>, P. Brand<sup>2</sup>, P. Achenbach<sup>1</sup>, S. Aulenbacher<sup>1</sup>, S. Bacca<sup>1</sup>, J. C. Bernauer<sup>3</sup>, M. Biroth<sup>1</sup>, D. Bonaventura<sup>2</sup>, D. Bosnar<sup>4</sup>, M. Christmann<sup>1</sup>, E. Cline<sup>3,5</sup>, A. Denig<sup>1</sup>, M. Distler<sup>1</sup>, A. Esser<sup>1</sup>, I. Frišić<sup>6</sup>, J. Geimer<sup>1</sup>, P. Gülker<sup>1</sup>, M. Hoek<sup>1</sup>, P. Klag<sup>1</sup>, A. Khoukaz<sup>2</sup>, M. Laub<sup>1</sup>, S. Lunkenheimer<sup>1</sup>, T. Manoussos<sup>1</sup>, D. Markus<sup>1</sup>, H. Merkel<sup>1</sup>, M. Mihovilović<sup>6,7</sup>, U. Müller<sup>1</sup>, J. Pochodzalla<sup>1</sup>, B. S. Schlimme<sup>1</sup>, C. Sienti<sup>1</sup>, J. E. Sobczyk<sup>1</sup>, S. Stengel<sup>1</sup>, E. Stephan<sup>8</sup>, M. Thiel<sup>1</sup>, S. Vestrick<sup>2</sup>, A. Wilczek<sup>8</sup>, L. Wilhelm<sup>1</sup>

<sup>1</sup> Institut für Kernphysik, Johannes Gutenberg-Universität, 55128 Mainz, Germany  
<sup>2</sup> Institut für Kernphysik, Universität Münster, 48149 Münster, Germany  
<sup>3</sup> Department of Physics and Astronomy, Center for Frontiers in Nuclear Science, Stony Brook University, New York 11794, USA  
<sup>4</sup> Department of Physics, Faculty of Science, University of Zagreb, Zaireb, Croatia  
<sup>5</sup> Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, MA 02139, USA  
<sup>6</sup> Jožef Stefan Institute, 1000 Ljubljana, Slovenia  
<sup>7</sup> Faculty of Mathematics and Physics, University of Ljubljana  
<sup>8</sup> Institute of Physics, University of Silesia in Katowice, 40000 Katowice, Poland



P. Brand

Nuclear Inst. and Methods in Physics Research, A 1013 (2021) 165668

Contents lists available at ScienceDirect

Nuclear Inst. and Methods in Physics Research, A

journal homepage: www.elsevier.com/locate/nima


### Operation and characterization of a windowless gas jet target in high-intensity electron beams

B.S. Schlimme<sup>a,\*</sup>, S. Aulenbacher<sup>a,1</sup>, P. Brand<sup>b,1,2</sup>, M. Littich<sup>a,3</sup>, Y. Wang<sup>1,3</sup>, P. Achenbach<sup>a,4,5</sup>, M. Ball<sup>6</sup>, J.C. Bernauer<sup>b,7</sup>, M. Biroth<sup>8</sup>, D. Bonaventura<sup>b</sup>, D. Bosnar<sup>9</sup>, S. Caiazza<sup>10</sup>, M. Christmann<sup>a,4</sup>, E. Cline<sup>3</sup>, A. Denig<sup>a,11</sup>, M.O. Distler<sup>12</sup>, L. Doria<sup>a,13</sup>, P. Eckert<sup>14</sup>, A. Esser<sup>15</sup>, I. Frišić<sup>16</sup>, S. Gagneur<sup>17</sup>, J. Geimer<sup>18</sup>, S. Grieser<sup>19</sup>, P. Gülker<sup>20</sup>, P. Herrmann<sup>21</sup>, M. Hoek<sup>22</sup>, S. Kegel<sup>23</sup>, J. Kelsey<sup>24</sup>, P. Klag<sup>25</sup>, A. Khoukaz<sup>26</sup>, M. Kohl<sup>27</sup>, T. Kolar<sup>28</sup>, M. Laub<sup>29</sup>, L. Leßmann<sup>30</sup>, S. Lunkenheimer<sup>31</sup>, J. Marekovič<sup>32</sup>, D. Markus<sup>33</sup>, M. Mauch<sup>34</sup>, H. Merkel<sup>35</sup>, M. Mihovilović<sup>36,37</sup>, R.G. Milner<sup>38</sup>, J. Müller<sup>39</sup>, U. Müller<sup>40</sup>, T. Petrovič<sup>41</sup>, J. Pochodzalla<sup>42</sup>, J. Rausch<sup>43</sup>, J. Schlaadt<sup>44</sup>, H. Schürig<sup>45</sup>, C. Sienti<sup>46</sup>, S. Širca<sup>47</sup>, R. Spreckels<sup>48</sup>, S. Stengel<sup>49</sup>, Y. Stöttinger<sup>50</sup>, C. Szyska<sup>51</sup>, M. Thiel<sup>52</sup>, S. Vestrick<sup>53</sup>, C. Vidal<sup>54</sup>, for the A1 and MAGIX Collaborations

<sup>a</sup> Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany  
<sup>b</sup> Institut für Kernphysik, Westfälische Wilhelms-Universität, D-48149 Münster, Germany  
<sup>c</sup> PRISMA<sup>+</sup> Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany  
<sup>d</sup> Helmholtz Institute Mainz, GSI Helmholtzaccelerator für Schwerionenforschung, Darmstadt, Johannes Gutenberg-Universität, D-55099 Mainz, Germany  
<sup>e</sup> Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität, D-53115 Bonn, Germany  
<sup>f</sup> Department of Physics, University of Zagreb, HR-10002 Zagreb, Croatia  
<sup>g</sup> Jožef Stefan Institute, SI-1000 Ljubljana, Slovenia  
<sup>h</sup> Faculty of Mathematics and Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia  
<sup>i</sup> Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA  
<sup>j</sup> MIT Bates Research and Engineering Center, Middlesex, MA, 01948, USA  
<sup>k</sup> Center for Frontiers in Nuclear Science, Department of Physics and Astronomy, Stony Brook University, New York 11794, USA  
<sup>l</sup> RIKEN INFL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA  
<sup>m</sup> Department of Physics, Hampton University, Hampton, Virginia 23068, USA



S. Schlimme et al., 10.1016/j.nima.2021.165668

# Benefits of a gas jet target.



# Benefits of a gas jet target.


Typical  
liquid H<sub>2</sub>  
target



- Large energy straggling and multiple scattering
- Background from target foils/cell


# Benefits of a gas jet target.

Typical liquid H<sub>2</sub> target




- Large energy straggling and multiple scattering
- Background from target foils/cell

D. Bonaventura



S. Schlimme et al., 10.1016/j.nima.2021.165668




H<sub>2</sub> gas jet target




- Sharper elastic peak in electron-proton scattering
- Background effects drastically reduced


# The MAGIX setup.




# The MAGIX setup.






# The MAGIX setup.



# The MAGIX TPC.





# The MAGIX TPC.




- TPC based on a stack of four Gas Electron Multipliers (GEMs)
- Segmented readout at the pad plane-anode
- 3D track reconstruction via projection on pad plane-anode (2D) plus drift time (1D)
- Expected accuracy in the focal plane coordinates:  
 $\Delta x = 100\mu\text{m}$  and  $\Delta\theta = 3.5\text{mrad}$

# The MAGIX TPC.




- TPC based on a stack of four Gas Electron Multipliers (GEMs)
- Segmented readout at the pad plane-anode
- 3D track reconstruction via projection on pad plane-anode (2D) plus drift time (1D)
- Expected accuracy in the focal plane coordinates:  
 $\Delta x = 100\mu m$  and  $\Delta\theta = 3.5 mrad$
- Novel open field cage design to minimize the material budget


# The MAGIX Trigger Veto System.




- Combine triggering and PID in one modular system
- Segmented trigger layer at the top, made of plastic scintillators read out by PMTs
- A flexible veto system underneath, built from:
  - Several veto layers, made of plastic scintillators read out by SiPMs
  - Passive lead absorber layers in between




# The MAGIX Trigger Veto System.




- Combine triggering and PID in one modular system
- Segmented trigger layer at the top, made of plastic scintillators read out by PMTs
- A flexible veto system underneath, built from:
  - Several veto layers, made of plastic scintillators read out by SiPMs
  - Passive lead absorber layers in between




# The MAGIX Trigger Veto System.





- Combine triggering and PID in one modular system
- Segmented trigger layer at the top, made of plastic scintillators read out by PMTs
- A flexible veto system underneath, built from:
  - Several veto layers, made of plastic scintillators read out by SiPMs
  - Passive lead absorber layers in between






# Working principle of the trigger veto system.







# Working principle of the trigger veto system.






# Working principle of the trigger veto system.



# Working principle of the trigger veto system.



# Recoil detector array.



- Additional detector array for low-energy recoil nuclei
- Mounted inside the scattering chamber with no material between reaction vertex and detectors
- Individual detectors built from silicon strip + scintillation detectors



# MAGIX


**EX**periment

Exploration


Excellence

Exattezza


# A versatile physics program.



# A versatile physics program.




# Dark sector searches.





Nollett and Steigman, 10.1103/PhysRevD.89.083508

Griest and Kamionkowski, 10.1103/PhysRevLett.64.615




[https://www.researchgate.net/figure/Estimated-distribution-of-matter-and-energy-in-the-Universe\\_fig2\\_353762159](https://www.researchgate.net/figure/Estimated-distribution-of-matter-and-energy-in-the-Universe_fig2_353762159)

# Dark sector searches.



# Dark sector searches.




# Dark sector searches.




## Light Dark Matter (LDM)

- Dark sector that communicates with the SM through one (or more) dark mediator particles?
- Popular mediator model: Dark photon  $\gamma'$  with a mass  $m_{\chi}$  that couples e.g. via kinetic mixing




# Dark sector searches.




## Light Dark Matter (LDM)



- Dark sector that communicates with the SM through one (or more) dark mediator particles?
- Popular mediator model: Dark photon  $\gamma'$  with a mass  $m_X$  that couples e.g. via kinetic mixing



# Dark photon - visible decay.





# Dark photon - visible decay.




Coincidence measurement  
between both spectrometers



# Dark photon - visible decay.





Coincidence measurement between both spectrometers





# Dark photon - visible decay.



Coincidence measurement between both spectrometers





# Dark photon - visible decay.




- **MESA in commissioning phase:** EB mode with  $100\mu A$  beam current and beam energies of **55MeV** and **105MeV**
- **Solid target:** stack of 17 tantalum foils ( $^{181}\text{Ta}$ ) with  $1\mu m$  thickness each, spaced 2.5mm apart
- $2\sigma$  confidence level
- Background included
- 2 weeks per run

# Dark photon - visible decay.



- **MESA in commissioning phase:** EB mode with  $100\mu A$  beam current and beam energies of **55MeV** and **105MeV**
- **Solid target:** stack of 17 tantalum foils ( $^{181}\text{Ta}$ ) with  $1\mu m$  thickness each, spaced 2.5mm apart
- $2\sigma$  confidence level
- Background included
- 2 weeks per run
- **X17 range covered!**


# Dark photon - displaced vertex.



$$l_{\text{decay}} \approx 1.5 \text{ mm} \left( \frac{E_{\text{beam}}}{55 \text{ MeV}} \right) \cdot \left( \frac{10^{-4}}{\varepsilon} \right)^2 \cdot \left( \frac{17 \text{ MeV}/c^2}{m_{\gamma'}} \right)^2$$

$\uparrow$   
 $= \gamma c \tau$


# Dark photon - displaced vertex.



$$\uparrow \boxed{l_{\text{decay}}} \approx 1.5 \text{ mm} \left( \frac{E_{\text{beam}}}{55 \text{ MeV}} \right) \cdot \left( \frac{10^{-4}}{\boxed{\varepsilon}} \right)^2 \cdot \left( \frac{17 \text{ MeV}/c^2}{m_{\gamma'}} \right)^2$$




$= \gamma c \tau$

# Dark photon - displaced vertex.






$$\begin{aligned}
 \uparrow \boxed{l_{\text{decay}}} &\approx 1.5 \text{ mm} \left( \frac{E_{\text{beam}}}{55 \text{ MeV}} \right) \cdot \left( \frac{10^{-4}}{\boxed{\varepsilon}} \right)^2 \cdot \left( \frac{17 \text{ MeV}/c^2}{m_{\gamma'}} \right)^2 \\
 &= \gamma c \tau
 \end{aligned}$$

# Dark photon - displaced vertex.




Coincidence measurement between both spectrometers




# Dark photon - displaced vertex.



Coincidence measurement between both spectrometers




# Dark photon - displaced vertex.





$l_{\text{decay}} \approx 1.5 \text{ mm} \left( \frac{E_{\text{beam}}}{55 \text{ MeV}} \right) \cdot \left( \frac{10^{-4}}{\epsilon} \right)^2 \cdot \left( \frac{17 \text{ MeV}/c^2}{m_{\gamma'}} \right)^2$


$= \gamma c \tau$




Coincidence measurement between both spectrometers



# Dark photon - displaced vertex.





- **MESA in commissioning phase:** EB mode with  $100\mu A$  beam current and beam energies of **55MeV** and **105MeV**



- **Solid target:** stack of 17 tantalum foils ( $^{181}\text{Ta}$ ) with  $1\mu m$  thickness each, spaced 2.5mm apart
- $2\sigma$  confidence level
- Background not included
- 2 weeks per run


# Dark photon - displaced vertex.





Bachelor's thesis: S. Merkel

- **MESA in commissioning phase:** EB mode with  $100\mu A$  beam current and beam energies of **55MeV** and **105MeV**
- **Solid target:** stack of 17 tantalum foils ( $^{181}\text{Ta}$ ) with  $1\mu m$  thickness each, spaced 2.5mm apart
- $2\sigma$  confidence level
- Background not included
- 2 weeks per run
- **X17 range covered!**

# Outlook - invisible dark photon decay.





# Outlook - invisible dark photon decay.




Coincidence measurement between one spectrometer and the silicon strip detectors



# Outlook - invisible dark photon decay.





Coincidence measurement between one spectrometer and the silicon strip detectors





# Outlook - invisible dark photon decay.





Coincidence measurement between one spectrometer and the silicon strip detectors



# Outlook - invisible dark photon decay.



Coincidence measurement between one spectrometer and the silicon strip detectors



# Outlook - effective neutron target.

PHYSICAL REVIEW LETTERS **128**, 091802 (2022)

## X17 Discovery Potential in the $\gamma N \rightarrow e^+ e^- N$ Process at Electron Scattering Facilities

Johannes Backens<sup>1</sup> and Marc Vanderhaeghen<sup>1</sup>

<sup>1</sup>Institut für Kernphysik and PRISMA<sup>+</sup> Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

PHYSICAL REVIEW D **109**, 095010 (2024)

## Low-mass dark sector searches with deuteron photodisintegration

Cornelis J.G. Mommers<sup>1,\*</sup> and Marc Vanderhaeghen<sup>1</sup>

<sup>1</sup>Institut für Kernphysik and PRISMA<sup>+</sup> Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany


EPJ Web of Conferences **303**, 05004 (2024)  
MENU 2023

<https://doi.org/10.1051/epjconf/202430305004>

## X17 discovery potential in $\gamma d \rightarrow e^+ e^- pn$ at MAGIX@MESA

Cornelis J.G. Mommers<sup>1,\*</sup> and Marc Vanderhaeghen<sup>1</sup>

<sup>1</sup>Institut für Kernphysik and PRISMA<sup>+</sup> Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany



# Outlook - effective neutron target.

PHYSICAL REVIEW LETTERS 128, 091802 (2022)

## X17 Discovery Potential in the $\gamma N \rightarrow e^+ e^- N$ Process at Electron Scattering Facilities

Johannes Backens<sup>1</sup> and Marc Vanderhaeghen<sup>1</sup>  
*Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany*

PHYSICAL REVIEW D 109, 095010 (2024)

## Low-mass dark sector searches with deuteron photodisintegration

Cornelis J.G. Mommers<sup>1,\*</sup> and Marc Vanderhaeghen<sup>1</sup>  
*Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany*



EPJ Web of Conferences 303, 05004 (2024)  
MENU 2023

<https://doi.org/10.1051/epjconf/202430305004>

## X17 discovery potential in $\gamma d \rightarrow e^+ e^- pn$ at MAGIX@MESA

Cornelis J.G. Mommers<sup>1,\*</sup> and Marc Vanderhaeghen<sup>1</sup>

<sup>1</sup>Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany



Coincidence measurement between both spectrometers

# Outlook - effective neutron target.

PHYSICAL REVIEW LETTERS 128, 091802 (2022)

## X17 Discovery Potential in the $\gamma N \rightarrow e^+ e^- N$ Process at Electron Scattering Facilities

Johannes Backens and Marc Vanderhaeghen  
 Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

PHYSICAL REVIEW D 109, 095010 (2024)

## Low-mass dark sector searches with deuteron photodisintegration

Cornelis J.G. Mommers and Marc Vanderhaeghen  
 Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany



EPJ Web of Conferences 303, 05004 (2024)  
 MENU 2023

<https://doi.org/10.1051/epjconf/202430305004>


## X17 discovery potential in $\gamma d \rightarrow e^+ e^- pn$ at MAGIX@MESA

Cornelis J.G. Mommers<sup>1,\*</sup> and Marc Vanderhaeghen<sup>1</sup>

<sup>1</sup>Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany



Coincidence measurement between both spectrometers



# Outlook - effective neutron target.

PHYSICAL REVIEW LETTERS 128, 091802 (2022)

## X17 Discovery Potential in the $\gamma N \rightarrow e^+ e^- N$ Process at Electron Scattering Facilities

Johannes Backens and Marc Vanderhaeghen  
 Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

PHYSICAL REVIEW D 109, 095010 (2024)

## Low-mass dark sector searches with deuteron photodisintegration

Cornelis J.G. Mommers and Marc Vanderhaeghen  
 Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany



EPJ Web of Conferences 303, 05004 (2024)  
 MENU 2023

<https://doi.org/10.1051/epjconf/202430305004>

## X17 discovery potential in $\gamma d \rightarrow e^+ e^- pn$ at MAGIX@MESA


Cornelis J.G. Mommers<sup>1,\*</sup> and Marc Vanderhaeghen<sup>1</sup>

<sup>1</sup>Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, D-55099 Mainz, Germany



Coincidence measurement between both spectrometers

No exclusion limits yet



# The **MAGIX** Experiment at MESA.

## **MA**inz

MESA

Multi-purpose Apparatus

Massima Accuratezza

## **Gas Injection**

Gas Internal

Gas Interaction

Grande Innovazione

Target



## **EX**periment

Exploration

Excellence

Exattezza



# Summary



## MESA

- New high-intensity, low-energy electron accelerator
- Two modes: energy-recovery linac mode and extracted beam mode
- Exciting experimental program

# Summary




## MESA



- New high-intensity, low-energy electron accelerator
- Two modes: energy-recovery linac mode and extracted beam mode
- Exciting experimental program

## MAGIX

- High-resolution, two-spectrometer setup utilizing an internal gas jet target
- Varied and rich physics program in nuclear, particle, and few-body physics



# Summary





## MESA

- New high-intensity, low-energy electron accelerator
- Two modes: energy-recovery linac mode and extracted beam mode
- Exciting experimental program

## MAGIX

- High-resolution, two-spectrometer setup utilizing an internal gas jet target
- Varied and rich physics program in nuclear, particle, and few-body physics



## X17 searches

- Several dark photon searches (visible, displaced vertex, ...) are planned at MAGIX, all of which cover the interesting X17 range


# Impressions.



MESA cryomodule 2



Pre-accelerator MAMBO



Photogun STEAM and beam preparation system MELBA



First beam









MESA cryomodule 1



MAGIX spectrometer setup

# Impressions.



# Impressions.



**Thank you for your attention!**

## Contact

Institute for Nuclear Physics  
Johannes Gutenberg University Mainz  
Johann-Joachim-Becher-Weg 45  
55128 Mainz  
Germany

[www.kernphysik.uni-mainz.de](http://www.kernphysik.uni-mainz.de)

Sebastian Stengel  
MAGIX Collaboration  
[sestenge@uni-mainz.de](mailto:sestenge@uni-mainz.de)

[www.magix.uni-mainz.de](http://www.magix.uni-mainz.de)