



Charting the spin structure of weak-boson pairs at the LHC

Giovanni Pelliccioli


University and INFN of Milano-Bicocca

Introduction

Motivations (1)

Polarisations of weak bosons at the **LHC**:

- important probes of **SM gauge and scalar sectors**




Quantum Diaries

Motivations (2)

Polarisations of weak bosons at the LHC:

- discrimination between BSM and SM



[Szeleper 1412.8367]


Motivations (3)

Polarisations of weak bosons at the LHC:


- non trivial to extract from data

Fits of LHC Run-2 data with polarised templates


[ATLAS 1902.05759, 2211.09435, 2310.04350, 2402.16365, 2503.11317, CMS 2009.09429, 2110.11231]



WZ [ATLAS 2402.16365]




ZZ [ATLAS 2310.04350]




$W^\pm W^\pm jj$ [CMS 2009.09429]

What do we need (from theory)?




What do we need (from theory)?



Theory aspects: polarised templates

Separating polarisations in amplitudes

Natural definition for resonant diagrams (in PA/NWA) [Denner GP 2006.14867]




$$\begin{aligned}\mathcal{A}^{\text{unpol}} &= \mathcal{P}_\mu \frac{-g^{\mu\nu}}{k^2 - M_V^2 + iM_V\Gamma_V} \mathcal{D}_\nu \\ &= \mathcal{P}_\mu \frac{\sum_{\lambda'} \epsilon_{\lambda'}^\mu \epsilon_{\lambda'}^{*\nu}}{k^2 - M_V^2 + iM_V\Gamma_V} \mathcal{D}_\nu \\ &\rightarrow \mathcal{P}_\mu \frac{\epsilon_\lambda^\mu \epsilon_\lambda^{*\nu}}{k^2 - M_V^2 + iM_V\Gamma_V} \mathcal{D}_\nu = \mathcal{A}_\lambda\end{aligned}$$

$$|\mathcal{A}^{\text{unpol}}|^2 = \underbrace{\sum_\lambda |\mathcal{A}_\lambda|^2}_{\text{incoherent sum}} + \underbrace{\sum_{\lambda \neq \lambda'} \mathcal{A}_\lambda^* \mathcal{A}_{\lambda'}}_{\text{interference terms}} \rightarrow |\mathcal{A}_\lambda|^2 \propto \text{polarised cross section}$$

Polarisation states defined in **specific Lorentz frame**

Selecting resonant diagrams

Factorised amplitude **not possible for all contributions**



(1) Non-resonant diagrams regarded as **non-resonant background**.

(2) Resonant diagrams treated with

DPA: double-pole approximation [Denner et al. 0006307]


NWA: spin-correlated narrow-width approximation [Artoisenet et al. 1212.3460]

(3) **Separate polarisations**

Radiative corrections to the production

(N)NLO QCD corr. with leptonic decays [Denner GP 2006.14867, Poncelet Popescu 2102.13583]

- ▶ Virtual (V) and real (R) contributions, $V + R$ free of IR singularities




- ▶ subtraction counterterms needed, e.g. dipoles [Catani, Seymour 9605323]:

$$d\sigma_{\text{nlo}}/d\xi = \int d\phi_n (B + V + \int d\phi_{\text{rad}} D)_{d=4} \delta_\xi^{(n)} + \int d\phi_{n+1} (R \delta_\xi^{(n+1)} - D \delta_\xi^{(n)})_{d=4}$$

- ▶ DPA/NWA and separation of polarisations for all contributions

Radiative corrections to the decays

NLO EW (QCD) corr. to Z/W bosons with leptonic (hadronic) decays



Z bosons [Denner GP 2107.06579, Denner Haitz GP 2211.09040]

W bosons [Le Baglio 2203.01470, Denner Haitz GP 2311.16031, 2409.03620, Dao Le 2311.17027]

Fixed-order SM predictions

- $W^+(\ell^+\nu_\ell)W^-(\ell'^-\bar{\nu}_{\ell'})$: NNLO QCD + NLO EW [Denner GP 2006.14867, Poncelet Popescu 2102.13583, Denner Haitz GP 2311.16031, Dao Le 2311.17027, 2409.06396]
- $W^\pm(\ell^\pm\nu_\ell)Z(\ell'^+\ell'^-)$: NNLO QCD + NLO EW [Denner GP 2010.07149, Le Baglio 2203.01470, 2208.09232, GP Poncelet 2510.25898]
- $Z(\ell^+\ell^-)Z(\ell'^+\ell'^-)$: NNLO QCD + NLO EW [Denner GP 2107.06579, Carrivale GP et al. 2505.09686]
- $W^\pm(\ell^\pm\nu_\ell)+j$: NNLO QCD [Pellen et al. 2109.14336]
- $W^\pm(jj/J)Z(\ell'^+\ell'^-)$: NLO QCD [Denner Haitz GP 2211.09040]
- $W^+(\ell^+\nu_\ell)W^+(\ell'^+\nu_{\ell'})+jj$: NLO EW + QCD [Denner Haitz GP 2409.03620]
- $W^+(\ell^+\nu_\ell)Z(\ell'^+\ell'^-)+jj$: NLO EW [Denner GP et al. 2510.26462]

Matching to parton shower

QCD PS does not factorise from spin of the EW system

At least NLO QCD accuracy required

Event generators for intermediate polarised bosons (matched to PS):

1. PHANTOM: LO [Ballestrero Maina GP 1710.09339, 1907.04722, 2007.07133, Maina GP 2105.07972]
2. MG5_AMC: LO and MJM / [Buarque-Franzosi et al. 1912.01725], loop-ind. [Javurkova et al. 2401.17365] (also MJM), mass effects [Basu Ruiz 2512.10015]
3. SHERPA: approx. NLO and MJM [Hoppe et al. 2310.14803]
4. POWHEG-BOX-RES: NLO [GP Zanderighi 2311.05220, Haisch GP et al. 2507.21768]

Effort needed to incorporate EW effects


What about BSM effects?

Polarised signals with BSM effects:

- **VBS**: LO in HSESM with PHANTOM [Ballestrero Maina GP 1907.04722], in CH models with MADGRAPH5 [Buarque-Franzosi et al. 1912.01725]
- **inc. VV** : NLO QCD + PS in SMEFT (dim. 6) with POWHEG-BOX-RES [Haisch GP et al. 2507.21768]

Inclusive di-boson production

General features



- sensitive to anomalous **TGC** and **Vff** couplings [Azatov et al. 1707.08060, Baglio et al. 1708.03332, Panico et al. 1708.07823, ...]
- huge **QCD corr.** from large IR logs [Rubin et al. 1006.2144]
- interplay with **top sector** in **gg** [Grazzini et al. 1811.09593, Rossia et al. 2306.09963, ...]
- sizeable **loop EW corr.** in the tails from Sudakov logs [Denner Pozzorini 0010201]
- **VV-CM** natural polarisation frame

ZZ (1): a community effort

Coordination within **COMETA** network [Carrivale GP et al. 2505.09686]

$pp \rightarrow e^+e^-\mu^+\mu^-$ in ATLAS fiducial setup [ATLAS 2310.04350]

Involved **all MC tools available** on the market and **all experts** in the field:

MC tool	OS approx.	LO tree	LO loop-ind.	NLO QCD	NNLO QCD	NLO EW	LO \times PS	NLO \times PS	MJ merging	code public
MoCANLO	DPA	✓	✓	✓	✗	✓	✗	✗	✗	(✓)
STRIPPER	DPA	✓	✓	✓	✓	✗	✗	✗	✗	✗
MuLBos	DPA	✓	✓	✓	✗	✓	✗	✗	✗	✗
BBMC	DPA	✓	✗	✓	✗	✓	✗	✗	✗	✗
SHERPA	NWA	✓	✗	(✓)	✗	✗	✓	(✓)	✓	✓
MG5_AMC	BW	✓	✓	✗	✗	✗	✓	✗	✓	✓
POWHEG-BOX	DPA	✓	✗	✓	✗	✗	✓	✓	✗	✓

Our best predictions:

$$\text{NNLO}_{\text{QCD, gg}}^{\text{EW}} \rightarrow \sigma_{q\bar{q}}^{\text{LO}} \left(1 + \delta_{\text{QCD}}^{(\text{NLO})} + \delta_{\text{QCD}}^{(\text{NNLO})} + \delta_{\text{EW}}^{(\text{NLO})} \right) + \sigma_{\text{gg}}^{\text{LO}}$$

$$\text{NLOPS}_{\text{had}} \rightarrow \text{NLO}_{\text{QCD}} \times \text{PS}_{\text{QCD, QED}} + (\text{hadr.} + \text{MPI})$$

$$\text{nLO-MJM}_{\text{had}} \rightarrow (\text{nLO}_{\text{QCD}}^{(0j)} + \text{nLO}_{\text{QCD}}^{(1j)} + \text{LO}^{(2j)}) + \text{PS}_{\text{QCD, QED}} + (\text{hadr.} + \text{MPI})$$

Pol. fraction = cross-sect. ratio, e.g. $f_{\text{LL}} = \sigma_{\text{LL}} / \sigma_{\text{unp}}$


Comparison with pre/post-fit results by ATLAS [[ATLAS 2310.04350](#)]

ZZ (3): joint polarisation fractions


our predictions	LL [%]	LT [%]	TL [%]	TT [%]
LO _{q\bar{q}}	5.85 ^{+0.07} _{-0.08}	11.86 ^{+0.09} _{-0.11}	11.89 ^{+0.09} _{-0.11}	69.27 ^{+0.30} _{-0.26}
LO _{g\bar{g}}	5.39 ^{+0.02} _{-0.02}	2.12 ^{+0.01} _{-0.01}	2.10 ^{+0.01} _{-0.01}	90.48 ^{+0.03} _{-0.03}
NLO _{EW}	5.89 ^{+0.06} _{-0.08}	11.97 ^{+0.09} _{-0.11}	12.01 ^{+0.09} _{-0.11}	68.98 ^{+0.29} _{-0.25}
NLO _{QCD}	5.87 ^{+0.03} _{-0.05}	12.74 ^{+0.07} _{-0.06}	12.69 ^{+0.07} _{-0.06}	67.35 ^{+0.14} _{-0.16}
NNLO _{QCD}	6.07 ^{+0.05} _{-0.04}	13.11 ^{+0.09} _{-0.08}	13.04 ^{+0.08} _{-0.07}	66.20 ^{+0.19} _{-0.24}
NNLO _{QCD} ^{EW}	6.12 ^{+0.06} _{-0.04}	13.29 ^{+0.09} _{-0.08}	13.21 ^{+0.08} _{-0.07}	65.75 ^{+0.20} _{-0.25}
NNLO_{QCD, g\bar{g}}^{EW}	6.05^{+0.03}_{-0.03}	12.15^{+0.10}_{-0.15}	12.07^{+0.11}_{-0.16}	68.29^{+0.30}_{-0.21}
NLOPS _{QCD}	5.88 ^{+0.03} _{-0.04}	12.76 ^{+0.08} _{-0.06}	12.71 ^{+0.07} _{-0.06}	67.30 ^{+0.13} _{-0.15}
nLOPS _{QCD}	6.02 ^{+0.05} _{-0.08}	13.04 ^{+0.04} _{-0.09}	12.97 ^{+0.04} _{-0.09}	66.61 ^{+0.14} _{-0.47}
NLOPS_{had}	5.86^{+0.03}_{-0.04}	12.74^{+0.08}_{-0.06}	12.69^{+0.07}_{-0.06}	67.38^{+0.13}_{-0.15}
nLOPS _{had}	5.98 ^{+0.03} _{-0.07}	12.99 ^{+0.02} _{-0.09}	12.96 ^{+0.02} _{-0.09}	66.70 ^{+0.22} _{-0.46}
LO-MJM _{QCD}	5.79 ^{+0.08} _{-0.09}	12.91 ^{+0.06} _{-0.05}	12.84 ^{+0.06} _{-0.06}	66.81 ^{+0.24} _{-0.22}
LO-MJM _{had}	5.91 ^{+0.01} _{-0.10}	12.84 ^{+0.13} _{-0.23}	12.79 ^{+0.12} _{-0.23}	67.14 ^{+1.08} _{-0.98}
nLO-MJM_{had}	6.14^{+0.12}_{-0.11}	13.35^{+0.47}_{-0.32}	13.23^{+0.32}_{-0.28}	65.85^{+1.11}_{-0.96}
ATLAS measurement				
pre-fit	6.1 ± 0.4	22.9 ± 0.9	69.9 ± 3.9	
post-fit	7.1 ± 1.7	22.8 ± 1.1	69.0 ± 2.7	

ZZ (4): comparison with ATLAS

Best predictions [Carrivale GP et al. 2505.09686] vs ATLAS modelling [ATLAS 2310.04350]:



(qq) agreement, small QCD unc.





(gg) NLO QCD urgent

$pp \rightarrow e^+ \nu_e \mu^+ \mu^-$ in ATLAS fiducial setup [2211.09435]

Improved SM pred., NNLO QCD (δ_s) $+/\times$ NLO EW (δ_{ew}) [GP Poncelet 2510.25898]

$$d\sigma_{\text{NNLO}}^{(+)} = d\sigma_{\text{LO}} \left(1 + \delta_{s, u\bar{d}}^{(1)} + \delta_{ew, u\bar{d}}^{(1)} + \delta_{s, u\bar{d}}^{(2)} \right) \\ + d\sigma_{qg} \left(1 + \delta_{s, qg}^{(1)} \right) + d\sigma_{q\gamma} + d\sigma_{qq'} + d\sigma_{gg}$$

state	$\delta_{s, u\bar{d}}^{(1)}$	$\frac{d\sigma_{qg}}{d\sigma_{\text{LO}}}$	$\delta_{s, u\bar{d}}^{(2)}$	$\frac{d\sigma_{qg}}{d\sigma_{\text{LO}}} \delta_{s, qg}^{(1)}$	$\frac{d\sigma_{qq'}}{d\sigma_{\text{LO}}}$	$\frac{d\sigma_{gg}}{d\sigma_{\text{LO}}}$	$\frac{d\sigma_{q\gamma}}{d\sigma_{\text{LO}}}$	$\delta_{ew, u\bar{d}}^{(1)}$
LL	31.9%	-2.1%	7.7%	0.2%	2.7%	0.5%	0.6%	-5.1%
LT	50.5%	108.5%	13.3%	19.5%	10.6%	-0.8%	1.4%	-4.9%
TL	50.9%	110.9%	13.7%	20.1%	10.7%	-0.8%	4.1%	-5.0%
TT	29.7%	31.3%	5.7%	-0.8%	3.9%	-0.1%	1.4%	-6.5%

Differential results in $\Delta y(\ell_W, Z)$ [GP Poncelet 2510.25898]

QCD unc. for K-factors from scale var. (7pSV) vs TNP [Lim Poncelet 2412.14910]

Joint fractions $f_{ij} = d\sigma_{ij}/d\sigma_{\text{unp}}$ [GP Poncelet 2510.25898] compared to ATLAS [2211.09435]

QCD-scale unc. on joint fractions: **correlated 7pSV** vs **uncorrelated 7pSV** vs **TNPs**



state	$f_{\text{NLO}}^{(\text{QCD})}$ [%]	$f_{\text{NNLO}}^{(\text{QCD})}$ [%]	$f_{\text{NNLO}}^{(+)}$ [%]	f_{meas} [%]
off-sh.	1.88	1.97	1.99	
unpol.	100	100	100	
LL	$5.71^{+0.17(+0.34)}_{-0.18(-0.27)} (\pm 0.56)$	$5.69^{+0.04(+0.18)}_{-0.03(-0.15)} (\pm 0.25)$	$5.63^{+0.04(+0.19)}_{-0.02(-0.16)}$	7.2 ± 1.6
LT	$15.42^{+0.30(+1.38)}_{-0.29(-1.11)} (\pm 2.36)$	$16.46^{+0.27(+0.77)}_{-0.22(-0.65)} (\pm 1.13)$	$16.65^{+0.26(+0.81)}_{-0.22(-0.68)}$	11.9 ± 3.4
TL	$14.68^{+0.29(+1.32)}_{-0.29(-1.06)} (\pm 2.26)$	$15.71^{+0.22(+0.71)}_{-0.20(-0.61)} (\pm 1.08)$	$16.03^{+0.21(+0.75)}_{-0.19(-0.63)}$	15.2 ± 3.3
TT	$63.55^{+0.40(+4.38)}_{-0.45(-3.53)} (\pm 7.24)$	$61.52^{+0.45(+1.72)}_{-0.58(-1.56)} (\pm 3.08)$	$61.07^{+0.45(+1.84)}_{-0.57(-1.64)}$	66.0 ± 4.0
interf.	0.63	0.63	0.62	

First SMEFT study with polarised-boson signals [Haisch GP et al. 2507.21768].

- automation in DPA based on SM package [GP Zanderighi 2311.05220] in POWHEG-BOX-RES [Alioli et al. 1002.2581, Jezo Nason 1509.09071, Chiesa et al. 2005.12146]
- pol. selection enabled in RECOLA 2 lib. [Denner Lang Uccirati 1711.07388]
- aTGC operators in Warsaw (new UFO) and HISZ basis [Chiesa Denner Lang 1804.01477]



$$\begin{aligned}
 Q_{HB} &= H^\dagger H B_{\mu\nu} B^{\mu\nu}, & Q_{H\tilde{B}} &= H^\dagger H B_{\mu\nu} \tilde{B}^{\mu\nu}, \\
 Q_{HW} &= H^\dagger H W_{\mu\nu}^i W^{i,\mu\nu}, & Q_{H\tilde{W}} &= H^\dagger H W_{\mu\nu}^i \tilde{W}^{i,\mu\nu}, \\
 Q_{HWB} &= H^\dagger \sigma^i H W_{\mu\nu}^i B^{\mu\nu}, & Q_{H\tilde{W}B} &= H^\dagger \sigma^i H \tilde{W}_{\mu\nu}^i B^{\mu\nu}, \\
 Q_W &= \epsilon_{ijk} W_\mu^{i,\nu} W_\nu^{j,\lambda} W_\lambda^{k,\mu}, & Q_{\tilde{W}} &= \epsilon_{ijk} W_\mu^{i,\nu} W_\nu^{j,\lambda} \tilde{W}_\lambda^{k,\mu}.
 \end{aligned}$$

Validation at f.o. (NLO QCD) with updated [SMEFT@NLO](#) [El Faham GP Vryonidou 2405.19083] and [SM_ATGC](#) [Chiesa Denner Lang 1804.01477] UFOs



Pheno. for $C_W, C_{\widetilde{W}} = 1\text{TeV}^{-2}$, $C_{H\widetilde{W}B} = 0.01\text{TeV}^{-2}$: large effects on **T** modes

σ [fb], ATLAS fiducial [2211.09435]					
contribution	unpolarized	$W_L Z_L$	$W_L Z_T$	$W_T Z_L$	$W_T Z_T$
SM	34.64(2)	1.968(1)	5.357(2)	5.100(2)	21.99(2)
Q_W (lin.)	-0.975(1)	0	0.1229(3)	0.1435(3)	-1.264(1)
Q_W (quad.)	6.299(2)	0	0.0803(0)	0.1001(0)	6.110(2)
$Q_{\widetilde{W}}$ (lin.)	-0.013(1)	0	-0.0039(1)	0.0056(1)	0.0102(9)
$Q_{\widetilde{W}}$ (quad.)	6.443(2)	0	0.1312(1)	0.1661(1)	6.132(2)



Quantum tomography (1)

DPA calculation allows for sound **quantum tomography** of four-lepton final states [Del Gratta GP et al. 2509.20456, GP Re 2601.09540]



$$\begin{aligned}\frac{d\sigma}{d\Omega_1 d\Omega_2 dX} / \frac{d\sigma}{dX} &= \frac{1}{(4\pi)^2} + \frac{1}{4\pi} \sum_{l=1}^2 \sum_{m=-l}^l \alpha_{lm}^{(1)}(X) Y_{lm}(\Omega_1) + \frac{1}{4\pi} \sum_{l'=1}^2 \sum_{m'=-l'}^{l'} \alpha_{l'm'}^{(2)}(X) Y_{l'm'}(\Omega_2) \\ &+ \sum_{l,l'=1}^2 \sum_{m=-l}^l \sum_{m'=-l'}^{l'} \gamma_{lm'l'm'}(X) Y_{lm}(\Omega_1) Y_{l'm'}(\Omega_2) \\ &= \left(\frac{3}{4\pi}\right)^2 \sum_{i,i',j,j'=1}^3 \rho_{i i' j j'}(X) \Gamma_{i i'}(\Omega_1) \Gamma_{j j'}(\Omega_2)\end{aligned}$$

→ $\{\alpha_{lm}, \gamma_{lm'l'm'}\}$ to reconstruct **spin-density matrix** ρ (entanglement, purity ...)

$$9\text{Tr}[\rho^2] = 1 + \sum_{l=1}^2 \sum_{m=-l}^l (\kappa_l^{(1)} \alpha_{lm}^{(1)})^2 + \sum_{l'=1}^2 \sum_{m'=-l'}^{l'} (\kappa_{l'}^{(2)} \alpha_{l'm'}^{(2)})^2 + \sum_{l,l'=1}^2 \sum_{m=-l}^l \sum_{m'=-l'}^{l'} (\kappa_l^{(1)} \kappa_{l'}^{(2)} \gamma_{lm'l'm'})^2$$

Quantum tomography (2)


Quadratic EFT terms crucial for semi-positive-definite ρ [GP Re 2601.09540]



Conclusions & Outlook

VV_POL publicly available in POWHEG-BOX-RES git repo

[Haisch Linder GP Re Zanderighi 2507.21768]



Diboson-specific Monte Carlo generator for PS-matched calculations at next-to-leading order QCD in SM and aTGC SMEFT

allows for calculations with polarised bosons in the DPA

MoCANLO public by end of February

[Denner Lombardi Lopez Pellen GP, to appear]



General-purpose (LHC, lepton colliders, heavy-ion UPC, ..) Monte Carlo **integrator** for **fixed-order** calculations (no PS matching) at complete **next-to-leading order** (α_S , α_{EW})

allows for **calculations with polarised bosons at NLO QCD + EW in the DPA**

Where do we stand?

- fixed-order in the SM ✓ [(N)NLO QCD + NLO EW]
- MC-tool comparison & LHC recommendations ✓ [ZZ]
- matching to PS / merging ✓ [NLO QCD × PS & MJM]
- SMEFT studies ✓ [NLO QCD × PS]


What is next?

- NNLO+PS automation
- ML / polarisation tagging



Backup

General features of VBS



- Sensitive to gauge-Higgs unitarity cancellations
- Large loop EW corr. from EW Sudakov logs
[Denner Pozzorini 0010201, Biedermann et al. 1611.02951]
- Sensitive to anomalous TGC, QGC and hVV couplings
[Ethier et al. 2101.03180, Bellan et al. 2108.03199, ...]
- Overlap with Higgs-strahlung and triple-boson
[Denner et al. 2406.12301, Denner Haitz GP 2409.03620, ...]

W^+W^+ (1)

$pp \rightarrow W^+(e^+\nu_e)W^+(\mu^+\nu_\mu)jj$ [Denner Hartz GP 2409.03620] @ NLO EW + QCD



CMS setup [CMS 2009.09429], Run-3 energy, WW-CM def. of polarised states

mode	σ_{LO} [fb]	δ_{EW}	δ_{QCD}	$\sigma_{\text{NLO EW+QCD}}$ [fb]
full	1.4863(1) ^{+9.2%} _{-7.8%}	-0.140	-0.047	1.208(1) ^{+1.6%} _{-3.1%}
unp.	1.46455(9) ^{+9.2%} _{-7.8%}	-0.142	-0.050	1.1836(5) ^{+1.7%} _{-3.3%}
LL	0.14879(1) ^{+8.3%} _{-7.2%}	-0.101	-0.044	0.12715(8) ^{+1.0%} _{-2.1%}
LT,TL	0.23209(2) ^{+9.1%} _{-7.8%}	-0.131	-0.042	0.1919(1) ^{+1.4%} _{-2.8%}
TT	0.87702(7) ^{+9.4%} _{-8.0%}	-0.154	-0.054	0.6944(4) ^{+1.9%} _{-3.7%}
int.	-0.0254(1) ^{-8.9%} _{+10.6%}	-0.139	-0.007	-0.0217(7) ^{-1.6%} _{+0.7%}

NLO pol. fractions similar to LO

LL: 11%, LT+TL: 32%, TT: 59%, int: -2%, non-res: 2%

Ratio between subleading- and leading-lepton p_T 's.




$pp \rightarrow W^+(e^+\nu_e)Z(\mu^+\mu^-)jj$ [Denner Haitz Franken Lombardi GP 2510.26462] @ NLO EW

ATLAS fid. setup [ATLAS 2403.15296], Run-3 energy, WZ-CM pol. def.

mode	σ_{LO} [fb]	δ_{EW} [%]	σ_{NLOEW} [fb]	f_{LO} [%]	f_{NLOEW} [%]
off-sh.	0.2723(1)	-18.7	0.2215(2)	101.7	101.4
unp.	0.26779(1)	-18.4	0.21838(4)	100.0	100.0
LL	0.023045(2)	-14.4	0.019697(5)	8.6	9.0
LT+TL	0.098391(6)	-17.9	0.08076(2)	36.7	37.0
TT	0.141723(9)	-19.6	0.11394(3)	52.9	52.2
int.	0.00463(2)	-13.9	0.00398(4)	1.7	1.7

Even larger EW corr. (-20% for TT) than in $W^\pm W^\pm$.

New: fudge factors to regulate un-subtractions in WZV topologies.

p_T of the di-jet systemrapidity sep. between ℓ_W and Z