



Contribution ID: 131

Type: Oral presentation

## Fusion and neutron transfer reactions with weakly bound nuclei within time-dependent and coupled channel approaches

Thursday, 25 June 2015 15:40 (20 minutes)

Click here to download the template: <https://agenda.infn.it/materialDisplay.py?materialId=2&confId=5235> >Lat Word </a>

The numerical solving of the time-dependent Schrödinger equation (TDSE) provides new possibilities for theoretical study of transfer reactions and the first (capture) stage of fusion reactions [1, 2]. In this model the motion of nuclei cores is described on basis of the classical physics. The small typical grid spacing ( $\sim 0.2$  fm) of TDSE method leads to correct calculations of the spatial structure of the external (valence) neutron wave function with the formation of two center (molecular) states (Fig. 1a) [3, 4]. The traditional coupled channel approach was combined with the TDSE method [2, 5]. The value of the coupling strength was determined by time-dependent two-center level populations. The coupling matrix elements were determined by the two-center wave functions of the valence neutron. They were calculated within the two-center shell model based on the Bessel series [2]. Results of the cross section calculation for the formation of the  $^{198}\text{Au}$  (Fig. 1b) and fusion (Fig. 1c) in the  $6\text{He}+^{197}\text{Au}$  reaction [6, 7] and for the formation  $^{65}\text{Zn}$  isotopes and fusion in  $6\text{He}+^{64}\text{Zn}$  reaction [8] agree satisfactorily with the experimental data near the barrier. A few additional three-body and two-body quantum models were used for more careful study the processes of neutron transfer, breakup of the weakly bound nucleus  $6\text{He}$  and sub-barrier fusion. They are: one and two nuclear cores plus one [1] and two valence neutrons, one and two cores plus a di-neutron.

Fig. 1. a) The probability density of the valence neutrons of the  $6\text{He}$  nucleus during a frontal collision with the  $^{64}\text{Zn}$  at energy in the center of mass system MeV, a scale factor is 1 fm, and radii of the circumferences equal to radii of the nuclei. b, c) The excitation functions for the formation of the  $^{198}\text{Au}$  isotope (b) and fusion (c) in the reaction  $6\text{He}+^{197}\text{Au}$ . Experimental data (circles) is from [6, 7]. Theoretical curves were calculated within the coupled channel approach (solid lines) and the TDSE method (dashed line); VB is the Coulomb barrier.

- [1] V. I. Zagrebaev, V. V. Samarin and W. Greiner, Phys. Rev. C. 75, 035809 (2007)
- [2] V. V. Samarin, Phys. Atom. Nucl. 78, 128 (2015)
- [3] V. V. Samarin, EPJ Web Conf. 66, 03075 (2014)
- [4] V. V. Samarin, EPJ Web Conf. 86, 00040 (2015)
- [5] V. V. Samarin, EPJ Web Conf. 86, 00039 (2015)
- [6] Yu. E. Penionzhkevich et al., Eur. Phys. J. A 31, 185 (2007)
- [7] A. Kulko et al., J. Phys. G 34, 2297 (2007)
- [8] V. Scuderi et al., Phys. Rev. C. 84, 064604 (2011)

**Primary author:** Prof. SAMARIN, Viacheslav (Flerov Laboratory of Nuclear Reactions, the Joint Institute for Nuclear Research, Dubna, Russia)

**Presenter:** Prof. SAMARIN, Viacheslav (Flerov Laboratory of Nuclear Reactions, the Joint Institute for Nuclear Research, Dubna, Russia)

**Session Classification:** Fusion and Fission

**Track Classification:** Fusion and Fission