



Contribution ID: 190

Type: Poster

Octupole bands in the neutron-rich nucleus ^{143}Ba

Click here to download the template: <https://agenda.infn.it/materialDisplay.py?materialId=2&confId=5235>

Yong-Jing Chen¹, Zao-Chun Gao¹, Yong-Shou Chen¹, Ya Tu²

¹ China Institute of Atomic Energy, Beijing 102413, China

² College of Physics Science and Technology, Shenyang Normal University, ShenYang 110034, China

The reflection asymmetric shell model (RASM) was developed to describe the high spin states in octupole deformed nuclei in Ra-Th octupole deformed nuclei [1] and neutron-rich nucleus ^{145}Ba in Ba-Sm region [2]. In this work, RASM was performed to investigate the reflection asymmetry in ^{143}Ba . All the observed four rotational bands are well reproduced by the present calculation with a proper octupole deformation (~ 0.08), which is consistent with the macroscopic-microscopic calculations in the literature. The two octupole deformed neutron Nilsson single-particle orbitals just below the octupole shell gap 88, with $K=1/2$ and $K=3/2$ dominate the intrinsic structure of the observed low-lying states. Based on the analysis of the calculated RASM wave functions and the pure-configuration calculation results, the assignments for the observed bands have been given. The $s=+i$ and $s=-i$ octupole bands are all based on the $K=1/2$ orbit, and they are really “parity doublets” in octupole deformed island around $Z=56$ and $N=88$. The calculated results show that the $s=+i$ and $s=-i$ octupole bands in ^{143}Ba are different from the ^{145}Ba . In ^{145}Ba , the $s=+i$ and $s=-i$ octupole bands originate from the different K orbits.

[1] Y. S. Chen and Z. C. Gao, Phys. Rev. C 63, 014314 (2000).

[2] Yong-Jing Chen, Zao-Chun Gao, Yong-Shou Chen, and Ya Tu, Phys. Rev. C 91, 014317 (2015).

Primary author: Dr CHEN, Yongjing (China Institute of Atomic Energy)

Presenter: Dr CHEN, Yongjing (China Institute of Atomic Energy)

Track Classification: Nuclear Structure