



СКОЙ И ЭКСПЕРИМЕНТАЛЬНОЙ ФИЗИКИ ПОВИ Российской Федерации



#### Light fragments from (C + Be) interactions at 0.6 GeV/nucleon

B.M.Abramov, P.N.Alekseev, Yu.A.Borodin, S.A.Bulychjov, I.A.Dukhovskoy, A.I.Khanov, A.P.Krutenkova, V.V.Kulikov, M.A.Martemianov, M.A.Matsyuk, E.N.Turdakina

Institute for Theoretical and Experimental Physics, Moscow, Russia (National Research Centre «Kurchatov Institute»)

S.G. Mashnik

Los Alamos National Laboratory, New Mexico, USA

P. I. Zarubin

Joint Institute for Nuclear Research, Dubna, Russia

XIIth International Conference on Nucleus-Nucleus Collisions (21 – 26 June 2015, Catania, Italy)



➢ FRAGM detector is optimized to measure yields of nuclear fragments produced at ion−ion interactions and operated at TWAC accelerated complex at ITEP (Moscow)

≻ Reaction that is being studied in our experiment :  ${}^{12}C + Be \rightarrow f + X$ , where f – proton or nuclear fragment registered by detector at small angle (~ 3.5<sup>0</sup>)

Experimental setup permits us to detect a large set of the fragments (p, d, t, <sup>3</sup>He, <sup>4</sup>He, <sup>6</sup>He, <sup>8</sup>He, etc.) for projectile kinetic energies  $T_0 = 0.2 - 3.2$  GeV/nucleon

Precise measurement of high energy fragment spectra allows :

- Study cumulative (high momentum) effect for protons produced in the kinematic region forbidden for interaction with free nucleon. Cumulative particles were discovered in 70's (JINR, Dubna), but the nature of effect is still under discussion
- ✓ There is a lack of data on fragment emission at intermediate energies in ion−ion collisions
- Test of the different models of ion-ion interactions covering large kinematic region (fragmentation region and cumulative part)

 $\succ$  This study is important as input to transportation codes for radiotherapy with ions and for radioactive ion beam design for TWAC future upgrade



## **ITEP accelerator complex TWAC**



TWAC= TeraWatt Accumulator Complex

TWAC current parameters

 Proton acceleration : 50 - 10000 MeV
Ion acceleration : up to 4 GeV/nucleon
Ion accumulation : up to 700 MeV/nucleon
Accelerating ions : up to <sup>56</sup>Fe
Typical intensity : 10<sup>11</sup> nucleons / s



### **Experiment FRAGM**





# **Fragment production in FRAGM**

Example: C – Be collisions at 0.3 GeV/nucleon



**TDC** is a function of A, so it gives a mark for clear fragment identification



Beamline has several construction features (beam pipe break ~ 3 m, stubs etc.); all counters are positioned through beam. Detection efficiency depends on beam momentum

MC for FRAGM is performed with GEANT4-based code developed by our group
Protons and light ions are used (<sup>2</sup>H, <sup>3</sup>H, <sup>3</sup>He, <sup>4</sup>He and etc.) at 0.6 < P/Z < 6 GeV/c</li>
Values of the magnet currents are recomputed for different momenta



Program transports particles in the magneto – optical channel taking into account multiple scattering effects, ionization losses and absorption in the detector materials.

 $\blacktriangleright$  Efficiency correction is essential for P/Z < 2 GeV/c



➢ Our experimental data are useful to test different ion − ion interaction models. We selected four more reliable (BC, QMD, INCL and LAQGSM) to test :

✓ Binary Cascade (BC, GEANT4 toolkit) :

- > Useable when either projectile or target is  ${}^{12}C$  or lighter
- It uses a 3-dimensional model of the nucleus
- Model is based exclusively on binary scattering between reaction participants and nucleons within this nuclear model
- Novel approach of the intra-nuclear cascade is implemented. Cascade stops if the mean kinetic energy of the participants are below 15 MeV

(G. Folger et al., EPJA 21 (2004) 407)

✓ Quantum Molecular Dynamics (QMD, GEANT4 toolkit) :

- Available for light and heavy ions
- All nucleons (target and projectile) are considered as participants and are propagated by means of a phenomenological nucleon-nucleon potential
- Time evolution of the system is stopped at 100 fm/c where it is assumed that equilibrium has been achieved
- (T. Koi et al., AIP Conf. Proc. 896 (2007) 21)

✓ Liege Intranuclear Cascade (INCL++):

- Model is implemented in the GEANT4 toolkit
- ➢ It has recently shown promising result comparable with the BC and QMD
- > INCL is not able to use projectile heavier than A = 18
- Clustering procedure is applied to nucleus and based on a coalescence model
- Cascade procedure is time-dependent and stopping time :  $t = 70 \times (A_{TARGET} / 208)^{0.16}$  fm/c
- (J. Dudouet et al., PR C89 (2014)054616)

✓ Los Alamos version of the Quark Gluon String Model (LAQGSM03.03) (by courtesy of

S. Mashnik) :

- First stage is the internuclear time-dependent cascade developed initially at JINR
- Second stage: Fermi breakup disintegration of light excited residual nuclei
- Model also accounts for coalescence of light fragments (complex particles) from energetic nucleons emitted during first stage
- It was tested in a wide energy region till 1 TeV/nucleon and large number of ions (LA-UR-11-01887)

 $\checkmark$  It was generated 10M ion – ion interactions to have a reasonable comparison with our data

## Momentum spectra :BC and INCL models

Plots present data (points) in comparison with BC and INCL models (histrograms)

Data are normalized to
BC for protons at frag mentation maximum

In the region of fragmentation peaks, all models give reasonable description of the data
In the cumulative region INCL model is in a good agreement with our data, especially for light fragments



«NN 2015», S. Mashnik





✓ Within the QMD model, fragmentation peaks are too narrow (apart from protons)
✓ High momentum region is strongly underestimated
✓ Shapes of fragmentation peaks for protons and deuterons are reasonably described by LAQGSM
✓ For <sup>2</sup>H, <sup>3</sup>He, <sup>4</sup>He the

fragmentation peaks are narrower in the model than in the data

✓ High momentum region is in a reasonable agreement with out data



## **Kinetic energy spectra : data vs models**





➢ Kinetic energy spectra (T, GeV) – fragment energy in the projectile (<sup>12</sup>C) rest frame

Plots present Data in comparison with different models :
BC, QMD, INCL and LAQGSM for different fragments



#### $Ed^{3}\sigma/d^{3}p \sim A_{S}e^{(-T/T_{S})} + A_{C}e^{(-T/T_{C})}$

- > Two slope parameters Ts and Tc (in MeV) are obtained from fit
- > Double exponential function well describes the kinetic energy spectra till <sup>4</sup>He
- ➢ Odeh: T. Odeh et al., PRL 84 (2000) 4557, <sup>197</sup>Au − <sup>197</sup>Au collisions at 1 GeV/n

Fragment	Data		INCL model		BC model		Odeh
	Ts (MeV)	Tc (MeV)	Ts (MeV)	Tc (MeV)	Ts (MeV)	Tc (MeV)	Tc (MeV)
р	$5.4 \pm 0.3$	$26.5\pm0.5$	$4.3\pm0.7$	$20.5\pm0.3$	$2.0 \pm 0.1$	$11.6\pm0.7$	$25.5 \pm 1.0$
<sup>2</sup> H	$8.1 \pm 0.2$	$17.0 \pm 0.4$	$8.1 \pm 0.1$	$20.9\pm0.3$	$6.0 \pm 0.2$	$10.6 \pm 0.4$	$16.0 \pm 1.0$
<sup>3</sup> H	$8.2 \pm 0.2$	$18.6\pm0.8$	$8.9 \pm 0.2$	$21.0 \pm 1.0$	$7.4 \pm 0.1$	$15.8 \pm 5.3$	$15.0 \pm 1.0$
<sup>3</sup> He	$9.4 \pm 0.2$	$28.0\pm6.6$	$9.3 \pm 0.2$	$22.0 \pm 1.0$	$7.7 \pm 0.1$	$21.2 \pm 6.7$	$19.0 \pm 1.0$
<sup>4</sup> He	$6.9 \pm 0.1$	$14.2 \pm 1.0$	$6.7 \pm 0.1$	$17.0 \pm 1.0$	$3.5 \pm 0.1$	$6.1\pm0.2$	$14.0 \pm 1.0$

Slope parameters Ts in Odeh change in the range:  $5 \div 7$  MeV

> INCL model is in a reliable agreement with our data and «Odeh» data



## **Fit parameters comparison**

> Plots show a comparison of the different parameters ( $\sigma^2/M$ , Ts) for two models (INCL and BC)

Analyzed fragments : p, <sup>2</sup>H, <sup>3</sup>H,
<sup>3</sup>He, <sup>4</sup>He, <sup>6</sup>He, <sup>6</sup>Li, <sup>7</sup>Li, <sup>8</sup>Li, <sup>9</sup>Li, <sup>7</sup>Be,
<sup>9</sup>Be, <sup>10</sup>Be, <sup>8</sup>B, <sup>10</sup>B, <sup>11</sup>B)

> σ<sup>2</sup>/M is RMS<sup>2</sup> of the fragmentation peak divided by fragment mass, Ts − slope parameter

Our experimental data are shown by the filled points, models by the open points

Fragments with atomic number Z > 2 are well described only by one exponential component with slope parameter Ts



«NN 2015», S. Mashnik



✓ Fragment yields from reaction  ${}^{12}C + Be \rightarrow f + X$  were measured at projectile kinetic energy T0 = 0.6 GeV/nucleon at 3.5<sup>0</sup>

✓ Several models (BC, QMD, INCL and LAQGSM) of the ion – ion collisions were compared to data

✓ In the region of the fragmentation peak all models give reasonable description of the data. INCL model is closer to our data in comparison with other models
✓ Kinetic energy spectra in the projectile rest frame can be parameterized as A<sub>s</sub>e<sup>(-T/Ts)</sup> + A<sub>c</sub>e<sup>(-T/Tc)</sup>, where Ts and Tc are slope parameters («temperatures»)
✓ Ts values for protons and other fragments is in a good agreement with predictions of the INCL model

 $\checkmark$  Tc can be used for a description of the kinetic energy spectra up to <sup>4</sup>He. Our data are in a reasonable agreement with INCL model and with other experimental data



