



Contribution ID: 154

Type: Oral presentation

Shear Viscosity to electric conductivity ratio for the Quark-Gluon Plasma

Tuesday, June 23, 2015 5:50 PM (20 minutes)

Click here to download the template: <https://agenda.infn.it/materialDisplay.py?materialId=2&confId=5235>

Transport coefficients of strongly interacting matter are currently subject of intense studies due to their relevance for the characterization of the quark-gluon plasma (QGP) produced in ultra-relativistic heavy-ion collisions (uRHIC).

We discuss the connection between the shear viscosity to entropy density ratio, η/s , and the electric conductivity, σ_{el} : we find that a minimal η/s is consistent with a low value of electric conductivity as measured in recent lattice QCD calculations.

More generally we show that the ratio of η/s over σ_{el}/T supplies a measure of the quark to gluon scattering rates whose knowledge would allow to significantly advance in the understanding of the QGP phase.

We also predict that $(\eta/s)/(\sigma_{el}/T)$, independently on the running coupling $\alpha_s(T)$, should increase up to about ~ 20 for $T \rightarrow T_c$, while it goes down to a nearly flat behaviour around $\simeq 4$ for $T \geq 4T_c$.

Primary author: PUGLISI, Armando (INFN-LNS, Catania, Italy)

Co-authors: PLUMARI, Salvatore (LNS); Dr GRECO, Vincenzo (LNS)

Presenter: PUGLISI, Armando (INFN-LNS, Catania, Italy)

Session Classification: Relativistic Heavy-Ion Collisions

Track Classification: Relativistic Heavy-Ion Collisions