

Coulomb excitation of re-accelerated $^{208,210}\text{Rn}$ and ^{206}Po at MINIBALL

martedì 11 giugno 2013 12:15 (20 minuti)

In regions near magic nuclei, seniority can be regarded as a good quantum number. In the trans-Pb nuclei near the $Z=82$ and $N=126$ shell closures, relative high- j single-particle proton orbitals dominate the structure and thus levels up to $I=2j-1$ could, in principle, be understood within the seniority scheme. In $N=122$, $N=124$ and especially in the closed shell $N=126$ isotones with $Z \geq 82$, behaviour of the $B(E2)$ values resembling the seniority scheme predictions has been observed. These nuclei lie in, or at the boundary of the region where seniority scheme could persist. However, contributions from collective excitations can not be ignored when moving away from the $N=126$ closed shell. To date, surprisingly little is known of the transition probabilities between the low-spin states in this region.

In the present study, $B(E2;0^+ \rightarrow 2^+)$ values have been measured in $^{208,210}\text{Rn}$ and ^{206}Po nuclei through Coulomb excitation of re-accelerated radioactive beams in inverse kinematics. The radioactive beams were produced at ISOLDE by bombarding a UCx primary target with 1.4 GeV protons. The mass separated radioactive beams were re-accelerated with the REX-ISOLDE linear accelerator to 2.8 MeV/u and delivered to the target position of the MINIBALL γ -ray spectrometer, which recorded γ -rays following Coulomb excitation. The resulting $B(E2;0^+ \rightarrow 2^+)$ values in $^{208,210}\text{Rn}$ and ^{206}Po are presented and discussed in terms of systematics and relevant nuclear models. The present study provides new insight into the interplay between collective excitations and single-particle regime near $N=126$.

Autore principale: Dr. GRAHN, Tuomas (University of Jyväskylä)

Relatore: Dr. GRAHN, Tuomas (University of Jyväskylä)

Classifica Sessioni: Session 6