



Contribution ID: 33

Type: **not specified**

Searches for rare and forbidden kaon decays at the NA62 experiment at CERN

Wednesday, 11 September 2013 12:50 (20 minutes)

The decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ is highly suppressed in the Standard Model (SM), while its rate can be predicted with minimal theoretical uncertainty. The branching ratio for this decay is thus a sensitive probe of the flavor sector of the SM.; however, the smallness of this BR (8×10^{-11}) and challenging experimental signature make it very difficult to measure. The primary goal of the NA62 experiment at the CERN SPS is to measure BR($K^+ \rightarrow \pi^+ \nu \bar{\nu}$) with $\sim 10\%$ precision. This will require the observation of 10^{13} K^+ decays in the experiment's fiducial volume, as well as the use of high-performance systems for precision tracking, particle identification, and photon vetoing. These aspects of the experiment will also allow NA62 to carry out a rich program of searches for lepton flavor and/or number violating K^+ decays. Such searches can probe new physics scenarios involving, for example, heavy Majorana neutrinos or R-parity violating SUSY. Part of the experimental apparatus was commissioned during a technical run in 2012; installation continues and data taking is expected to begin in late 2014. The physics prospects and the status of the NA62 experiment will be reviewed.

Primary author: MASSAROTTI, Paolo (NA)

Presenter: MASSAROTTI, Paolo (NA)

Session Classification: Flavour Physics (II)