

doubly heavy exotics

Marek Karliner

JHEP 7,153(2013) – arXiv:1304.0345, with S. Nussinov

JHEP 8,96(2013) – arXiv:1305.6457, with Y. Frishman



Tel Aviv University



From ϕ to ψ , Rome, Sep. 2013

Possibility of Exotic States in the Upsilon system

Marek Karliner^{a*}
and
Harry J. Lipkin^{a,b†}

Abstract

Recent data from Belle show unusually large partial widths $\Upsilon(5S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ and $\Upsilon(5S) \rightarrow \Upsilon(2S) \pi^+ \pi^-$. The $Z(4430)$ narrow resonance also reported by Belle in $\psi' \pi^+$ spectrum has the properties expected of a $\bar{c}cud\bar{d}$ charged isovector tetraquark $T_{\bar{c}c}^\pm$. The analogous state $T_{\bar{b}b}^\pm$ in the bottom sector might mediate anomalously large cascade decays in the Upsilon system, $\Upsilon(mS) \rightarrow T_{\bar{b}b}^\pm \pi^\mp \rightarrow \Upsilon(nS) \pi^+ \pi^-$, with a tetraquark-pion intermediate state. We suggest looking for the $\bar{b}bud\bar{d}$ tetraquark in these decays as peaks in the invariant mass of $\Upsilon(1S) \pi$ or $\Upsilon(2S) \pi$ systems. The $\bar{b}bu\bar{s}$ tetraquark can appear in the observed decays $\Upsilon(5S) \rightarrow \Upsilon(1S) K^+ K^-$ as a peak in the invariant mass of $\Upsilon(1S) K$ system. We review the model showing that these tetraquarks are below the two heavy meson threshold, but respectively above the $\Upsilon \pi \pi$ and $\Upsilon K \bar{K}$ thresholds.

Observation of two charged bottomonium-like resonances


The Belle Collaboration

(Dated: May 24, 2011)


Abstract

We report the observation of two narrow structures at $10610 \text{ MeV}/c^2$ and $10650 \text{ MeV}/c^2$ in the $\pi^\pm \Upsilon(nS)$ ($n = 1, 2, 3$) and $\pi^\pm h_b(mP)$ ($m = 1, 2$) mass spectra that are produced in association with a single charged pion in $\Upsilon(5S)$ decays. The measured masses and widths of the two structures averaged over the five final states are $M_1 = 10608.4 \pm 2.0 \text{ MeV}/c^2$, $\Gamma_1 = 15.6 \pm 2.5 \text{ MeV}$ and $M_2 = 10653.2 \pm 1.5 \text{ MeV}/c^2$, $\Gamma_2 = 14.4 \pm 3.2 \text{ MeV}$. Analysis favors quantum numbers of $I^G(J^P)=1^+(1^+)$ for both states. The results are obtained with a 121.4 fb^{-1} data sample collected with the Belle detector near the $\Upsilon(5S)$ resonance at the KEKB asymmetric-energy e^+e^- collider.


arXiv:1105.4583v1 [hep-ex] 23 May 2011




$\Upsilon(3S)\pi^+$



$\Upsilon(2S)\pi^+$



$\Upsilon(1S)\pi^+$



Comparison of $Z_b(10610)$ and $Z_b(10650)$ parameters obtained from different decay channels. The vertical dotted lines indicate $B^*\bar{B}$ and $B^*\bar{B}^*$ thresholds.

$$J^P = 1^+ \quad \text{for both } Z_b(10610) \text{ and } Z_b(10650)$$

The Z_b resonances decay into

$\Upsilon(nS)$ and a charged pion

\implies must contain both $\bar{b}b$ and $\bar{d}u$



\rightarrow manifestly exotic

Neutral member of the $I=1$ multiplet
very recently also observed
by Belle in Dalitz plot analysis

■ $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^0\pi^0$ decay

In this fit mass and width are fixed from
the charged Z_b result.

— fit result with Z_b
— fit result without Z_b



Simultaneous fit gives 6.3σ for $Z_b(10610)^0$

After the discovery of Z_b -s by Belle,
natural to expect analogous states
in the charm system

one caveat:

a priori unknown whether charmed quarks
are heavy enough to allow for binding

encouraging indications from toy model
of QCD in $D=1+1$


[JHEP 8,96(2013) - arXiv:1305.6457]

in March 2013 BES in Beijing,
followed by Belle in KEK provided the answer:



Observation of a Charged Charmoniumlike Structure in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at $\sqrt{s} = 4.26$ GeV


We study the process $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at a center-of-mass energy of 4.260 GeV using a 525 pb^{-1} data sample collected with the BESIII detector operating at the Beijing Electron Positron Collider. The Born cross section is measured to be $(62.9 \pm 1.9 \pm 3.7) \text{ pb}$, consistent with the production of the $Y(4260)$. We observe a structure at around $3.9 \text{ GeV}/c^2$ in the $\pi^\pm J/\psi$ mass spectrum, which we refer to as the $Z_c(3900)$. If interpreted as a new particle, it is unusual in that it carries an electric charge and couples to charmonium. A fit to the $\pi^\pm J/\psi$ invariant mass spectrum, neglecting interference, results in a mass of $(3899.0 \pm 3.6 \pm 4.9) \text{ MeV}/c^2$ and a width of $(46 \pm 10 \pm 20) \text{ MeV}$. Its production ratio is measured to be $R = (\sigma(e^+e^- \rightarrow \pi^\pm Z_c(3900)^\mp \rightarrow \pi^+\pi^-J/\psi)/\sigma(e^+e^- \rightarrow \pi^+\pi^-J/\psi)) = (21.5 \pm 3.3 \pm 7.5)\%$. In all measurements the first errors are statistical and the second are systematic.





Study of $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and Observation of a Charged Charmoniumlike State at Belle

The cross section for $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ between 3.8 and 5.5 GeV is measured with a 967 fb^{-1} data sample collected by the Belle detector at or near the $Y(nS)$ ($n = 1, 2, \dots, 5$) resonances. The $Y(4260)$ state is observed, and its resonance parameters are determined. In addition, an excess of $\pi^+\pi^- J/\psi$ production around 4 GeV is observed. This feature can be described by a Breit-Wigner parametrization with properties that are consistent with the $Y(4008)$ state that was previously reported by Belle. In a study of $Y(4260) \rightarrow \pi^+\pi^- J/\psi$ decays, a structure is observed in the $M(\pi^\pm J/\psi)$ mass spectrum with 5.2σ significance, with mass $M = (3894.5 \pm 6.6 \pm 4.5) \text{ MeV}/c^2$ and width $\Gamma = (63 \pm 24 \pm 26) \text{ MeV}/c^2$, where the errors are statistical and systematic, respectively. This structure can be interpreted as a new charged charmoniumlike state.




$$M_{Z_c} = 3899.0 \pm 3.6 \pm 4.9 \text{ MeV}$$

$$\Gamma_{Z_c} = 46 \pm 10 \pm 20 \text{ MeV}$$


$Z_c^+(3900)$ decays to $J/\psi \pi^+$

should also be seen in $\bar{D}D^*$


a) pion



b) proton



c) $Z_c(3900)$



tetraquark or a “molecule” ?

The molecule idea has a long history:
Voloshin & Okun 1976,
de Rujula, Georgi & Glashow 1977
Tornqvist, Z. Phys. C61,525 (1993)

Z_b -s sit 3 MeV above the $\bar{B}B^*$ and \bar{B}^*B^* thresholds

$X(3872)$ sits at $\bar{D}D^*$ threshold

strong hints in favor of the molecular interpretation

what about $Z_c(3900)$?

Heavy-light $Q\bar{q}$ mesons have $I=1/2$

→ they couple to pions

→ deuteron-like meson-meson bound states,
“deusons”

via pion exchange - no $\bar{D}D$, only $\bar{D}D^*$

$D\bar{D}^*$ ($I=0$) at threshold \leftrightarrow **X(3872) !**

S-wave $\rightarrow J^P = 1^+$

$I=1$ attraction x3 weaker than $I=0$

→ $I=1$ expected well above threshold

What about $\bar{B}B^*$ analogue ?...

B B* vs D D*:

-- same attractive potential

-- much heavier, so smaller kinetic energy

→ expect $B\bar{B}^*$ and $B^*\bar{B}^*$ $l=1$ states near threshold

→ $Z_b(10610)$ and $Z_b(10650)$ seen by Belle !!!

$l=0$ binding much stronger

→ $l=0$ states expected well below threshold

EXP signature:

$$X_b(I = 0) \longrightarrow \Upsilon(nS)\omega, \quad \chi_b \pi^+ \pi^-$$

perhaps also

$$X'_b(I = 0) \longrightarrow \Upsilon(nS)\bar{B}^* B \gamma \text{ via EM } B^* \rightarrow B \gamma$$

→ **LHCb!**

in the $M_Q \longrightarrow \infty$ limit attractive potential between the two heavy mesons becomes universal, as kinetic energy vanishes:

$$\text{Kinetic } E \sim \frac{p^2}{M_Q} \longrightarrow 0 \quad \text{as } M_Q \rightarrow \infty$$

→ treat kinetic E as perturbation:

$$H = a \cdot p^2 + V(r) \quad \text{where } a \equiv 1/2\mu_{\text{red}}$$

convert the parameter $a \sim 1/M_Q$ into a dimensionless parameter \tilde{a}

“natural” unit of ~ 0.8 Fermi $\sim 4.0 \text{ GeV}^{-1}$

With $m_D \sim 2 \text{ GeV}$ and $m_B \sim 5.3 \text{ GeV}$

$$\tilde{a}(D) = 1/8 \qquad \tilde{a}(B) = 1/21$$

→ small: can use 1-st order P.T.

for $l=1$ potential have 2 data points:

$Z_c(3900)$ at $\tilde{a}(D)$ approximately 27 MeV above $\bar{D}D^*$ threshold

$Z_b(10610)$ at $\tilde{a}(B)$ approximately 3 MeV above $\bar{B}B^*$ threshold

Linear extrapolation to $\tilde{a} = 0$ yields

$$E_b^{I=1}(\tilde{a}=0) \approx -11.7 \text{ MeV}$$

In view of the convexity, the actual binding energy likely to slightly exceed this linear extrapolation

→ use this result for the isovector channel to estimate the $\bar{B}B^*$ binding in the isoscalar channel

Assuming that the isoscalar binding energy in the $m_Q \rightarrow \infty$ limit is 3 times larger than for the isovector,

$$E_b^{I=0}(\tilde{a}=0) \approx 3 \cdot (-11.7) = -35 \text{ MeV}$$

$X(3872)$ at $\bar{D}D^*$ threshold → $E_b^{I=0}(\tilde{a}(D)) \approx 0$

Linear extrapolation to $\tilde{a}(B)$ yields $\bar{B}B^*$ binding energy in the isoscalar channel $\approx -20 \text{ MeV}$

Heavy Quark Nuclear Physics!

the newly discovered $Z_c(3900)$ isovector resonance confirms and refines the estimates for the mass of the putative $\bar{B}B^*$ isoscalar bound state.

immediately leads to several predictions:

- two $l=0$ narrow resonances in bottomonium system,
~23 MeV below $Z_b(10610)$ and $Z_b(10650)$, i.e.
~20 MeV below $\bar{B}B^*$ and \bar{B}^*B^* thresholds
- $l=0$ resonance near \bar{D}^*D^* threshold
- $l=1$ resonance slightly above \bar{D}^*D^* threshold

the newly discovered $Z_c(3900)$ isovector resonance confirms and refines the estimates for the mass of the putative $\bar{B}B^*$ isoscalar bound state.

immediately leads to several predictions:

- two $l=0$ narrow resonances in bottomonium system,
~23 MeV below $Z_b(10610)$ and $Z_b(10650)$, i.e.
~20 MeV below $\bar{B}B^*$ and \bar{B}^*B^* thresholds

- $l=0$ resonance near \bar{D}^*D^* threshold

- $l=1$ resonance slightly above \bar{D}^*D^* threshold

reported by BES: Aug 13, arXiv:1308.2760

$Z_c^+(4025)$: $M=4026.3 \pm 2.6 + 3.7$, $\Gamma = 24.8 \pm 5.6 \pm 7.7$ MeV

& Sep 10 arXiv:1309.1896

$Z_c^+(4020)$: $M=4022.9 \pm 0.8 + 2.7$, $\Gamma = 7.9 \pm 2.7 \pm 2.6$ MeV (mass a bit low)

Likely observable at LHC and Tevatron:

Guo, Meißner & Wang, arXiv:1308.0193

\sim nb x-section for $Z_b(10610)$ and $Z_b(10650)$

x-section for $Z_c(3900)$ and $Z_c(4020)$

larger by a factor of 20-30

large enough to be observed

x-section for neutral exotic states ?

Null result from CMS:

CERN-PH-EP/2013-157
2013/09/03

CMS-BPH-11-016

Search for a new bottomonium state decaying to $Y(1S)\pi^+\pi^-$ in pp collisions at $\sqrt{s} = 8$ TeV

The CMS Collaboration*

Abstract

The results of a search for the bottomonium counterpart, denoted as X_b , of the exotic charmonium state $X(3872)$ is presented. The analysis is based on a sample of pp collisions at $\sqrt{s} = 8$ TeV collected by the CMS experiment at the LHC, corresponding to an integrated luminosity of 20.7 fb^{-1} . The search looks for the exclusive decay channel $X_b \rightarrow Y(1S)\pi^+\pi^-$ followed by $Y(1S) \rightarrow \mu^+\mu^-$. No evidence for an X_b signal is observed. Upper limits are set at the 95% confidence level on the ratio of the inclusive production cross sections times the branching fractions to $Y(1S)\pi^+\pi^-$ of the X_b and the $Y(2S)$. The upper limits on the ratio are in the range 0.9–5.4% for X_b masses between 10 and 11 GeV. These are the first upper limits on the production of a possible X_b at a hadron collider.

The null result from CMS in search for

$$X_b \rightarrow Y(1S)\pi^+\pi^-$$

is excellent news for the molecular picture,

since isoscalar X_b with $J^{PC} = 1^{++}$

cannot decay into $\Upsilon(1S)\pi^+\pi^-$

It can decay into $\Upsilon(1S)\omega$ or $\chi_b\pi^+\pi^-$

$\Sigma_b^+ \Sigma_b^-$ dibaryon ?

Σ_b heavier, with $I = 1 \rightarrow$ stronger binding via π

\rightarrow deuteron-like $J=1, I=0$ bound state: “beautron”

electric charges contribute extra ~ 3 MeV to binding energy

exp. signature:

$$(\Sigma_b^+ \Sigma_b^-) \rightarrow \Lambda_b \Lambda_b \pi^+ \pi^-$$

$$\Gamma(\Sigma_b^-) = 4.9 \pm 3 \text{ MeV}, \quad \Gamma(\Sigma_b^+) = 9.7 \pm 3 \text{ MeV}$$

so might be visible

should be seen in lattice QCD

$$(\Sigma_c^0 \Sigma_c^+) \rightarrow \Lambda_c \Lambda_c \pi^- \pi^0 \text{ as well?}$$

doubly heavy baryons QQq (bbq, ccq, bcq)

- not exotic, must exist
- excellent challenge for EXP (LHCb!)

$(bbq) \rightarrow (\bar{c}cs) (\bar{c}cs)q \rightarrow J/\psi J\psi \Xi$
unique signature, w/o background

- QQq and $QQ\bar{q}\bar{q}$ have the same color structure

→ once QQq mass is known, can immediately predict $QQ\bar{q}\bar{q}$ mass :

$$m(cc\bar{u}\bar{d}) = m(\Xi_{ccu}) + m(\Lambda_c) - m(D^0) - \frac{1}{4}[m(D^*) - m(D)]$$

Summary

- a simple and consistent picture emerges from Belle and BES data:
- the new exotic resonances are loosely bound states of $\bar{D}D^*$, \bar{D}^*D^* , $\bar{B}B^*$, and \bar{B}^*B^*
- prediction: \bar{D}^*D^* resonances in $l=0$ and $l=1$ channels seen!
- predictions: new $\bar{B}B^*$ and \bar{B}^*B^* states below threshold
- heavy “deuteron”: $\Sigma_b^+ \Sigma_b^-$ (and $\Sigma_c^0 \Sigma_c^+$?)
- challenge for EXP: doubly heavy baryons QQq (LHCb?)
- $QQq \rightarrow$ accurate prediction for $QQ\bar{q}\bar{q}$ tetraquark
- challenge for TH: derive from QCD