

## Dielectric laser acceleration of 28 keV electrons with the inverse Smith-Purcell effect

*Monday, June 3, 2013 3:45 PM (25 minutes)*

Dielectric laser acceleration exploiting the large optical field strength of short laser pulses and the proximity of a dielectric structure can support high acceleration gradients and may therefore lead to much smaller accelerators, with future potential application in table-top free electron lasers. We report a proof-of-concept experiment demonstrating dielectric laser acceleration of non-relativistic 28keV electrons derived from a conventional scanning electron microscope column at a single fused-silica grating. The electrons pass the grating as closely as 50nm and interact with the third spatial harmonic, which is excited by 110fs long 800nm laser pulses with a peak electric field of 2.85GV/m. The observed maximum acceleration gradient of 25MeV/m is already comparable to state-of-the-art radio-frequency structures. This work represents the first demonstration of scalable laser acceleration and of the inverse Smith-Purcell effect in the optical regime. For relativistic electrons and otherwise identical conditions up to 2 orders of magnitude larger acceleration gradients are expected.

**Author:** Prof. HOMMELHOFF, Peter (University of Erlangen and Max Planck Institute of Quantum Optics)

**Co-author:** Mr BREUER, John (Max Planck Institute for Quantum Optics)

**Presenter:** Prof. HOMMELHOFF, Peter (University of Erlangen and Max Planck Institute of Quantum Optics)

**Session Classification:** WG3 - Electron beams from electromagnetic structures, including dielectric and laser-driven

**Track Classification:** WG3 RF - Electron beams from electromagnetic structures, including dielectric and laser-driven