

## First EXL experiment with radioactive beam: Proton scattering on $^{56}\text{Ni}$

M. von Schmid<sup>1</sup>, S. Bagachi<sup>2</sup>, S. Bönig<sup>1</sup>, M. Castlós<sup>3</sup>, I. Dillmann<sup>4</sup>, C. Dimopoulou<sup>4</sup>, P. Egelhof<sup>4</sup>, V. Eremin<sup>5</sup>, H. Geissel<sup>4</sup>, R. Gernhäuser<sup>6</sup>, M.N. Harakeh<sup>2</sup>, A.-L. Hartig<sup>1</sup>, S. Ilieva<sup>1</sup>, N. Kalantar-Nayestanaki<sup>2</sup>, Y. Ke<sup>4,7</sup>, O. Kiselev<sup>4</sup>, H. Kollmus<sup>4</sup>, C. Kozhuharov<sup>4</sup>, A. Krasznahorkay<sup>3</sup>, T. Kröll<sup>1</sup>, M. Kuilman<sup>2</sup>, S. Litvinov<sup>4</sup>, Y. Litvinov<sup>4</sup>, M. Mahjour-Shafiei<sup>2</sup>, M. Mutterer<sup>4</sup>, D. Nagae<sup>8</sup>, M.A. Najafi<sup>2</sup>, C. Nociforo<sup>4</sup>, F. Nolden<sup>4</sup>, U. Popp<sup>4</sup>, C. Rigollet<sup>2</sup>, S. Roy<sup>2</sup>, C. Scheidenberger<sup>4</sup>, M. Steck<sup>4</sup>, B. Streicher<sup>2,4</sup>, L. Stuhl<sup>3</sup>, M. Thürauf<sup>1</sup>, T. Uesaka<sup>9</sup>, H. Weick<sup>4</sup>, J. Winfield<sup>4</sup>, D. Winters<sup>4</sup>, P.J. Woods<sup>10</sup>, T. Yamaguchi<sup>11</sup>, J.C. Zamora<sup>1</sup>, J. Zenihiro<sup>9</sup>

<sup>1</sup> Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

<sup>2</sup> Kernfysisch Versneller Instituut (KVI), Rijksuniversiteit Groningen, 9700 AB Groningen, Netherlands

<sup>3</sup> ATOMKI, Institute of Nuclear Research, Hungarian Academy of Sciences, 4026 Debrecen, Hungary

<sup>4</sup> GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

<sup>5</sup> Ioffe Physico-Technical Institute, Russian Academy of Sciences, St.Petersburg, 194021 Russia

<sup>6</sup> Physik-Department E12, Technische Universität München, 85748 Garching, Germany

<sup>7</sup> Institute of Modern Physics, Chinese Academy of Sciences, 730000 Lanzhou, China

<sup>8</sup> Department of Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan

<sup>9</sup> RIKEN Nishina Center, Wako, Saitama 351-0198, Japan

<sup>10</sup> Institute for Particle and Nuclear Physics, University of Edinburgh, Edinburgh, EH9 3JZ, United Kingdom

<sup>11</sup> Department of Physics, Saitama University, Saitama, Saitama 338-8570, Japan

Contact email: [schmid@ikp.tu-darmstadt.de](mailto:schmid@ikp.tu-darmstadt.de)

EXL (EXotic nuclei studied in Light-ion induced reactions at the NESR storage ring) is a project within NUSTAR [1] at FAIR [2]. It aims for the investigation of light-ion induced direct reactions in inverse kinematics with radioactive ions cooled and stored in the future NESR (New Experimental Storage Ring). A universal detector system will be built around an internal target of the NESR in order to detect the target-like recoils. One of the key interests of EXL is the investigation of reactions at very low momentum transfers where, for example, the nuclear matter distribution, giant monopole resonances or Gamow-Teller transitions can be studied [3].

The existing ESR (Experimental Storage Ring) at GSI, together with its internal gas-jet target, provides a unique opportunity to perform this kind of experiments on a smaller scale already today. In the last years we have developed a UHV compatible detector setup mainly based on DSSDs (Double-sided Silicon-Strip Detector) for the target-like recoils [4] and an in-ring detection system for the projectile-like heavy ions. With this setup we were able to successfully investigate reactions with a stored radioactive beam for the very first time. As a part of the first EXL campaign we investigated the reaction  $^{56}\text{Ni}(p,p)^{56}\text{Ni}$  in order to measure the differential cross section for elastic proton scattering and deduce the nuclear matter distribution of  $^{56}\text{Ni}$ . The present contribution will present the current status of the project and preliminary results.

This work was supported by BMBF (06DA9040I und 05P12RDFN8) and HIC for FAIR.

[1] <http://www.fair-center.eu/for-users/experiments/nustar.html>

[2] <http://www.fair-center.eu>

[3] FAIR Baseline Technical Report, GSI Darmstadt (2006), <https://www-alt.gsi.de/documents/DOC-2006-Jul-44-1.pdf>

[4] B. Streicher et al, Nucl. Instr. And Meth. A 654 (2011) 604