



Contribution ID: 73

Type: Poster

P59 - Enhanced RBE of submicron focused low LET protons

Friday, 11 July 2014 13:00 (1 hour)

Due to their physical and radiobiological properties, in particular their increased relative biological effectiveness (RBE), high linear energy transfer (LET) radiation qualities are of special interest for tumour therapy. To study the influence of spatial dose distribution on the biological effectiveness we use the ion microprobe SNAKE to concentrate the deposited dose in cells by focusing a certain number of low LET protons or ions to a submicron spot which approximate a high LET dose distribution. By changing the spot size and the number of focused ions one has opened a wide field to study RBE effects in dependence of dose distributions on submicron scale.

For our experiments we compare homogeneous proton irradiation with spots of focused 20 MeV protons (LET in water: 2.65 keV/μm) or 45 MeV Lithium ions (LET in water: 60 keV/μm) in a matrix pattern with certain numbers of ions per matrix point resulting in a constant dose of 1.7 Gy. For induction of chromosome aberrations we observed an increased effectiveness for higher number of ions per point and thus for a higher ionisation load per spot. 1.7 Gy irradiation with 541 protons per spot increases induction of dicentric by a factor of 2.7 with respect to homogeneous proton application. However, the ionisation load per spot, the product of LET times the applied ions per spot, is not enough to understand RBE. At constant ionisation load at a single point of the matrix the effectiveness for the yield for dicentric increases with LET. At the same mean dose of 1.7 Gy and the same matrix pattern 5 Li ions increase the yield for dicentric by 45%, one Carbon (LET in water: 310 keV/μm) ion by 130%, with respect to 117 protons per spot.

The results indicate that at least two effects cause the increased RBE of high LET particles. On the one hand the focused dose application leads to a locally enhanced DNA double strand break (DSB) density, which increases the probability for misrepair, e.g. joining wrong ends of DSB. On the other hand the very high doses in the core of a heavy ion lead to a higher amount or more complex DNA damage, e.g. due to nonlinear effect in induction of DSB.

Supported by the DFG-Cluster of Excellence 'Munich-Centre for Advanced Photonics' and by Bmbf-project 02NUK031A "LET-Verbund"

Primary author: Mr GREUBEL, Christoph (Institut für Angewandte Physik und Messtechnik, Universität der Bundeswehr München, Germany)

Co-authors: SIEBENWIRTH, Christian (Department of Radiation Oncology, Technische Universität München, Germany); WALSH, Dietrich (Institut für Angewandte Physik und Messtechnik, Universität der Bundeswehr München, Germany); SCHMID, Ernst (Institute for Cell Biology, Ludwig-Maximilians Universität München, Germany); MULTHOFF, Gabriele (Department of Radiation Oncology, Technische Universität München, Germany); REINDL, Judith (Institut für Angewandte Physik und Messtechnik, Universität der Bundeswehr München, Germany); ILICIC, Katarina (Department of Radiation Oncology, Technische Universität München, Germany); ZLOBINSKAYA, Olga (Department of Radiation Oncology, Technische Universität München, Germany); GIRST, Stefanie (Institut für Angewandte Physik und Messtechnik, Universität der Bundeswehr München, Germany); SCHMID, Thomas E. (Department of Radiation Oncology, Technische Universität München, Germany)

Presenter: Mr GREUBEL, Christoph (Institut für Angewandte Physik und Messtechnik, Universität der Bundeswehr München, Germany)

Session Classification: Poster Session with Cheese and Wine