

QCD at the LHC

Recent progress and open problems

Juan Rojo
CERN, PH Division, TH Unit

Les Rencontres de Physique de la Vallée d'Aoste 2014
La Thuile, 25/02/2014

QCD: The Toolbox for Discoveries at the LHC


The days of “guaranteed” discoveries or of no-lose theorems in particle physics are over, at least for the time being

.... but the big questions of our field remain wild open (hierarchy problem, flavour, neutrinos, DM, BAU,) **Mangano, Aspen14**

This simply implies that, more than for the past 30 years, future HEP’s progress is to be driven by experimental exploration, possibly renouncing/reviewing deeply rooted theoretical bias

Improving our **quantitative understanding of the Standard Model** is essential in this new era for HEP, where we need to hunt, unbiased, for **answers to the big questions of our field**

Now, more than ever, **sharpening our QCD tools** could be the **key for new discoveries at the LHC**




Prime example: **extraction of Higgs couplings** from LHC data soon to be **limited by QCD uncertainties**

Better QCD predictions


Improved indirect **sensitivity to New Physics** via deviations of Higgs couplings from SM expectations

QCD at the LHC




Drawing by K. Hamilton

QCD at the LHC




Drawing by K. Hamilton

QCD at the LHC




Drawing by K. Hamilton

QCD at the LHC




Drawing by K. Hamilton

QCD at the LHC



Drawing by K. Hamilton

QCD at the LHC



Drawing by K. Hamilton

The inner life of the proton

The **Master Formula** for LHC cross-sections:


$$\sigma_X(s, M_X^2) = \sum_{a,b} \int_{x_{\min}}^1 dx_1 dx_2 f_{a/h_1}(x_1, M_X^2) f_{b/h_2}(x_2, M_X^2) \hat{\sigma}_{ab \rightarrow X}(x_1 x_2 s, M_X^2)$$

Parton Distributions:


- ☑ Energy distribution of quarks and gluons in proton
- ☑ Determined by non-perturbative dynamics
- ☑ Extract from *experimental data* + pQCD evolution

Matrix Elements:

- ☑ Hard-scattering between quarks, gluons, electroweak bosons, Higgs
- ☑ Compute in perturbation theory as series expansion in α_s , α_{QED} , α_{EW}




Experimental data



QCD Theory

Statistical Methodology


PDFs




Predictions @ LHC

Parton Distributions: fundamental limit to theory predictions at LHC

Parton Distributions and LHC phenomenology




1) PDFs fundamental limit for Higgs boson characterization in terms of couplings



Juan Rojo

2) Very large PDF uncertainties ($>100\%$) for new heavy particle production



3) PDFs dominant systematic for precision measurements, like W boson mass, that test internal consistency of the Standard Model

La Thuile, 25/02/2014

PDFs and LHC data

- ✓ A major recent development in global PDF fits is the **inclusion of constraints from LHC data**
- ✓ The impact of **new data into PDFs** has been also studied by ATLAS and CMS themselves using the open-source QCD analyses framework **HERAFitter**


LHC data already included in PDF fits:

- ✓ *Inclusive W,Z production*
- ✓ *W production with charm quarks*
- ✓ *Isolated photon production*
- ✓ *Inclusive jet and dijet production*
- ✓ *Low and high-mass off-shell Drell-Yan*
- ✓ *Top quark pair cross-sections*
- ✓ *Ratios of cross-sections between different E_{cm}*

LHC data with potential PDF constraints

- ✓ *Z+jets, high- p_T Z production*
- ✓ *Photon+jet production*
- ✓ *Photon+charm, Z+charm*
- ✓ *Single top production*
- ✓ *Top quark pair differential distributions*
- ✓ *Ratios between 13 and 8 TeV*


Pinning down large-x gluon with top quark data




Czakon et al 13

Juan Rojo

Gluon PDF from ratio of
7 TeV and 2.76 TeV jet data






W+charm: accurate strangeness from LHC data




La Thuile, 25/02/2014

Theory Developments on PDFs

- Consistent inclusion of **QED** effects in LHC calculations require PDFs with **QED corrections**, and in particular a determination of the **photon PDF** from experimental data (**NNPDF 13**, see **S. Carrazza talk**)




- Impact of **Fixed-Flavor** vs **Variable-Flavor-Number** heavy quark schemes on PDFs (**NNPDF13**, **Thorne 14**)




Use of different heavy flavor schemes responsible for part differences between some PDF sets
Fits in the FFN worse fit quality to DIS data than VFN fits

- Parton Distributions with **Intrinsic Charm** PDF (**CT 13**)

Intrinsic charm still allowed to carry up to 2% of the proton momentum
Accessible at LHC via photon+charm and Z+charm data



QCD at the LHC




Drawing by K. Hamilton

The NLO revolution

- During many years, the needs for NLO calculations were summarized in the Les Houches wishlist
- The NLO revolution in the last years makes computations of NLO cross-sections a solved process
- Key has been the automation of NLO real emission and subtraction and of virtual corrections (**MadFKS, MadLoop, GoSam, Sherpa, OpenLoops, HelacNLO, ...**)
- Despite automation, for high final state multiplicities, tailored calculations still required for efficiency (**BlackHat, NJet, Rocket,**)

Process ($V \in \{Z, W, \gamma\}$)	Comments
1. $pp \rightarrow VV \text{ jet}$	WW jet completed by Dittmaier/Kallweit/Uwer; Campbell/Ellis/Zanderighi ZZ jet completed by Binoth/Gleisberg/Karg/Kauer/Sanguinetti
2. $pp \rightarrow \text{Higgs}+2 \text{ jets}$	WZ jet, $W\gamma$ jet completed by Campanario et al. NLO QCD to the gg channel completed by Campbell/Ellis/Zanderighi NLO QCD+EW to the VBF channel completed by Ciccolini/Denner/Dittmaier Interference QCD-EW in VBF channel
3. $pp \rightarrow VVV$	ZZZ completed by Lazopoulos/Melnikov/Petriello and WWZ by Hankele/Zeppenfeld see also Binoth/Ossola/Papadopoulos/Pittau VBFNLO meanwhile also contains WWW, ZZW, ZZZ, WW γ , ZZ γ , WZ γ , W $\gamma\gamma$, Z $\gamma\gamma$, $\gamma\gamma\gamma$, W $\gamma\gamma j$
4. $pp \rightarrow t\bar{t} b\bar{b}$	relevant for $t\bar{t}H$, computed by Bredenstein/Denner/Dittmaier/Pozzorini and Bevilacqua/Czakon/Papadopoulos/Pittau/Worek
5. $pp \rightarrow V+3 \text{ jets}$	W+3 jets calculated by the Blackhat/Sherpa and Rocket collaborations Z+3 jets by Blackhat/Sherpa
6. $pp \rightarrow t\bar{t}+2 \text{ jets}$	relevant for $t\bar{t}H$, computed by Bevilacqua/Czakon/Papadopoulos/Worek
7. $pp \rightarrow VV b\bar{b}$	Pozzorini et al. Bevilacqua et al.
8. $pp \rightarrow VV+2 \text{ jets}$	$W^+W^++2 \text{ jets}, W^+W^-+2 \text{ jets}$, relevant for VBF $H \rightarrow VV$ VBF contributions by (Bozzi/Jäger/Oleari/Zeppenfeld Binoth et al.
9. $pp \rightarrow b\bar{b}b\bar{b}$	
10. $pp \rightarrow V+4 \text{ jets}$	top pair production, various new physics signatures Blackhat/Sherpa: W+4 jets, Z+4 jets see also HEJ for W+ n jets
11. $pp \rightarrow Wb\bar{b}j$	top, new physics signatures, Reina/Schutzmeier
12. $pp \rightarrow t\bar{t}t\bar{t}$	various new physics signatures, Bevilacqua/Worek

Current frontier of NLO calculations
pp \rightarrow W + 5 jets @ NLO, BlackHat 13



NLO crucial for reliable scale uncertainties

The NNLO revolution

📍 Until recently, few processes were known **differentially at NNLO**, in particular only processes with either **colorless initial state** or **colorless final state**

Process	Calculation	Relevance
pp -> H	Anastasiou, Melnikov, Petriello Catani, Grazzini	Higgs production
pp -> V	Melnikov, Petriello Catani, Cieri, de Florian, Ferrera, Grazzini	Electroweak precision tests Quark flavor separation
e+e- -> 3 jets	Gerhman, Glover, Heinrich	Fits of α_s
pp -> gamma gamma pp -> VH	Catani, Ferrera, Grazzini, Tramontano	Background to Higgs production Higgs associated production


📍 The development of new calculational techniques, *Antenna Subtraction* and *Sector-Improved subtraction*, lead to the **2013 NNLO breakthrough**: it is now possible to compute NNLO QCD corrections to processes with both **colored initial and final states**

Process	Calculation	Relevance
pp -> tt	Czakon, Fiedler, Mitov	Precision studies of top sector Large-x gluon PDF
gg -> dijets	Gehrmann-De Ridder, Gehrmann, Glover, Pires	Background to New Physics Gluon PDF + alphas fits
pp -> H + jets	Boughezal, Caola, Melnikov, Petriello, Schulze	Higgs production in association with hard jets


Other recent NNLO calculations include **pp -> HH** in gluon fusion (**De Florian, Mazzitelli 13**), and **pp -> HHjj VBF** (**Liu-Sheng et al 14**)

The NNLO revolution

NNLO dijets: large perturbative corrections



$t\bar{t}$ bar: scale uncertainties now at 2-3% level



- ⦿ Whats in the pipeline? We have new **Les Houches wishlist**, now for **NNLO + EWK calculations**
- ⦿ **NNLO is crucial for many precision measurements**, expect lots of rapid progress in the following years

Process	known	desired	details
$t\bar{t}$	σ_{tot} @ NNLO QCD $d\sigma(\text{top decays})$ @ NLO QCD $d\sigma(\text{stable tops})$ @ NLO EW	$d\sigma(\text{top decays})$ @ NNLO QCD + NLO EW	precision top/QCD, gluon PDF, effect of extra radiation at high rapidity, top asymmetries
$t\bar{t} + j$	$d\sigma(\text{NWA top decays})$ @ NLO QCD	$d\sigma(\text{NWA top decays})$ @ NNLO QCD + NLO EW	precision top/QCD top asymmetries
single-top	$d\sigma(\text{NWA top decays})$ @ NLO QCD	$d\sigma(\text{NWA top decays})$ @ NNLO QCD (t channel)	precision top/QCD, V_{tb}
dijet	$d\sigma$ @ NNLO QCD (g only) $d\sigma$ @ NLO weak	$d\sigma$ @ NNLO QCD + NLO EW	Obs.: incl. jets, dijet mass \rightarrow PDF fits (gluon at high x) $\rightarrow \alpha_s$ CMS http://arxiv.org/abs/1212.6660
3j	$d\sigma$ @ NLO QCD	$d\sigma$ @ NNLO QCD + NLO EW	Obs.: $R3/2$ or similar $\rightarrow \alpha_s$ at high scales dom. uncertainty: scales CMS http://arxiv.org/abs/1304.7498
$\gamma + j$	$d\sigma$ @ NLO QCD $d\sigma$ @ NLO EW	$d\sigma$ @ NNLO QCD +NLO EW	gluon PDF $\gamma + b$ for bottom PDF

*Differential NNLO calculations
bring QCD to a new level of
precision at LHC*

QED and Electroweak corrections


- ✓ At present level of precision in QCD calculations, **electroweak corrections** become **comparable if not larger**
- ✓ **Electroweak Sudakov logarithms** grow with energy, more important at LHC 13 TeV

Typical impact on $2 \rightarrow 2$ reactions at $\sqrt{s} \sim 1$ TeV:

$$\delta_{LL}^{1\text{-loop}} \sim -\frac{\alpha}{\pi s_W^2} \ln^2\left(\frac{s}{M_W^2}\right) \simeq -26\%, \quad \delta_{NLL}^{1\text{-loop}} \sim +\frac{3\alpha}{\pi s_W^2} \ln\left(\frac{s}{M_W^2}\right) \simeq 16\%$$

- ✓ Electroweak corrections affect the **TeV scale phenomenology**, both for **New Physics searches** in the high-mass tails, **Higgs characterization** and **precision SM measurements**, such as PDF fits

Electroweak corrections to high- p_T jets @ LHC8



QED photon-induced and EW effects in high-mass Drell-Yan




TABLE V: Are we in the Sudakov zone yet?


Process	$\sqrt{s} = 8$ TeV	$\sqrt{s} = 14$ TeV	$\sqrt{s} = 33, 100$ TeV
Inclusive jet, dijet	Yes	Yes	Yes
Inclusive W/Z tail	\sim Yes	Yes	Yes
$W\gamma, Z\gamma$ tail ($l\nu\gamma, ll\gamma$)	No	\sim Yes	Yes
W/Z+jets tail	\sim Yes	Yes	Yes
WW leptonic	Close	\sim Yes	Yes
WZ, ZZ leptonic	No	No	Yes

☛ The region where EW corrections become relevant known as the **Sudakov zone**

☛ At LHC 13 TeV, many **crucial processes** will require these EW corrections (see review in [arxiv:1308.1430](https://arxiv.org/abs/1308.1430))

☛ **PDF sets** which include **non only QED, but also electroweak corrections**, are required for consistent implementation of EW effects


QCD at the LHC



Drawing by K. Hamilton

NLO parton showers


- ✓ **Fixed order calculation** do not provide a realistic description of final states in hadronic collisions
- ✓ They need to be supplemented with **parton showers, all-order resummations** of QCD soft and collinear radiation (**Pythia6/8, Herwig/++, Sherpa, Ariadne,**)
- ✓ In addition, **merging matrix elements** with high multiplicity improves final state description
- ✓ Matching to parton showers trivial at LO. LO merging requires **prescriptions to avoid double counting**



Leading Order
QCD jet production

NLO parton showers

- ✓ **Fixed order calculation** do not provide a realistic description of final states in hadronic collisions
- ✓ They need to be supplemented with **parton showers, all-order resummations** of QCD soft and collinear radiation (**Pythia6/8, Herwig/++, Sherpa, Ariadne ...**)
- ✓ In addition, **merging matrix elements** with high multiplicity improves final state description
- ✓ Matching to parton showers trivial at LO. LO merging requires **prescriptions to avoid double counting**




Leading Order +
Parton Shower
QCD jet production

NLO parton showers

- ✓ **Fixed order calculation** do not provide a realistic description of final states in hadronic collisions
- ✓ They need to be supplemented with **parton showers, all-order resummations** of QCD soft and collinear radiation (**Pythia6/8, Herwig/++, Sherpa, Ariadne, ...**)
- ✓ In addition, **merging matrix elements** with high multiplicity improves final state description
- ✓ Matching to parton showers trivial at LO. LO merging requires **prescriptions to avoid double counting**


Leading Order + Parton Shower + Merging
QCD jet production



LO merging requires a prescription to avoid double counting
CKKW, Catani, Krauss, Kuhn, Webber 02
MLM, Mangano 02


NLO parton showers

- ✓ At NLO matching to parton showers non-trivial, requires either i) modify / veto the shower first emission or ii) subtract from the NLO the first shower emission, to avoid **double counting**
- ✓ Two main methods: MCatNLO (Frixione, Webber 02) and POWHEG (Nason 04, Alioli, Oleari, Nason, Re 10) are of common use. These approaches are now largely automated: aMCatNLO, POWHEG-Box, also in Sherpa, Herwig++, ...
- ✓ NLO+PS calculations now available for virtually all relevant LHC process



A NLO+PS matched calculation provides improved description of a wider range of final state configurations that NLO or PS alone

Multileg NLO+PS Merging



✓ The current frontier in NLO+PS is the merging of matched NLO+PS samples with different multiplicities into a common sample

✓ As illustration, consider H+jet in the FxFx merging approach (Frederix, Frixione 12)

✓ NLO+PS H+0jet treats the extra jet at LO (fails at high jet pt)

✓ NLO+PS H+1jet misses the bulk of cross section, which comes from events with no hard radiation


✓ The merged sample successfully interpolated between the two regimes avoiding double counting

✓ Various different approaches have been proposed: FxFx, MEPS@NLO (Sconherr, Hoeche, Krauss, Siebert 13), UNLOPS (Lonnblad and Prestel 12), ...


✓ Multileg NLO+PS merging should become the standard for realistic NLO Monte Carlo simulations in the following years

Towards NNLO matched to parton showers

- ✓ Matching **fully differential NNLO** calculations to **parton showers** would provide the ultimate accuracy to QCD simulations at the LHC
- ✓ Many new conceptual issues need to be tackled, but already encouraging progress (**Hamilton, Nason, Re, Zanderighi 13, Alioli et al. (GENEVA) 13,**)




QCD at the LHC



Drawing by K. Hamilton

Jets at LHC

Jets are ubiquitous at LHC, and required for almost all analysis from SM measurements, Higgs physics and BSM searches. Paradigm is Anti-kt jets (Cacciari, Salam, Soyez 08) with radius R in a range between 0.4 and 0.7. Virtually all jet reconstruction tools available in the FastJet framework (Cacciari, Salam, Soyez)




Jets@LHC



- Standard Model: PDF determination, extraction of α_s , top quark reconstruction, hadronic V decays,
- Higgs physics: discrimination between production models, hadronic Higgs decays (bb, cc), associated production,
- Beyond the Standard Model: searches for compositeness, supersymmetry in the jets + missing ET, TeV scale gravity via quantum black holes, jet substructure

Jet Veto Efficiency
in $gg \rightarrow H$



Recently a lot of effort has been done in understanding the (large) theoretical uncertainties associated to vetoing jets, as done for instance in Higgs analysis to separate gluon fusion from VBF

Resummed NNLO+NNLL calculations allow to reduce higher-order uncertainties in the jet veto efficiency as function of the jet transverse momentum

Banfi, Monni, Salam, Zanderighi 12-13

Becher, Neubert 12-13


Tackmann, Walsh, Zuberi 13

Liu, Petriello 12

La Thuile, 25/02/2014

Jet substructure

- In the decays of a **massive enough resonances**, **boosted prongs** can often be collimated into a **single jet**
- Different **jet substructure** in these jets and QCD jets provide strong background suppression in **BSM searches**




QCD jet

Boosted W jet

Boosted top quark jet


- LHC analysis are using more and more jet substructure techniques (also for Higgs)
- As illustration, recent CMS search for $q^* \rightarrow qV$ in the **tagged dijet final state**
- Discriminating variable: **different jet shape/mass in signal and in QCD background**

CMS PAS-EXO-12-024



Boosted W jet


Juan Rojo



La Thuile, 25/02/2014

Jet substructure



Wide variety to **jet substructure tools**, useful to **sharpen interesting signals** and to **reduce QCD backgrounds**



See the **proceedings of the BOOST workshops** for a complete set of references

[arxiv:1311.2708](https://arxiv.org/abs/1311.2708)

[arxiv:1012.5412](https://arxiv.org/abs/1012.5412)



However, it should be avoided to **use blindly** these tools.

It is essential to back then not only with **Monte Carlo studies** but also with **analytical calculations** (Dasgupta, Fregoso, Marzani, Salam 13)

These calculations in turn show to to **further improve jet taggers**

Note that even for something as fundamental as **jet mass**, different taggers lead to quite different results, and also a dependence with the scale $m^2/p_t^2 R^2$

Summary

- 🔧 **Quantum Chromodynamics** is an **essential ingredient** of the **LHC physics program**
- 🔧 **Precision QCD calculations** are required for most LHC analysis, from **Higgs boson characterization**, **searches for new massive particles** to the **precision determination of Standard Model parameters**
- 🔧 Huge progress in QCD in the last years including:
 - ☑ *Robust, statistically meaningful PDFs with LHC data and QED corrections*
 - ☑ *NNLO calculations for many LHC process including partons in both initial and final state*
 - ☑ *Automation of NLO matched to parton shower calculations and multileg NLO+PS merging*
 - ☑ *Precision jet reconstruction including resummed calculations, and standardization of jet tools in the FastJet framework*
 - ☑ *New taggers for jet substructure, and improved analytical understanding of these*
- 🔧 And much more to come, to match the requirements of LHC data, so stay tuned!

Summary

- 📌 Quantum Chromodynamics is an essential ingredient of the LHC physics program
- 📌 Precision QCD calculations are required for most LHC analysis, from Higgs boson characterization, searches for new massive particles to the precision determination of Standard Model parameters
- 📌 Huge progress in QCD in the last years including:
 - ✓ Robust, statistically meaningful PDFs with LHC data and QED corrections
 - ✓ NNLO calculations for many LHC process including partons in both initial and final state
 - ✓ Automation of NLO matched to parton shower calculations and multileg NLO+PS merging
 - ✓ Precision jet reconstruction including resummed calculations, and standardization of jet tools in the FastJet framework
 - ✓ New taggers for jet substructure, and improved analytical understanding
- 📌 And much more to come, to match the requirements

and apologies for the missing references....

**Thanks for your
attention!**