



Contribution ID: 43

Type: **Poster**

## **PS3-21 Interaction of Ions with Graphene on a Substrate**

*Thursday, October 9, 2014 5:00 PM (1h 30m)*

In this paper analyzes the process of dynamic screening of external charges, moving over the surface of graphene disposed on a substrate. We are considered here relatively slow ions with velocities from thermal speeds to Bohr velocity. We explore here the effects of dynamic polarization of graphene on a substrate by ion motion parallel to its surface. We limited the study of the effects associated with the presence of an insulating substrate, such as , with a dielectric constant of . Ishigami et al. [1] have found that the distance  $h$  between graphene and substrate is on the order of the distance between graphene layers in graphite or even larger. Only (according to our information) in the paper [2] considered the effects of finite  $h$ . Our principal goal was to demonstrate how strong are the effects of finite  $h$  on the degree of dynamic polarization of graphene moving ions. This meant the need to include the gap size graphene-substrate as a parameter in modeling phenomena screening graphene. The simulation was performed by molecular dynamics using ReaxFF potential of free package for classical molecular dynamics LAMMPS (Large-scale Atomic/Molecular Massively Parallel Simulator).

### References

- [1] M. Ishigami, J.H. Chen, W.G. Cullen, M.S. Fuhrer, E.D. Nano Lett. (2007) 7, 1643.
- [2] I. Radović, Lj. Hadžievski PHYSICAL REVIEW (2008) 77, 075428.

**Primary author:** Mr IVANOV, Aleksei (Russian)

**Presenter:** Mr IVANOV, Aleksei (Russian)

**Session Classification:** Poster Session