## **Channeling 2014**



Contribution ID: 40

Type: Poster

## PS3-23 Hydrogen Atoms Channeling through Carbon Nanotubes

Thursday, 9 October 2014 17:00 (1h 30m)

Using molecular dynamics method with Brenner interatomic potential [1], we studied the channeling of ideally collimated beam of hydrogen atoms (1000 atoms, 100-250eV/atom) through carbon nanotubes (CNT). CNT with diameter of 13.6 angstroms were placed in one plane, parallel to each other at a distance apart 3 angstroms. Hydrogen atoms are directed along the axes of the tubes. During H interaction with CNT a part of H atoms is scattered by the face of the tube, some of them (having undergone one or more collisions inside the tube) are dissipated through the wall, and other part of H atoms is channeled through tubes. We studied these channeled atoms. As results show that these atoms consist of 40-53% of the total number of H atoms.

We have analyzed the angular distributions of H atoms channeled through the tubes. These H atoms are slightly deflected from an initial direction, basically, in the range from 0 to 20 degrees. 20-25% (of total) atoms is scattered in the range from 0 to 1 degree. 20-28% (of total) atoms is scattered, mainly, in the range from 1 to 20 degrees. Distribution is continuous with main peak at 0 degree. With sequential increase of the energy of the incident beam (from 100 to 250 eV/atom), the number of hydrogen atoms channeled through the CNTs is increased, however, the range of scattering angles remained the same - 0-20 degrees.

Setting a screen at some distance (100-1000 angstroms) from the ends of the nanotubes, we got a projection of H atoms on the screen after channeling them through CNT. Diameter of these projections is 10-11 angstroms, which is slightly smaller than the diameter of nanotubes (13.6 angstroms). Diameter of these projections does not depend on the energy of beam (in 100-250eV/atom range). Dispersion of these projections is not significant and the reason of this is the small scattering angles of atoms channeled through the CNTs.

## References

1. D.W. Brenner, O.A. Shenderova, J.A. Harrison, S.J. Stuart, B. Ni, S.B. Sinnot. "A second-generation reactive empirical bond order (REBO) potential energy expression for hydrocarbons", J.Phys: Condens. Matter, 14, pp.783-802 (2002)

**Primary author:** YADGAROV, Ishmumin (1Institute of Ion-Plasma and Laser Technologies, Tashkent, Uzekistan)

**Co-authors:** Prof. DZHURAKHALOV, Abdiravuf (Dept. of Mathematics and Computer Sciences, University of Antwerp, Antwerp, Belgium); ALYABEV, Danila (Institute of Ion-Plasma and Laser Technologies)

Presenter: YADGAROV, Ishmumin (1Institute of Ion-Plasma and Laser Technologies, Tashkent, Uzekistan)

Session Classification: Poster Session