



Contribution ID: 72

Type: Invited talk

## Spectral Characteristics of Radiation from Thomson and Compton Scattering of an Intense Laser Field by Relativistic Electrons

Thursday, October 9, 2014 9:00 AM (30 minutes)

The mechanisms of Thomson and Compton scattering under linear and nonlinear interactions between relativistic electrons and counter-propagated intense laser wave are considered. The quantum consideration of the Compton scattering process allows to calculate a probability of a few successive collisions  $k$  of an electron with laser photons accompanying by absorption of  $n$  photons (nonlinear regime for  $n$  more than unit) when the number of collisions and the number of absorbed photons are random quantities. The cross-section of the nonlinear Thomson scattering process was obtained from the classical formula for intensity using the Planck's law. An electron interacts with a few laser photons subsequently, emitting a "hard photon" in each collision if a laser pulse intensity is high enough (multiple Compton scattering process, MCS) [1]. A mean number of emitted photons  $\kappa$  is determined by a luminosity of the MCS process and its cross-section. We have showed that spectra of emitted photons can be described by the classical Thomson formula if the condition  $4\kappa\gamma\hbar\omega/E \ll 1$  ( $\gamma$  is Lorentz-factor,  $\hbar\omega$  is energy of laser photon,  $E$  is electron rest energy) is fulfilled. In opposite case one has to use the Compton cross-section formula.

We developed an approach based on Monte-Carlo technique allowing to simulate spectral distributions of photons emitted into a narrow aperture for both linear and nonlinear MCS processes. In contrast with other models we took into account a multiplicity of collisions of an electron with laser photons and showed that monochromaticity of radiation is worsening due to such a reason. In each collision an electron loses an energy during emission process and a subsequent photon will have an energy less than the first one (in the average). For modern projects (such as the ELI-NP [2, 3]) the average number of emitted photons by each electron can enhance the value  $\kappa \geq 1$ .

For typical parameters of the ELI-NP project ( $\kappa = 1.1$ , the electron energy 720 MeV ( $\gamma = 1409$ ), acceptance angle 0.1 mrad) we have simulated the spectral line shape and found that the line has a "tail" of soft photons (about 20% from the total number of photons accepted into an aperture). Such an effect should be taken into account for design of monochromatic gamma-beam.

### References

- [1] A. Potylitsyn, A. Kol'chuzhkin, Nucl. Phys. and Meth. B 309 (2013) 15-19.
- [2] [www.extreme-light-infrastructure.eu](http://www.extreme-light-infrastructure.eu).
- [3] V. Petrillo, A. Bacci, R. Ben Ali Zinati et al., Nucl. Instrum. and Methods A 693 (2012) 109.

**Primary author:** Prof. POTYLITSYN, Alexander (Tomsk Polytechnic University)

**Co-author:** Prof. KOLCHUZHKIN, Anatoliy (Moscow State University of Technology Stankin)

**Presenter:** Prof. POTYLITSYN, Alexander (Tomsk Polytechnic University)

**Session Classification:** S2: Channeling & Radiation in Various Fields