Recent results from DAMA/ LIBRA-phase1 and perspectives





RICAP 2014 Noto (SR), Italy Sept 30th – Oct 3rd, 2014



DAMA set-ups an observatory for rare processes @ LNGS



- DAMA/LIBRA (DAMA/Nal)
- DAMA/LXe
- DAMA/R&D
- DAMA/Crys
- DAMA/Ge

Collaboration:

Roma Tor Vergata, Roma La Sapienza, LNGS, IHEP/Beijing
+ by-products and small scale expts.: INR-Kiev
+ in some studies on ββ decays (DST-MAE and Inter-Universities projects):
IIT Kharagpur and Ropar, India

Web Site: http://people.roma2.infn.it/dama

Some direct detection processes:



The annual modulation: a model independent signature for the investigation of DM particles component in the galactic halo

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions can point out its presence.

Requirements of the annual modulation

- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about 2 June)
- 5) Just for single hit events in a multidetector set-up
- 6) With modulation amplitude in the region of maximal sensitivity must be <7% for usually adopted halo distributions, but it can be larger in case of some possible scenarios



the DM annual modulation signature has a different origin and peculiarities (e.g. the phase) than those effects correlated with the seasons

To mimic this signature, spurious effects and side reactions must not only - obviously - be able to account for the whole observed modulation amplitude, but also to satisfy simultaneously all the requirements

The pioneer DAMA/NaI: ≈100 kg highly radiopure NaI(Tl)

Performances:

N.Cim.A112(1999)545-575, EPJC18(2000)283, Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

Results on rare processes:

- Possible Pauli exclusion principle violation
- CNC processes
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell)
- Search for solar axions
- Exotic Matter search
- Search for superdense nuclear matter
- Search for heavy clusters decays

Results on DM particles:

- PSD
- Investigation on diurnal effect
- Exotic Dark Matter search
- Annual Modulation Signature

PLB408(1997)439 PRC60(1999)065501

PLB460(1999)235 PLB515(2001)6 EPJdirect C14(2002)1 EPJA23(2005)7 EPJA24(2005)51





PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512, PLB480(2000)23, EPJC18(2000)283, PLB509(2001)197, EPJC23(2002)61, PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)2127, IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)023506, MPLA23(2008)2125

Model independent evidence of a particle DM component in the galactic halo at 6.3 σ C.L.

total exposure (7 annual cycles) 0.29 ton×yr

The DAMA/LIBRA set-up ~250 kg NaI(Tl) (Large sodium Iodide Bulk for RAre processes)



As a result of a 2nd generation R&D for more radiopure NaI(TI) by exploiting new chemical/physical radiopurification techniques (all operations involving - including photos - in HP Nitrogen atmosphere)



Residual contaminations in the new DAMA/LIBRA Nal(TI) detectors: ²³²Th, ²³⁸U and ⁴⁰K at level of 10⁻¹² g/g







 Radiopurity, performances, procedures, etc.: NIMA592(2008)297, JINST 7 (2012) 03009
 Results on DM, Annual Modulation Signature: EPJC56(2008)333, EPJC67(2010)39, EPJC73(2013)2648 related results: PRD84(2011)055014, EPJC72(2012)2064, IJMPA28 (2013)1330022, EPJC74(2014)2827, arXiv:1409.3516
 Rare processes: PEP viol.: EPJC62(2009)327, CNC in I: EPJC72(2012)1920, IPP in Am: EPJA49(2013)64

DAMA/LIBRA calibrations

Low energy: various external gamma sources (²⁴¹Am, ¹³³Ba) and internal X-rays or gamma's (⁴⁰K, ¹²⁵I, ¹²⁹I), routine calibrations with ²⁴¹Am







High energy: external sources of gamma rays (e.g. ¹³⁷Cs, ⁶⁰Co and ¹³³Ba) and gamma rays of 1461 keV due to ⁴⁰K decays in an adjacent detector, tagged by the 3.2 keV X-rays



Thus, here and hereafter keV means keV electron equivalent

Complete DAMA/LIBRA-phase1

	Period	Mass (kg)	Exposure (kg×day)	$(lpha - eta^2)$
DAMA/LIBRA-1	Sept. 9, 2003 - July 21, 2004	232.8	51405	0.562
DAMA/LIBRA-2	July 21, 2004 - Oct. 28, 2005	232.8	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 - July 18, 2006	232.8	39445	0.591
DAMA/LIBRA-4	July 19, 2006 - July 17, 2007	232.8	49377	0.541
DAMA/LIBRA-5	July 17, 2007 - Aug. 29, 2008	232.8	66105	0.468
DAMA/LIBRA-6	Nov. 12, 2008 - Sept. 1, 2009	242.5	58768	0.519
DAMA/LIBRA-7	Sep. 1, 2009 - Sept. 8, 2010	242.5	62098	0.515
DAMA /I IBDA phase1	Sept 0 2002 Sept 8 2010		$270705 \approx 1.04$ ton yum	0 518
DAMA/N-L+DAMA/I	1.04 toll x yr	0.518		
[DAMA/NaI + DAMA/L	1.33 ton×yr			

a ton × yr experiment? done

EPJC56(2008)333, EPJC67(2010)39 EPJC73(2013)2648

- calibrations: ≈96 M events from sources
- acceptance window eff: 95 M events (≈3.5 M events/keV)





- replacement of some PMTs in HP N_{2} atmosphere
- restore 1 detector to operation
- new Digitizers installed (U1063A Acqiris 1GS/s 8-bit High-Speed cPCI)
- new DAQ system with optical read-out installed

START of DAMA/LIBRA – phase 2

- Second upgrade on Oct./Nov. 2010
- Replacement of all the PMTs with higher Q.E. ones from dedicated developments
- \diamond Goal: lowering the software energy threshold

Fall 2012: new preamplifiers installed + special trigger modules. Other new components in the electronic chain in development

... continuously running

Model Independent Annual Modulation Result DAMA/Nal + DAMA/LIBRA-phase1 Total exposure: 487526 kg×day = 1.33 ton×yr



The data favor the presence of a modulated behaviour with all the proper features for DM particles in the galactic halo at about 9.2 σ C.L.

Energy distribution of the modulation amplitudes

Maximum-likelihood analysis

 $R(t) = S_0 + S_m \cos[\omega(t - t_0)]$ T=2\pi/\omega=1 yr t_0= 152.5 day

DAMA/NaI + DAMA/LIBRA-phase1

total exposure: 487526 kg×day ≈**1.33 ton×yr**



A clear modulation is present in the (2-6) keV energy interval, while S_m values compatible with zero are present just above

The S_m values in the (6–20) keV energy interval have random fluctuations around zero with χ^2 equal to 35.8 for 28 degrees of freedom (upper tail probability 15%)



Summary of the results obtained in the additional investigations of possible systematics or side reactions – DAMA/LIBRA-phase1

(NIMA592(2008)297, EPJC56(2008)333, J. Phys. Conf. ser. 203(2010)012040, arXiv:0912.0660, S.I.F.Atti Conf.103(211), Can. J. Phys. 89 (2011) 11, Phys.Proc.37(2012)1095, EPJC72(2012)2064, arXiv:1210.6199 & 1211.6346, IJMPA28(2013)1330022)

Source	Main comment	Cautious upper limit (90%C.L.)	
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	<2.5×10 ⁻⁶ cpd/kg/keV	
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	<10 ⁻⁴ cpd/kg/keV	
NOISE	Effective full noise rejection near threshold	<10 ⁻⁴ cpd/kg/keV	
ENERGY SCALE	Routine + intrinsic calibrations	<1-2 ×10 ⁻⁴ cpd/kg/keV	
EFFICIENCIES	Regularly measured by dedicated calibrations	<10 ⁻⁴ cpd/kg/keV	
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV <i>multiple-hits</i> events; this limit includes all possible sources of background	<10 ⁻⁴ cpd/kg/keV	
SIDE REACTIONS	Muon flux variation measured at LNGS	<3×10 ⁻⁵ cpd/kg/keV	

+ they cannot satisfy all the requirements of annual modulation signature Thus, they cannot mimic the observed annual modulation effect

No role for μ in DAMA annual modulation result

✓ Direct μ interaction in DAMA/LIBRA set-up:

DAMA/LIBRA surface ≈0.13 m² µ flux @ DAMA/LIBRA ≈2.5 µ/day

It cannot mimic the signature: already excluded by R₉₀, by *multi-hits* analysis + different phase, etc.

Rate, R_n , of fast neutrons produced by μ :

- Φ_{μ} @ LNGS $\approx 20 \,\mu \, \text{m}^{-2} \text{d}^{-1}$ (±1.5% modulated)
- Annual modulation amplitude at low energy due to μ modulation:

$$S_m^{(m)} = R_n g \epsilon f_{\Delta E} f_{single} 2\% / (M_{setup} \Delta E)$$

Moreover, this modulation also induces a variation in other parts of the energy spectrum and in the *multi-hits* events



$S_m^{(\mu)} \le (0.3-2.4) \times 10^{-5} \text{ cpd/kg/keV}$

It cannot mimic the signature: already excluded by R₉₀, by *multi-hits* analysis + different phase, etc.

\checkmark Inconsistency of the phase between DAMA signal and μ modulation

µ flux @ LNGS (MACRO, LVD, BOREXINO) ≈ $3 \cdot 10^{-4}$ m⁻²s⁻¹; modulation amplitude 1.5%; **phase**: July 7 ± 6 d, June 29 ± 6 d (Borexino)

The DAMA phase: May 26 ± 7 days (stable over 13 years)

The DAMA phase is 5.7σ far from the LVD/BOREXINO phases of muons (7.1 σ far from MACRO measured phase)

Considering the seasonal weather al LNGS, quite impossible that the max. temperature of the outer atmosphere (on which μ flux variation is dependent) is observed e.g. in June 15 which is 3 σ from DAMA For many o



For many others arguments EPJC72(2012)2064

- Contributions to the total neutron flux at LNGS;
- Counting rate in DAMA/LIBRA for single-hit events, in the (2-6) keV energy region induced by:

$$\Phi_{k} = \Phi_{0,k} \left(1 + \eta_{k} \cos\omega \left(t - t_{k} \right) \right)$$

$$R_{k} = R_{0,k} \left(1 + \eta_{k} \cos\omega \left(t - t_{k} \right) \right)$$

Modulation

- \succ neutrons,
- \geq muons,

solar neutrinos.

(See e.g. also EPJC 56 (2008) 333, EPJC 72 (2012) 2064, IJMPA 28 (2013) 1330022)

arViv:1100 2514

						ampiltudes			
	Source	$\stackrel{\Phi^{(n)}_{0,k}}{(\mathrm{neutrons\ cm^{-2}\ s^{-1}})}$	η_k	t_k	$egin{array}{c c} R_{0,k} \ (\mathrm{cpd/kg/keV}) \end{array}$		$A_k = R_{0,k} \eta_k \ ({ m cpd/kg/keV})$	A_k/S_m^{exp}	
SLOW	thermal n $(10^{-2} - 10^{-1} \text{ eV})$	$1.08 \times 10^{-6} [15]$	$ \begin{array}{l} \simeq 0 \\ \text{however} \ll 0.1 \ [2, \ 7, \ 8] \end{array} $	_	$< 8 \times 10^{-6}$	[2, 7, 8]	$\ll 8 \times 10^{-7}$	$\ll 7 \times 10^{-5}$	
neutrons	epithermal n (eV-keV)	$2 imes 10^{-6}$ [15]	$ \simeq 0 \\ \text{however} \ll 0.1 \ [2, \ 7, \ 8] $	_	$< 3 \times 10^{-3}$	[2, 7, 8]	$\ll 3 \times 10^{-4}$	≪ 0.03	
	fission, $(\alpha, n) \rightarrow n$ (1-10 MeV)	$\simeq 0.9 \times 10^{-7} \ [17]$	$\simeq 0$ however $\ll 0.1$ [2, 7, 8]	_	$< 6 \times 10^{-4}$	[2, 7, 8]	$\ll 6 \times 10^{-5}$	$\ll 5 \times 10^{-3}$	
FAST neutrons	$\mu \rightarrow n \text{ from rock}$ (> 10 MeV)	$\simeq 3 \times 10^{-9}$ (see text and ref. [12])	0.0129 [23]	end of June [23, 7, 8]	$\ll 7 \times 10^{-4}$	(see text and [2, 7, 8])	$\ll 9 \times 10^{-6}$	$\ll 8 \times 10^{-4}$	
	$\mu \rightarrow$ n from Pb shield (> 10 MeV)	$\simeq 6 \times 10^{-9}$ (see footnote 3)	0.0129 [23]	end of June [23, 7, 8]	$\ll 1.4 \times 10^{-3}$	(see text and footnote 3)	$\ll 2 \times 10^{-5}$	$\ll 1.6 \times 10^{-1}$	
	$ \begin{array}{l} \nu \rightarrow \mathrm{n} \\ (\mathrm{few} \ \mathrm{MeV}) \end{array} $	$\simeq 3 \times 10^{-10}$ (see text)	0.03342 *	Jan. 4th *	$\ll 7 \times 10^{-5}$	(see text)	$\ll 2 \times 10^{-6}$	$\ll 2 \times 10^{-4}$	
	direct μ	$\Phi_0^{(\mu)} \simeq 20 \ \mu \ \mathrm{m}^{-2} \mathrm{d}^{-1} \ [20]$	0.0129 [23]	end of June [23, 7, 8]	$\simeq 10^{-7}$	[2, 7, 8]	$\simeq 10^{-9}$	$\simeq 10^{-7}$	
	direct ν	$\Phi_0^{(\nu)} \simeq 6 \times 10^{10} \ \nu \ \mathrm{cm}^{-2} \mathrm{s}^{-1}$ [26]	0.03342 *	Jan. 4th *	$\simeq 10^{-5}$	[31]	3×10^{-7}	3×10^{-5}	

* The annual modulation of solar neutrino is due to the different Sun-Earth distance along the year; so the relative modulation amplitude is twice the eccentricity of the Earth orbit and the phase is given by the perihelion.

All are negligible w.r.t. the annual modulation amplitude observed by DAMA/LIBRA and they cannot contribute to the observed modulation amplitude.

+ In no case neutrons (of whatever origin) can mimic the DM annual modulation signature since some of the **peculiar requirements of the signature** would fail, such as the neutrons would induce e.g. variations in all the energy spectrum, variation in the multiple hit events,... which were not observed.

Model-independent evidence by DAMA/NaI and DAMA/LIBRA

well compatible with several candidates (in several of the many possible astrophysical, nuclear and particle physics scenarios); other ones are open



Possible model dependent positive hints from Indirect searches (but interpretation, evidence itself, derived mass and cross sections depend e.g. on bckg modeling, on DM spatial velocity distribution in the galactic halo, etc.) not in conflict with DAMA results; null results not in conflict as well

Available results from direct searches using different target materials and approaches do not give any robust conflict & compatibility of possible positive hints

Just few <u>examples</u> of interpretation of the annual modulation in terms of candidate particles in <u>some scenarios</u>



Compatibility with several candidates; other ones are open



...models...

- Which particle?
- Which interaction coupling?
- Which Form Factors for each target-material?
- Which Spin Factor?
- Which nuclear model framework?
- Which scaling law?
- Which halo model, profile and related parameters?
- Streams?
- ..

About interpretation

See e.g.: Riv.N.Cim.26 n.1 (2003) 1, IJMPD13 (2004) 2127, EPJC47 (2006) 263, IJMPA21 (2006) 1445, EPJC56 (2008) 333, PRD84 (2011) 055014, JMPA28 (2013) 1330022

...and experimental aspects...

- Exposures
- Energy threshold
- Detector response (phe/keV)
- Energy scale and energy resolution
- Calibrations
- Stability of all the operating conditions.
- Selections of detectors and of data.
- Subtraction/rejection procedures and stability in time of all the selected windows and related quantities
- Efficiencies
- Definition of fiducial volume and nonuniformity
- Quenching factors, channeling

Uncertainty in experimental parameters, as well as necessary assumptions on various related astrophysical, nuclear and particle-physics aspects, affect all the results at various extent, both in terms of exclusion plots and in terms of allowed regions/volumes. Thus comparisons with a fixed set of assumptions and parameters' values are intrinsically strongly uncertain.

No experiment can be directly compared in model independent way with DAMA

... an example in literature...

Case of DM particles inducing elastic scatterings on target-nuclei





... many other interpretations available in literature

DAMA/LIBRA phase 2 - running

Second upgrade on end of 2010: all PMTs replaced with new ones of higher Q.E.



The second orders effects to be investigated by DAMA/LIBRA-phase2

The importance of studying second order effects and the annual modulation phase



8



New PMTs with higher Q.E.

Conclusions

- Positive evidence for the presence of DM particles in the galactic halo at 9.3σ C.L. (14 annual cycles DAMA/Nal and DAMA/LIBRAphase1: 1.33 ton × yr)
- The modulation parameters determined with better precision
- Full sensitivity to many kinds of DM candidates and interactions types (both inducing recoils and/or e.m. radiation), full sensitivity to low and high mass candidates



DAMA/LIBRA-phase2 in data taking at lower software energy threshold (below 2 keV)

- software energy threshold (below 2 keV) to investigate further features of DM signals and second order effects
- Continuing investigations of rare processes other than DM as well as further developments



Moreover, works and efforts for:

- further improvement (phase3);
- DAMA/1ton set up;
- ADAMO project, anisotropic scintillators for directionality