



Contribution ID: 38

Type: **Poster**

## The $a_0(980)$ revisited

*Thursday, 2 July 2015 13:10 (1h 20m)*

A way to understand the light scalar sector of hadron physics is by applying a type of dynamical generation in which additional resonances arise as companion poles from quark-antiquark seed states by incorporating mesonic loops at the level of S-wave propagators [1–3]. Along this line, we first repeat and complete the calculations of previous works of Tornqvist and Roos (TR) and Boglione and Pennington (BP), where the resonance  $a_0(980)$  appears as an additional pole of the propagator of the predominately quark-antiquark state  $a_0(1450)$ . Both works were important as they demonstrated the feasibility of the idea of companion poles – however, we show that in TR, the same as in BP, the pole width of both states is too large by a factor of 2 when compared to data, and that in BP the pole mass of  $a_0(1450)$  is too large of about 400 MeV. We then construct an effective model Lagrangian for  $a_0(1450)$  coupling to pseudoscalar mesons with both non-derivative and derivative interactions, and calculate its propagator. We demonstrate that it is indeed possible to obtain two poles, one for  $a_0(980)$  and one for  $a_0(1450)$ , in good agreement with data, thus showing that the mechanism of companion poles can deliver not only qualitative but also quantitative correct results.

### References

1. N. A. Tornqvist, Z. Phys. C 68, 647 (1995).
2. N. A. Tornqvist and M. Roos, Phys. Rev. Lett. 76, 1575 (1996).
3. M. Boglione and M. R. Pennington, Phys. Rev. D 65, 114010 (2002).

**Primary author:** WOLKANOWSKI-GANS, Thomas (Goethe-Universität Frankfurt am Main)

**Co-authors:** Prof. RISCHKE, Dirk H. (Institut für Theoretische Physik, Goethe-Universität Frankfurt am Main, Frankfurt am Main); Dr GIACOSA, Francesco (Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität, Frankfurt)

**Presenter:** WOLKANOWSKI-GANS, Thomas (Goethe-Universität Frankfurt am Main)

**Session Classification:** Poster Session

**Track Classification:** Goldstone Boson Working Group