



Contribution ID: 71

Type: **Talk**

## Resonances in the electroweak chiral lagrangian

*Thursday, 2 July 2015 15:00 (20 minutes)*

We study strongly coupled models of electroweak symmetry breaking with a light Higgs boson. We use a resonance effective Lagrangian with bosonic massive resonances together with the Standard Model degrees of freedom, including a light Higgs. We consider constraints from the phenomenology and from the assumed high-energy behavior of the underlying theory. This resonance effective theory can be used to estimate the low-energy constants (LECs) of the Electroweak Effective Theory (Electroweak Chiral Lagrangian) in terms of resonance parameters and to make predictions of low-energy observables like, for instance, the oblique parameters. Note that the theoretical framework is completely analogous to the Resonance Chiral Theory description of QCD at GeV energies.

**Primary author:** ROSELL, Ignasi (Universidad CEU Cardenal Herrera, Valencia)

**Co-authors:** Prof. PICH, Antonio (Valencia University, IFIC); SANTOS, Joaquin (Universitat de Valencia, IFIC); Dr SANZ-CILLERO, Juan Jose (Universidad Aut3noma de Madrid)

**Presenter:** ROSELL, Ignasi (Universidad CEU Cardenal Herrera, Valencia)

**Session Classification:** Parallel Session 5 - Goldstone Boson WG

**Track Classification:** Goldstone Boson Working Group