



Contribution ID: 20

Type: **Poster**

Two-loop cusp anomaly in ABJM at strong coupling

Monday, 15 September 2014 14:10 (20 minutes)

We compute the null cusp anomalous dimension of ABJM theory at strong coupling up to two-loop order. This is done by evaluating corrections to the corresponding superstring partition function, weighted by the AdS₄×CP³ action in AdS light-cone gauge. We compare our result, where we use an anomalous shift in the AdS₄ radius, with the cusp anomaly of N=4 SYM, and extract the two-loop contribution to the non-trivial integrable coupling $h(\lambda)$ of ABJM theory. It coincides with the strong coupling expansion of the exact expression for $h(\lambda)$ recently conjectured by Gromov and Sizov. Our work provides thus a non-trivial perturbative check for the latter, as well as evidence for two-loop UV-finiteness and quantum integrability of the Type IIA AdS₄×CP³ superstring in this gauge. Based on arXiv:1407.4788.

Primary author: Mr VESCOVI, Edoardo (Humboldt University Berlin)

Co-authors: Mr BRÈS, Alexis (École Normale Supérieure); Mr BIANCHI, Lorenzo (Humboldt University Berlin); Dr BIANCHI, Marco S. (Humboldt University Berlin); Dr FORINI, Valentina (Humboldt University Berlin)

Presenter: Mr VESCOVI, Edoardo (Humboldt University Berlin)

Session Classification: Discussion - Posters