

9th Bologna Workshop on CFT and Integrable Models



Bologna Workshop on:

CFT AND INTEGRABLE MODELS

and their applications to Quantum Field Theory, Statistical Mechanics and Condensed Matter Physics



Report of Contributions

Contribution ID: 9

Type: **Plenary Talk**

Non-Conformal Holography and Localization

Monday, September 15, 2014 11:30 AM (1 hour)

Primary author: ZAREMBO, Konstantin (Nordita)

Presenter: ZAREMBO, Konstantin (Nordita)

Session Classification: Monday morning

Contribution ID: 10

Type: **Talk**

Large-N asymptotic expansion of multiple integrals related to the quantum separation of variables method

Monday, September 15, 2014 4:30 PM (30 minutes)

The scalar products and certain correlation functions of models solvable by the quantum separation of variables

can be expressed in terms of N -fold multiple integrals which can be thought of as the partition function of a one dimensional

gas of particles trapped in an external potential V and interacting through repulsive two-body interactions of the type

$$\ln \left[\sinh[\pi \operatorname{om}_1(la - \mu)] \cdot \sinh[\pi \operatorname{om}_2(la - \mu)] \right].$$

The analysis of the large- N asymptotic behaviour of these integrals is of interest to the description of the continuum limit of the integrable model.

Although such partition functions present certain structural resemblances with those arising in the context of the so-called be -ensembles,

their large- N asymptotic analysis demands the introduction of several new ingredients. Such a complication in the analysis

is due to the lack of dilation invariance of the exponential of the two-body interaction.

In this talk, I shall discuss the main features of the method of asymptotic analysis which we have developed.

The method utilises large-deviation techniques on the one hand and the Riemann–Hilbert problem approach to

truncated Wiener-Hopf singular-integral equations on the other hand.

This is a joint work with G. Borot and A. Guionnet.

Primary author: Dr KOZLOWSKI, Karol (CNRS)

Presenter: Dr KOZLOWSKI, Karol (CNRS)

Session Classification: Monday afternoon

Contribution ID: 11

Type: **Plenary Talk**

The Quantum Spectral Curve of the ABJM theory

Tuesday, September 16, 2014 11:00 AM (1 hour)

The discovery of integrability in the planar limit of AdS/CFT has triggered a renewed interest toward the study of finite-size corrections of exactly solvable systems. In this talk, I will present a very recent idea which has made possible the reduction of the complicated Thermodynamic Bethe Ansatz equations for the anomalous dimensions of composite operators in AdS_5/CFT_4 and in AdS_4/CFT_3 to much simpler nonlinear matrix Riemann-Hilbert problems.

Primary author: TATEO, Roberto (TO)**Presenter:** TATEO, Roberto (TO)**Session Classification:** Tuesday morning

Contribution ID: 14

Type: **Plenary Talk**

Correlation functions in Minimal Liouville Gravity from Douglas string equation

Monday, September 15, 2014 10:00 AM (1 hour)

I am going to present some results about (p,q) Minimal Liouville gravity.

It is assumed that the generating function of the correlators in genus zero in Minimal Liouville gravity (MLG) is nothing but logarithm of the Sato tau-function for dispersionless Gelfand-Dikii hierarchy with the special initial condition given by Douglas string equation.

Using the connection between MLG and Frobenius manifold structure we get an explicit and useful expression for log Sato tau-function corresponding Douglas string equation in dispersionless limit.

We argue that the MLG correlators are the expansion coefficients of Log tau-function in respect to the new variables connected with KdV variables by a special nonlinear “resonance” transformation in such a way that the MLG correlators satisfy to the necessary fusion rules.

Primary author: Prof. BELAVIN, Alexander (Landau Institute for Theoretical Physics)

Presenter: Prof. BELAVIN, Alexander (Landau Institute for Theoretical Physics)

Session Classification: Monday morning

Contribution ID: 15

Type: **Plenary Talk**

Entanglement Entropy of non-Unitary Conformal Field Theory

Wednesday, September 17, 2014 9:00 AM (1 hour)

In this talk I will show that the entanglement entropy of a region of large size ℓ in a one-dimensional non-unitary critical model behaves as $S \sim (c_{\text{eff}}/3) \log \ell$, where $c_{\text{eff}} = c - 24\Delta > 0$ is the effective central charge, c (which may be negative) is the central charge of the conformal field theory and $\Delta \leq 0$ is the lowest holomorphic conformal dimension in the theory. This result generalizes the well known expressions for unitary models. I will provide a general proof, as well as numerical evidence for a non-unitary spin chain (an analytical computation using the corner transfer matrix method for a non-unitary lattice model will be discussed by Davide Bianchini). I will show how a new algebraic technique can be used for studying the branching that arise within the replica approach, and find a new expression for the entanglement entropy in terms of correlation functions of twist fields that is valid for non-unitary models. This expression will be further generalized in Olalla Castro-Alvaredo's talk to the massive Yang-Lee model of integrable quantum field theory.

Primary author: Dr DOYON, Benjamin (King's College London)

Presenter: Dr DOYON, Benjamin (King's College London)

Session Classification: Wednesday morning

Contribution ID: 16

Type: **Plenary Talk**

Energy flow correlations in CFT

Monday, September 15, 2014 2:30 PM (1 hour)

We present a new approach to computing energy flow correlations in a generic CFT. These infrared finite observables are familiar from collider physics studies and describe the angular distribution of energy in outgoing radiation created from the vacuum by some source. The energy correlations can be expressed in terms of Wightman correlation functions in a certain limit. We explain how to compute these quantities starting from their Euclidean analogues by means of a non-trivial analytic continuation which, in the framework of CFT, can elegantly be performed in Mellin space. We illustrate the general formalism in N=4 SYM, making use of the well-known results on the four-point correlation function of half-BPS scalar operators.

Primary author: KORCHEMSKY, Gregory (Institut de Physique Theorique, Saclay)

Presenter: KORCHEMSKY, Gregory (Institut de Physique Theorique, Saclay)

Session Classification: Monday afternoon

Contribution ID: 17

Type: **Plenary Talk**

Discretely holomorphic observables in the chiral Potts model

Wednesday, September 17, 2014 2:30 PM (1 hour)

Primary author: Dr WESTON, Robert (Heriot-Watt University)

Presenter: Dr WESTON, Robert (Heriot-Watt University)

Session Classification: Wednesday afternoon

Contribution ID: 19

Type: **Talk**

"Off-critical interfaces in two dimensions. Exact results from field theory"

Thursday, September 18, 2014 10:00 AM (30 minutes)

We will consider interfaces of systems of classical statistical mechanics in two dimensions. Interfaces at criticality are conformally invariant random curves and SLE together with field theory have been successfully applied to several models. On the other hand, below criticality exact results are available only from lattice calculations which are restricted to the Ising model, a circumstance that raises the question about the role of Ising solvability. Our aim is to fill this gap, presenting a new, exact, field-theoretic description of interfaces in two dimension which works directly in the continuum. Finally this framework allow us to clarify the role of integrability.

In particular we show how field theory yields the exact description of intermediate phases in the scaling limit close to a first order phase transition point. The ability of a third phase to form an intermediate wetting layer or only isolated bubbles is explicitly related to the spectrum of excitations of the field theory. The order parameter profiles are determined and interface properties such as passage probabilities and internal structure are deduced from them. The theory is illustrated through the application to the q -state Potts model and Ashkin-Teller model. The latter is shown to provide the first exact solution of a bulk wetting transition. The whole technique can be applied also to the study of interfaces at boundaries. We will develop the exact theory of phase separation in a two-dimensional wedge from the properties of the order parameter and boundary condition changing operators in field theory. For a shallow wedge we determine the passage probability for an interface with endpoints on the boundary. For generic opening angles we exhibit the fundamental origin of the filling transition condition and of the property known as wedge covariance. The limit of a straight opening angle corresponds to an interface on the half plane, the midpoint average distance of the interface from the boundary grows as the square root of the distance between the endpoints, unless the reflection amplitude of the bulk excitations on the boundary possesses a stable bound state pole. The contact angle of the phenomenological wetting theory is exactly related to the location of this pole. Results available from the lattice solution of the Ising model are recovered as a particular case.

References

- [1] G. Delfino and A.S., Interfaces and wetting transition on the half plane. Exact results from field theory, *J. Stat. Mech.* P05010 (2013)
- [2] G. Delfino and A.S., Exact theory of intermediate phases in two dimensions, *Annals of Physics* 342, 171 (2014)
- [3] G. Delfino and A.S., Phase separation in a wedge. Exact results, [arXiv1403.1138], to appear in *Physical Review Letters*

Primary author: SQUARCINI, Alessio (TS)

Co-author: DELFINO, Gesualdo (TS)

Presenter: SQUARCINI, Alessio (TS)

Session Classification: Thursday morning

Contribution ID: 20

Type: **Poster**

Two-loop cusp anomaly in ABJM at strong coupling

Monday, September 15, 2014 2:10 PM (20 minutes)

We compute the null cusp anomalous dimension of ABJM theory at strong coupling up to two-loop order. This is done by evaluating corrections to the corresponding superstring partition function, weighted by the AdS₄×CP³ action in AdS light-cone gauge. We compare our result, where we use an anomalous shift in the AdS₄ radius, with the cusp anomaly of N=4 SYM, and extract the two-loop contribution to the non-trivial integrable coupling $h(\lambda)$ of ABJM theory. It coincides with the strong coupling expansion of the exact expression for $h(\lambda)$ recently conjectured by Gromov and Sizov. Our work provides thus a non-trivial perturbative check for the latter, as well as evidence for two-loop UV-finiteness and quantum integrability of the Type IIA AdS₄×CP³ superstring in this gauge. Based on arXiv:1407.4788.

Primary author: Mr VESCOVI, Edoardo (Humboldt University Berlin)

Co-authors: Mr BRÈS, Alexis (École Normale Supérieure); Mr BIANCHI, Lorenzo (Humboldt University Berlin); Dr BIANCHI, Marco S. (Humboldt University Berlin); Dr FORINI, Valentina (Humboldt University Berlin)

Presenter: Mr VESCOVI, Edoardo (Humboldt University Berlin)

Session Classification: Discussion - Posters

Contribution ID: 21

Type: **Talk**

Form factor approach to correlation functions in massless quantum integrable models

Tuesday, September 16, 2014 12:00 PM (30 minutes)

We study the asymptotic behaviour of the dynamical correlation functions of massless quantum integrable models. We develop an approach based on the summation of the appropriate (critical) form factors. We show that this approach gives rise to a number of new hypergeometric identities. These identities lead to the leading terms of the asymptotics for the two-point and multi-point correlation functions as well as to the analysis of the edge behaviour of the dynamical structure factors.

Primary author: Prof. KITANINE, Nikolai (Université de Bourgogne)

Presenter: Prof. KITANINE, Nikolai (Université de Bourgogne)

Session Classification: Tuesday morning

Contribution ID: 27

Type: **Poster**

Junctions of one-dimensional hard-core bosons

Monday, September 15, 2014 1:50 PM (20 minutes)

We map a system of one-dimensional guides containing bosons with strongly repulsive interaction onto the topological Kondo model. We study the thermodynamic properties by computing the ground state energy and the temperature dependence of the free energy shift due to the junction.

Primary author: Dr BUCCHERI, Francesco (IIP-UFRN)

Presenter: Dr BUCCHERI, Francesco (IIP-UFRN)

Session Classification: Discussion - Posters

Contribution ID: 28

Type: **Talk**

Correlations after quantum quenches in the XXZ spin chain: Failure of the Generalized Gibbs Ensemble

Tuesday, September 16, 2014 4:30 PM (30 minutes)

In closed integrable systems, thermalization after a quantum quench is precluded by the existence of infinitely many conserved charges. The idea of the generalized Gibbs ensemble (GGE) is to include all local conserved charges into the canonical density operator with appropriate Lagrange-multipliers to ensure conservation of local charges. It is a nontrivial question whether the GGE correctly describes all local observables in post-quench steady states. While the validity of the GGE among free theories has been established, we showed that it breaks down for certain quenches of the spin-1/2 XXZ Heisenberg spin chain.

We studied local correlations in the steady state of the XXZ chain after quenches starting from the Majumdar–Ghosh dimer product and Néel states. We computed the correlations using a quench action approach (QA) variant of the thermodynamic Bethe ansatz (TBA) and independently using an infinite time-evolving block decimation (iTEBD) algorithm which simulated the real time evolution of correlations. While the steady-state correlations obtained through the QA and the iTEBD methods agreed consistently, predictions of the GGE failed to reproduce these values. Therefore the GGE, in its present form, is not a generally valid statistical description of post-quench steady states of closed integrable systems.

References:

- [1] B. Pozsgay, M. Mestyán, M. A. Werner, M. Kormos, G. Zaránd and G. Takács. Phys. Rev. Lett., in press, arXiv:1405.2843 (2014)
- [2] M. Mestyán and B. Pozsgay. J. Stat. Mech.: Theor. Exp., in press, arXiv:1405.0232 (2014)
- [3] B. Pozsgay. arXiv:1406.4613 (2014)

Primary author: Mr MESTYÁN, Márton (Budapest University of Technology and Economics)

Presenter: Mr MESTYÁN, Márton (Budapest University of Technology and Economics)

Session Classification: Tuesday afternoon

Contribution ID: 29

Type: **Talk**

Excited state entanglement in conformal field theory: extensivity and the role of microscopic detail

Wednesday, September 17, 2014 4:30 PM (30 minutes)

We study the excited state subsystem purity (equivalent to 2nd Renyi entropy) as a measure of entanglement in 1+1d conformal field theories. In order to access the highly excited states we generalize the replica approach and develop a systematic framework to treat arbitrary states (in particular descendants) by means of novel CFT techniques including transformation of descendant fields under conformal mappings and exact evaluation of descendant n-point functions. By examining a number of individual states in the critical Ising and three-state Potts models we uncover a new interesting regime of the subsystem purity as a function of the relative subsystem size: when the subsystems are comparable purity decays exponentially with an exponent depending only on the excitation energy and in a nonlinear way. For the Renyi entropy this translates as a quasi thermodynamic entropy, that is extensive but obeys a nonlinear 1st law analog. As an additional application of the present framework we show preliminary results for ground and excited state Renyi entropies in the massive Ising model using a truncated conformal space approach.

arXiv:1406.3182 [hep-th]

Primary author: PALMAI, Tamas (SISSA, INFN TS)

Presenter: PALMAI, Tamas (SISSA, INFN TS)

Session Classification: Wednesday afternoon

Contribution ID: 31

Type: **Talk**

Recurrence Relation for Conformal Blocks

Wednesday, September 17, 2014 4:00 PM (30 minutes)

We generalize the idea of 1307.6856 to build conformal blocks of a generic CFT (in dimension $d > 2$) using a recurrence relation, idea which has roots in a paper by Zamolodchikov (1984) originally applied to Virasoro blocks in 2 dimensions.

Generalizing this construction we are able to find a recurrence relation to build conformal blocks associated to the four point function of vector operators.

Primary author: Mr TREVISANI, Emilio (Porto University)

Presenter: Mr TREVISANI, Emilio (Porto University)

Session Classification: Wednesday afternoon

Contribution ID: 32

Type: **Talk**

Q-functions and classical affine Toda Field Theories

Tuesday, September 16, 2014 10:00 AM (30 minutes)

We discuss a connection between classical (1+1)-dimensional A_{n+1} affine Toda field theories (ATFTs) and Bethe Ansatz systems. We show that the connection coefficients for specific solutions of the associated linear problem for the classical ATFTs satisfy certain functional relations. This way we enlarge the so-called ODE/IM correspondence.

Based on: P. Adamopoulou, C. Dunning 2014 Bethe Ansatz equations for the classical A_{n+1} affine Toda field theories, J. Phys. A: Math. Theor. 47 205205

Primary author: Dr ADAMOPOULOU, Panagiota (University of Kent)

Co-author: Dr DUNNING, Clare (University of Kent)

Presenter: Dr ADAMOPOULOU, Panagiota (University of Kent)

Session Classification: Tuesday morning

Contribution ID: 33

Type: **Talk**

Revealing elementary excitations of one-dimensional Bose gases through their dynamical structure factor

Tuesday, September 16, 2014 4:00 PM (30 minutes)

Interactions are known to have dramatic effects on bosonic gases in low dimension: Not only does the ground state transform from a condensate-like state to an effective Fermi sea, but new fundamental excitations are predicted to appear which do not have any higher-dimensional equivalents. In this work, we trace these elusive excitations via their effects on the dynamical structure factor of arrays of one-dimensional strongly-interacting Bose gases at low temperature, probed by energy deposition through low-momentum Bragg excitations. The experimental signals are compared to recent theoretical predictions for the dynamical structure factor of the Lieb-Liniger model at finite temperatures. Our results demonstrate that the main contribution to the spectral widths can be explained from the dynamics of the interaction-induced excitations in the gas.

Primary author: PANFIL, Milosz

Presenter: PANFIL, Milosz

Session Classification: Tuesday afternoon

Contribution ID: 34

Type: **Talk**

Universal dynamics of a soliton after a quantum quench

Monday, September 15, 2014 4:00 PM (30 minutes)

We propose a new type of experimentally feasible quantum quench protocol. A quantum system is prepared in a coherent, localized excited state of a Hamiltonian, whose interaction strength is characterized by a control parameter. At some point during the evolution of this solitonic excitation, the parameter is suddenly changed. We study the dynamics of solitons after this global quench in a semi-classical, hydrodynamic, limit, for a wide class of systems, and we find it to be universal at short times, i.e. not depending on the microscopics of the physical system.

Numerical support for these results is presented using the Calogero model and the non-linear Schroedinger equation, relevant for the implementation of the proposed protocol with ultracold bosons. Finally, it is shown that the effects of integrability breaking by a parabolic potential and by a power-law non-linearity do not change the universality of the short-time dynamics.

Primary author: FRANCHINI, Fabio (TS)

Presenter: FRANCHINI, Fabio (TS)

Session Classification: Monday afternoon

Contribution ID: 35

Type: **Plenary Talk**

Breaking the area law in deformed critical spin chains

Thursday, September 18, 2014 9:00 AM (1 hour)

Primary author: Prof. SIERRA, German (Instituto de Fisica Teorica UAM-CSIC)

Presenter: Prof. SIERRA, German (Instituto de Fisica Teorica UAM-CSIC)

Session Classification: Thursday morning

Contribution ID: 36

Type: **Plenary Talk**

QCD Pomeron and AdS/CFT Integrability

Tuesday, September 16, 2014 9:00 AM (1 hour)

Presenter: Dr GROMOV, Nikolay (KCL)

Session Classification: Tuesday morning

Contribution ID: 37

Type: **Plenary Talk**

Quantum Quench Dynamics

Tuesday, September 16, 2014 2:30 PM (1 hour)

I will discuss in simple terms some topics emerging from recent studies of quantum systems driven away from equilibrium. Particular emphasis will be given to the integrable/non-integrable nature of the Hamiltonian and to its local properties.

Presenter: MUSSARDO, Giuseppe (TS)

Session Classification: Tuesday afternoon

Contribution ID: 38

Type: **Plenary Talk**

Logarithmic Minimal Models: Logging Progress

Thursday, September 18, 2014 11:00 AM (1 hour)

Presenter: Prof. PEARCE, Paul (University of Melbourne)

Session Classification: Thursday morning

Contribution ID: 39

Type: **Plenary Talk**

Constraints on CFTs in Diverse Dimensions from the Bootstrap Mechanism

Wednesday, September 17, 2014 11:30 AM (1 hour)

Recently an efficient numerical method has been developed to implement crossing symmetry and unitarity on the operator dimensions and on OPE coefficients in CFTs in various dimensions. These calculations can be done only for theories lying at the boundary of the allowed parameter space. I illustrate an alternative method that can be applied to a larger class of CFTs, whether unitary or not, though the obtained results are not yet as precise as in the usual method. Examples of this kind of calculations, where the usual method cannot be applied, include the CFTs associated with the Yang-Lee edge singularity in $D < 6$ or with the ordinary surface transition in 3D Ising model.

Primary author: GLIOZZI, Ferdinando (TO)

Presenter: GLIOZZI, Ferdinando (TO)

Session Classification: Wednesday morning

Contribution ID: 40

Type: **Talk**

Entanglement Entropy of the Lee-Yang Model from Branch Point Twist Fields

Wednesday, September 17, 2014 11:00 AM (30 minutes)

In this talk I will present new (unpublished) results regarding the entanglement entropy of one of the simplest integrable models: the Lee-Yang theory. Despite its simplicity, the Lee-Yang theory is non-unitary. One of the most famous consequences of this non-unitarity was the realization by Zamolodchikov that the form factor expansion of correlation functions is now given in terms of an alternating series. In my talk I will explain how form factors of twist fields and their correlators may be computed for this model which enter an expression for the entanglement entropy of the theory. I will show how this form factor expansion reproduces the predicted CFT behaviour for short distances and how next-to-leading order corrections to the entanglement entropy of large sub-systems differ from those found for unitary models.

Primary author: Dr CASTRO ALVAREDO, Olalla (City University London)

Presenter: Dr CASTRO ALVAREDO, Olalla (City University London)

Session Classification: Wednesday morning

Contribution ID: 41

Type: **Talk**

Entanglement Entropy of Forrester-Baxter RSOS Models

Wednesday, September 17, 2014 10:00 AM (30 minutes)

We analytically evaluate the Rényi entropy for the Forrester-Baxter RSOS models in the regime III. These models are lattice realisations of non-unitary minimal models of CFT. By studying the leading term of this expression it is possible to probe conformal properties of the ground state. Our results agree completely with recent CFT predictions for the entanglement entropy of non-unitary systems. Furthermore we discuss a class of possible quantum Hamiltonian related to off-critical RSOS models.

Primary author: Mr BIANCHINI, Davide (City University London)

Presenter: Mr BIANCHINI, Davide (City University London)

Session Classification: Wednesday morning

Contribution ID: 42

Type: **Poster**

On the scattering over the GKP Vacuum

Monday, September 15, 2014 1:30 PM (20 minutes)

Within the framework of the AdS₅/CFT₄ correspondence, the GKP string living on a AdS₅ × S⁵ background finds a counterpart in the antiferromagnetic vacuum state for the spin chain, fruitfully employed to investigate the dual N = 4 SYM superconformal gauge theory. Integrability underlying such a model allows us to analyse the excitations over the vacuum considered, by computing their scattering matrices and dispersion relations.

Primary author: PISCAGLIA, Simone

Presenter: PISCAGLIA, Simone

Session Classification: Discussion - Posters