



Contribution ID: 204

Type: poster

Plasma production for electron acceleration by resonant plasma wave.

Monday, 14 September 2015 20:00 (30 minutes)

Plasma wakefield acceleration is the most promising acceleration technique known nowadays, able to provide very high accelerating fields ($10\text{--}100\text{ GVm}^{-1}$), enabling acceleration of electrons to GeV energy in few centimetres. However, the quality of the electron bunches accelerated with this technique is still not comparable with that of conventional accelerators; Radiofrequency-based accelerators, in fact, are limited in accelerating field ($10\text{--}100\text{ MVm}^{-1}$) requiring therefore kilometric distances to reach the GeV energies, but can provide very bright electron bunches. To combine high brightness electron bunches from conventional accelerators and high accelerating fields reachable with plasmas could be a good compromise allowing to further accelerate high brightness electron bunches coming from LINAC while preserving electron beam quality. Following the idea of plasma wave resonant excitation driven by a train of short bunches, we have started to study the requirements in terms of plasma for SPARC_LAB. In particular here we focus on hydrogen plasma discharge, both from a theoretical and an experimental point of view.

Primary author: ANANIA, Maria Pia (LNF)

Co-authors: CIANCHI, Alessandro (ROMA2); BIAGIONI, Angelo (LNF); ZIGLER, Arie (LNF); DI GIOVENALE, Domenico (LNF); CHIADRONI, Enrica (LNF); FILIPPI, Francesco (ROMA1); FERRARIO, Massimo (LNF)

Presenter: ANANIA, Maria Pia (LNF)

Session Classification: Poster Session 1 (WG1-WG2-WG3-WG4) and Wine

Track Classification: WG4 - Application of compact and high-gradient accelerators/Advanced beam manipulation and control