



Contribution ID: 118

Type: talk

Wakefield-Induced Ionization Injection and Self-Similar Staging in Beam-driven Plasma Wakes.

Monday, September 14, 2015 6:00 PM (30 minutes)

We propose a simple strategy for controlled ionization-induced trapping of electrons in beam-driven plasma accelerators (PWFA) [1]. The presented method exploits the electric wakefields to ionize electrons from a dopant gas and to capture them into a well-defined volume of the accelerating and focusing wake phase, leading to the formation of high-quality witness bunches. This injection principle is explained by example of three-dimensional particle-in-cell simulations using the code OSIRIS. The produced bunches feature multi-kA peak currents, $\sim 1 \mu\text{m}$ transverse normalized emittances, uncorrelated energy spreads of $< 1\%$ on a GeV-energy scale, and femtosecond bunch lengths. The hereby generated witness bunches fulfill all the requirements to drive the same injection principle in a largely increased plasma density. This brings up a new concept of staging that has the potential of producing electron beams with unprecedented energy and quality in PWFA.

References

[1] A. Martinez de la Ossa et al., Phys. Rev. Lett. 111, 245003 (2013).

Primary author: Dr MARTINEZ DE LA OSSA, Alberto (DESY)

Co-authors: Dr OSTERHOFF, Jens (Deutsches Elektronen-Synchrotron DESY); Dr SCHAPER, Lucas (University Hamburg / DESY); Dr STREETER, Matthew (DESY); Dr MEHRLING, Timon (DESY)

Presenter: Dr MARTINEZ DE LA OSSA, Alberto (DESY)

Session Classification: WG1 - Electron beams from plasmas

Track Classification: WG1 - Electron beams from plasmas