



Contribution ID: 147

Type: **talk**

## **A collinear wakefield accelerator for a high repetition rate multi beamline soft x-ray free-electron laser facility**

*Monday, 14 September 2015 16:00 (15 minutes)*

A concept is presented for a multi beamline soft x-ray FEL facility where several free-electron laser (FEL) undulator lines are driven by equal number of high repetition rate single-stage collinear wakefield accelerators (CWA). Various means to mitigate the adverse collective effects are considered and a practical design of the CWA, embedded into a quadrupole wiggler to control a beam breakup instability, and extending over 30 meters is presented. It is shown that the maximum achievable strength of the quadrupole gradient defines a threshold charge in the drive beam and a maximum attainable acceleration gradient for the witness beam. It is also pointed out that the distance between the drive and witness bunches varies along the accelerator with a measurable impact on the energy gain of the witness bunch and its energy spread. Means to mitigate these effects are also considered. Obtaining the witness bunch with a minimum energy spread is discussed. Finally, results are presented for the expected FEL performance using an appropriately tapered undulator.

**Primary author:** Dr ZHOLENTS, Alexander (Argonne National Laboratory)

**Presenter:** Dr ZHOLENTS, Alexander (Argonne National Laboratory)

**Session Classification:** WG3 - Electron beams from electromagnetic structures, including dielectric and laser-driven structures

**Track Classification:** WG3 - Electron beams from electromagnetic structures, including dielectric and laser-driven structures