



Contribution ID: 78

Type: **talk**

## **Ion acceleration from laser driven collisionless shockwaves in optically shaped gas targets**

*Wednesday, 16 September 2015 17:10 (20 minutes)*

We report on recent experiments using the high power CO<sub>2</sub> laser ( $\lambda \sim 10 \mu\text{m}$ ,  $a_0 \sim 1$ ) at the Accelerator Test Facility, Brookhaven National Laboratory to generate  $\sim\text{MeV}$  protons and helium ions, which are accelerated by collisionless shockwave acceleration in overcritical gas targets. We have developed new techniques of shaping gas jet targets with optically induced hydrodynamic shocks, allowing us to tailor density profiles to optimise ion generation. Sufficiently steep gradients resulted in quasi-monoenergetic proton beams with a single laser pulse, in contrast to previous work relying on pulse trains. The laser plasma interaction also generates energetic electrons, which propagate into the target with beam to background ratio up to  $\sim 0.1$ . We have observed transverse filamentation in this region, which is reproduced by PIC modelling. These simulations also elucidate the key role of the laser-generated electrons in heating the plasma to create the conditions required for collisionless shockwave formation and subsequent ion generation.

**Primary author:** Dr DOVER, Nicholas (Imperial College London)

**Co-authors:** Dr MAHARJAN, Chakra (Stony Brook University); Dr POGORELSKY, Igor (Brookhaven National Laboratory); Dr POLYANSKIY, Mikhail (Brookhaven National Laboratory); Dr COOK, Nathan (Stony Brook University); Dr TRESKA, Olivier (Brookhaven National Laboratory); Dr SHKOLNIKOV, Peter (Stony Brook University); Dr KINGHAM, Robert (Imperial College London); Prof. NAJMUDIN, Zufikar (Imperial College London)

**Presenter:** Dr DOVER, Nicholas (Imperial College London)

**Session Classification:** WG2 - Ion beams from plasmas

**Track Classification:** WG2 - Ion beams from plasmas