

# Studio e investigazione di DM tramite una marcatura model-independent

## *Past:*



- Annual Modulation: DAMA/NaI and DAMA/LIBRA-phase1 results

## *Present:*

- Annual Modulation: DAMA/LIBRA-phase2

## *Future (WN):*

- Annual Modulation
- Studies at second order (ex: phase vs energy)
- Diurnal Modulation
- ...



- Directionality: see talk of Riccardo Cerulli

# Prospects

**a possible phase-3 at end of phase2  
(or data taking dedicated to other rare processes)**

The strong interest in the low energy range suggests the possibility of a new development of high Q.E. PMTs with increased radiopurity to directly couple them to the DAMA/LIBRA crystals, removing the special quartz light guides which act also as optical window obtaining an ultimate number of ph.e./keV.  
(many rare processes can be investigated as discussed in CSN2 many times in the past)

**a possible multipurpose fully sensitive DAMA/1ton**

Proposed by DAMA since 1996  
(DAMA/NaI and DAMA/LIBRA intermediate steps, some R&D funded and carried out)

**New anisotropic scintillator/nanotube detectors for directionality**

**at the end of LNGS DAMA/LIBRA underground data taking:  
New measurements of q.f., channelling, etc. for each detector @ Tor  
Vergata and neutron beam**

**MOREOVER:** with our international partners developments and use of many low background/new/enriched scintillators/samples to deeply investigate rare processes with the realization of specific high mass set-ups


# The annual modulation: a model independent signature for the investigation of DM particles component in the galactic halo

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions can point out its presence.

Drukier, Freese, Spergel PRD86; Freese et al. PRD88

## Requirements of the annual modulation

- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about 2 June)
- 5) Just for single hit events in a multi-detector set-up
- 6) With modulation amplitude in the region of maximal sensitivity must be <7% for usually adopted halo distributions, but it can be larger in case of some possible scenarios



- $v_{\text{sun}} \sim 232 \text{ km/s}$  (Sun vel in the halo)
- $v_{\text{orb}} = 30 \text{ km/s}$  (Earth vel around the Sun)
- $\gamma = \pi/3, \omega = 2\pi/T, T = 1 \text{ year}$
- $t_0 = 2^{\text{nd}} \text{ June}$  (when  $v_{\oplus}$  is maximum)


$$v_{\oplus}(t) = v_{\text{sun}} + v_{\text{orb}} \cos\gamma \cos[\omega(t-t_0)]$$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

the DM annual modulation signature has a different origin and peculiarities (e.g. the phase) than those effects correlated with the seasons

To mimic this signature, spurious effects and side reactions must not only - obviously - be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements

# DAMA/NaI & DAMA/LIBRA experiments main upgrades and improvements



PHASE2

Minimal upgrade in Fall

July 2000 new DAQ and new electronic chain installed (MULTIPLEXER removed, now one TD channel for each detector):  
(i) TD VXI Tektronix;  
(ii) Digital Unix DAQ system;  
(iii) GPIB-CAMAC.

July 2002 DAMA/NaI data taking completed


On 2003 DAMA/LIBRA has begun first operations

Sept.-Oct. 2008 – DAMA/LIBRA upgrade:  
① one detector recovered by replacing a broken PMT  
② a new optimization of some PMTs and HVs performed  
③ all the TD replaced with new ones (U1063A Acqiris 8-bit 1GS/s DC270 High-Speed cPCI Digitizers)  
④ a new DAQ with optical read-out installed.

The second DAMA/LIBRA upgrade in Fall 2010:  
Replacement of all the PMTs with higher Q.E. ones from dedicated developments  
(+new preamp in Fall 2012 and other developments in progress)  
**DAMA/LIBRA-phase2 in data taking**


# DAMA/LIBRA phase 2 - running

Second upgrade on end of 2010:  
all PMTs replaced with new ones of higher Q.E.



# DAMA/LIBRA phase 2 - running

## Quantum Efficiency features




## Residual Contamination

The limits are at 90% C.L.

PMT	Time (s)	Mass (kg)	<sup>226</sup> Ra (Bq/kg)	<sup>234m</sup> Pa (Bq/kg)	<sup>235</sup> U (mBq/kg)	<sup>228</sup> Ra (Bq/kg)	<sup>228</sup> Th (mBq/kg)	<sup>40</sup> K (Bq/kg)	<sup>137</sup> Cs (mBq/kg)	<sup>60</sup> Co (mBq/kg)
Average			0.43	-	47	0.12	83	0.54	-	-
Standard deviation			0.06	-	10	0.02	17	0.16	-	-

## Energy resolution



$\sigma/E$  @ 59.5 keV for each detector with new PMTs with higher quantum efficiency (blue points) and with previous PMT EMI-Electron Tube (red points).

## The light responses



Previous PMTs: 5.5-7.5 ph.e./keV  
New PMTs: up to 10 ph.e./keV

- To study the nature of the particles and features of related astrophysical, nuclear and particle physics aspects, and to investigate second order effects
- Special data taking for *other rare processes*

# Examples of low energy calibrations DAMA/LIBRA-ph2





Routine calibrations each about 10 days in the same running conditions with  $^{241}\text{Am}$  sources



**PRELIMINARY,  
CONFIDENTIAL**

$^{40}\text{K}$  ( $\delta=0.0117\%$ )

EC: (1461 keV  $\gamma$ ) + (3.2 keV X-rays/Auger electrons); b.r.=10.66%




**DAMA/LIBRA is the only set-up calibrated near the software energy threshold**


# Examples of low-energy distribution of the *single-hit* scintillation events in some DAMA/LIBRA-phase2\_2-a.c. detector

Single-hit events = each detector has all the others as anticoincidence

DAMA/LIBRA-phase1



DAMA/LIBRA-phase2-2a.c.





# WN: DAMA/LIBRA-phase3

- The light collection of the detectors can further be improved.
- Thus, the light yields and the energy thresholds will improve too.

The strong interest in the low energy range suggests the possibility of a new development of **high Q.E. PMTs** with **increased radiopurity** to directly couple them to the DAMA/LIBRA crystals, **removing** the special radio-pure quartz (Suprasil B) light guides (10 cm long), which act also as optical window.

The presently-reached PMTs features, **but not for the same PMT mod.:**

- Q.E. around 35-40% @ 420 nm (NaI(Tl) light)
- radiopurity at level of 5 mBq/PMT ( $^{40}\text{K}$ ), 3-4 mBq/PMT ( $^{232}\text{Th}$ ), 3-4 mBq/PMT ( $^{238}\text{U}$ ), 1 mBq/PMT ( $^{226}\text{Ra}$ ), 2 mBq/PMT ( $^{60}\text{Co}$ ).

R&D efforts to obtain PMTs matching the best performances... **feasible**

No longer need for light guides (a 30-40% improvement in the light collection is expected)




# Towards a possible multi-purpose DAMA/1ton



- 1) Proposed since 1996 (DAMA/NaI and DAMA/LIBRA intermediate steps)
- 2) Technology largely at hand (large experiences and fruitful collaborations among INFN and companies/industries)
- 3) Still room for further improvements in the low-background characteristics of the set-up (NaI(Tl) crystals, PMTs, shields, etc.)
- 4) 1 ton detector: the cheapest, the highest duty cycle, the clear signature, fast realization in few years



## A possible design: DAMA/1 ton can be realized by four replicas of DAMA/LIBRA:



- the detectors could be of similar size than those already used
- the features of low-radioactivity of the set-up and of all the used materials would be assured by many years of experience in the field
- electronic chain and controls would profit by the previous experience and by the use of compact devices already developed, tested and used.
- new digitizers will offer high expandibility and high performances
- the daq can be a replica of that of DAMA/LIBRA



- Some R&Ds on PMTs and on crystals carried out
- Other ideas

Electronic chain and example of the trigger system

# Features of the DM signal

The importance of studying **second order effects** and the **annual modulation phase**

High exposure and lower energy threshold can allow further investigation on:

## - the nature of the DM candidates

- ✓ to disentangle among the different astrophysical, nuclear and particle physics models (nature of the candidate, couplings, inelastic interaction, form factors, spin-factors ...)
- ✓ scaling laws and cross sections
- ✓ multi-component DM particles halo?

## - possible diurnal effects on the sidereal time


- ✓ expected in case of high cross section DM candidates (shadow of the Earth)
- ✓ due to the Earth rotation velocity contribution (it holds for a wide range of DM candidates)
- ✓ due to the channeling in case of DM candidates inducing nuclear recoils.

## - astrophysical models


- ✓ velocity and position distribution of DM particles in the galactic halo, possibly due to:
  - satellite galaxies (as Sagittarius and Canis Major Dwarves) tidal "streams";
  - caustics in the halo;
  - gravitational focusing effect of the Sun enhancing the DM flow ("spike" and "skirt");
  - possible structures as clumpiness with small scale size
  - Effects of gravitational focusing of the Sun

The annual modulation phase depends on :



- Presence of streams (as SagDEG and Canis Major) in the Galaxy
- Presence of caustics
- Effects of gravitational focusing of the Sun



The effect of the streams on the phase depends on the galactic halo model




Such investigations allowed by:  
**DAMA/LIBRA-phase3**  
**and DAMA/1ton**  
 with lower energy threshold and larger exposure



# Towards signatures for the presence of streams in the Galactic halo

The effect of the streams on the phase depends on the galactic halo model



DM Phase with no streams  
 $t_0 = 152.5$  d (2<sup>nd</sup> June)

Evans' log axisymmetric non-rotating,  $v_0=220$ km/s,  $R_c=5$ kpc,  $\rho_0$  max + 4% Sgr

NFW spherical isotropic non-rotating,  $v_0=220$ km/s,  $\rho_0$  max + 4% Sgr

The higher sensitivity of DAMA/LIBRA-phase2&3 and DAMA/1ton will allow the further investigation of possible contributions of streams in the galactic halo

If the sensitivity on the phase can be in DAMA/1ton at level of  $\sim 1$  d, discrimination power on the local velocity,  $V_0$ , will be obtained:

- $v_0 = 170$  km/s  $\rightarrow t_0 - t_{\text{equinox}} = 71.7$  d
- $v_0 = 220$  km/s  $\rightarrow t_0 - t_{\text{equinox}} = 73.2$  d
- $v_0 = 270$  km/s  $\rightarrow t_0 - t_{\text{equinox}} = 74.2$  d

Eur. Phys. J. C 73(2013) 2648  
 Eur. Phys. J. C 47(2006) 263

DAMA/NaI & DAMA/LIBRA-ph1  
 DAMA/NaI & DAMA/LIBRA-ph1 & DAMA/LIBRA-ph2&3  
 DAMA/1ton (10 ton × yr)

(2-6) keV:  $t_0 = (144 \pm 7)$  d  
 (0.5/1-6) keV:  $t_0 = (149 \pm 3/4)$  d  
 (0.5/1-6) keV:  $t_0 = (149 \pm 2)$  d

Assuming same mean value

- *Other signatures?*
- *Other second order effects*
- *...*

# A diurnal effect with the sidereal time is expected for DM because of Earth rotation

Eur. Phys. J. C 74 (2014) 2827

Velocity of the detector in the terrestrial laboratory:

$$\vec{v}_{lab}(t) = \vec{v}_{LSR} + \vec{v}_{\odot} + \vec{v}_{rev}(t) + \vec{v}_{rot}(t),$$

Since:

- $|\vec{v}_s| = |\vec{v}_{LSR} + \vec{v}_{\odot}| \approx 232 \pm 50$  km/s,
- $|\vec{v}_{rev}(t)| \approx 30$  km/s
- $|\vec{v}_{rot}(t)| \approx 0.34$  km/s at LNGS

$$v_{lab}(t) \simeq v_s + \hat{v}_s \cdot \vec{v}_{rev}(t) + \hat{v}_s \cdot \vec{v}_{rot}(t).$$

$\vec{v}_{LSR}$  velocity of the Local Standard of Rest (LSR) due to the rotation of the Galaxy

$\vec{v}_{\odot}$  Sun peculiar velocity with respect to LSR

$\vec{v}_{rev}(t)$  velocity of the revolution of the Earth around the Sun

$\vec{v}_{rot}(t)$  velocity of the rotation of the Earth around its axis at the latitude and longitude of the laboratory.

## Annual modulation term:



$$\hat{v}_s \cdot \vec{v}_{rev}(t) = V_{Earth} B_m \cos(\omega(t - t_0))$$

- $V_{Earth}$  is the orbital velocity of the Earth  $\approx 30$  km/s
- $B_m \approx 0.489$
- $t_0 \approx t_{equinox} + 73.25$  days  $\approx$  June 2

## Diurnal modulation term:

$$\hat{v}_s \cdot \vec{v}_{rot}(t) = V_r B_d \cos[\omega_{rot}(t - t_d)]$$

- $V_r$  is the rotational velocity of the Earth at the given latitude (for LNGS  $\approx 0.3435$  km/s)
- $B_d \approx 0.671$
- $t_d \approx 14.02$  h (at LNGS)





Velocity of the Earth in the galactic frame as a function of the sidereal time, with starting point March 21 (around spring equinox). The contribution of diurnal rotation has been dropped off. The maximum of the velocity (vertical line) is about 73 days after the spring equinox.

Sum of the Sun velocity in the galactic frame ( $v_s$ ) and of the rotation velocity of a detector at LNGS ( $\hat{v}_s \cdot \vec{v}_{rot}(t)$ ) as a function of the sidereal time. The maximum of the velocity is about at 14 h (vertical line).

# Model independent result on possible diurnal effect in DAMA/LIBRA-phase1


Eur. Phys. J. C 74 (2014) 2827



- Experimental *single-hit* residuals rate vs either sidereal and solar time and vs energy.
- These residual rates are calculated from the measured rate of the *single-hit* events after subtracting the constant part

Energy region where the annual modulation is observed.

Energy region just above.



Energy	Solar Time	Sidereal Time
2-4 keV	$\chi^2/\text{d.o.f.} = 35.2/24 \rightarrow P = 7\%$	$\chi^2/\text{d.o.f.} = 28.7/24 \rightarrow P = 23\%$
2-5 keV	$\chi^2/\text{d.o.f.} = 35.5/24 \rightarrow P = 6\%$	$\chi^2/\text{d.o.f.} = 24.0/24 \rightarrow P = 46\%$
2-6 keV	$\chi^2/\text{d.o.f.} = 25.8/24 \rightarrow P = 36\%$	$\chi^2/\text{d.o.f.} = 21.2/24 \rightarrow P = 63\%$
6-14 keV	$\chi^2/\text{d.o.f.} = 25.5/24 \rightarrow P = 38\%$	$\chi^2/\text{d.o.f.} = 35.9/24 \rightarrow P = 6\%$

no diurnal variation with a significance of 95% C.L.

+ run test to verify the hypothesis that the positive and negative data points are randomly distributed. The lower tail probabilities (in the four energy regions) are: 43, 18, 7, 26% for the solar case and 54, 84, 78, 16% for the sidereal case.

Thus, the presence of any significant diurnal variation and of time structures can be excluded at the reached level of sensitivity.

# The time dependence of the counting rate

Eur. Phys. J. C 74 (2014) 2827

Expected signal counting rate in a given k–th energy bin:

$$S_k [v_{lab}(t)] \simeq S_k [v_s] + \left[ \frac{\partial S_k}{\partial v_{lab}} \right]_{v_s} [V_{Earth} A_m \cos \omega(t - t_0) + V_r A_d \cos \omega_{rot}(t - t_d)]$$


The ratio  $R_{dy}$  of the diurnal over annual modulation amplitudes is a model independent constant

• Annual modulation amplitude:  $S_m = \left[ \frac{\partial S_k}{\partial v_{lab}} \right]_{v_s} V_{Earth} B_m$

• Diurnal modulation amplitude:  $S_d = \left[ \frac{\partial S_k}{\partial v_{lab}} \right]_{v_s} V_r B_d$

$$R_{dy} = \frac{S_d}{S_m} = \frac{V_r B_d}{V_{Earth} B_m} \simeq 0.016 \quad \text{at LNGS latitude}$$

- Observed annual modulation amplitude in DAMA/LIBRA–phase1 in the (2–6) keV energy interval:  $(0.0097 \pm 0.0013)$  cpd/kg/keV
- Thus, the expected value of the diurnal modulation amplitude is  $\simeq 1.5 \times 10^{-4}$  cpd/kg/keV.
- When fitting the *single-hit* residuals with a cosine function with amplitude  $A_d$  as free parameter, period fixed at 24 h and phase at 14 h: all the diurnal modulation amplitudes are compatible with zero.



Energy	$A_d^{exp}$ (cpd/kg/keV)	$\chi^2/\text{d.o.f.}$	P
2–4 keV	$(2.0 \pm 2.1) \times 10^{-3}$	27.8/23	22%
2–5 keV	$-(1.4 \pm 1.6) \times 10^{-3}$	23.2/23	45%
2–6 keV	$-(1.0 \pm 1.3) \times 10^{-3}$	20.6/23	61%
6–14 keV	$(5.0 \pm 7.5) \times 10^{-4}$	35.4/23	5%

$A_d < 1.2 \times 10^{-3}$  cpd/kg/keV (90%CL)

The  $A_d$  values are compatible with zero, having random fluctuations around zero with  $\chi^2$  equal to 19.5 for 18 dof

Present experimental sensitivity more modest than the expected diurnal modulation amplitude derived from the DAMA/LIBRA–phase1 observed effect.

larger exposure DAMA/LIBRA–phase2&3 and/or DAMA/1ton (+lower energy threshold) offers increased sensitivity to such an effect



Moreover, DAMA/LIBRA-phase2&3 and/or DAMA/1ton will also allow:

high sensitivities investigation on other rare processes:  
*possible PEP violating processes, various possible CNC processes in  $^{23}\text{Na}$  and  $^{127}\text{I}$ , nucleon and di-nucleon decay into invisible channels with new approach in  $^{23}\text{Na}$  and  $^{127}\text{I}$ , exotic particles (e.g. SIMPs, neutral nuclearities, Q-balls), solar axions by Primakoff effect in NaI(Tl), rare nuclear processes in  $^{23}\text{Na}$ ,  $^{127}\text{I}$ , hypothesized neutral particles (new QED phase) in  $^{241}\text{Am}$  decays, etc.*

... towards a 100 ton highly radiopure NaI(Tl) set-up for high-resolution full-spectroscopy solar neutrinos  
(Astrop.Phys.4(1995)45)