

AdS black hole thermodynamics with scalar hair

Ioannis Papadimitriou

SISSA and INFN - Sezione di Trieste, Italy



Gauge/Gravity Duality 2015

GGI, Florence

Based on [hep-th/0505190](https://arxiv.org/abs/hep-th/0505190), [hep-th/0703152](https://arxiv.org/abs/hep-th/0703152) and work in progress with K. Skenderis

- Asymptotically AdS black holes with scalar hair play a central role in applications of the AdS/CFT correspondence, especially to condensed matter physics.
- There has been a number of papers in the last year or two claiming that in order to satisfy the first law for certain hairy black holes one needs to introduce ‘scalar charges’ in the first law.
- Using the AdS/CFT dictionary as a guiding principle to formulate the variational problem for asymptotically AdS black holes with mixed boundary conditions on the scalars we demonstrate that this is not correct and the first law holds in its standard form once the conserved charges are properly defined.
- Although AdS/CFT language is used to study the problem, it is crucial to emphasize that there is no inherent assumption or requirement for a holographic duality – it is simply a matter of formulating a well posed variational problem in asymptotically locally AdS spacetimes.

- 1 Multi-trace deformations in the large- N limit
- 2 The AdS Dirichlet problem & holographic renormalization
- 3 Neumann or mixed scalar boundary conditions
- 4 An example
- 5 Concluding remarks

Outline

- 1 Multi-trace deformations in the large- N limit
- 2 The AdS Dirichlet problem & holographic renormalization
- 3 Neumann or mixed scalar boundary conditions
- 4 An example
- 5 Concluding remarks

Multi-trace deformations in the large-N limit

- Consider a local, gauge-invariant, scalar single-trace operator $\mathcal{O}(x)$, transforming in some representation of the relevant rank N (gauge/global symmetry) group.
- Let us deform the theory by a generic multi-trace operator built out of $\mathcal{O}(x)$:

$$S \rightarrow S + \int d^d x f(\mathcal{O})$$

where

$$f(\mathcal{O}) = c_2 \mathcal{O}^2 + \dots + c_k \mathcal{O}^k$$

is a polynomial in \mathcal{O} .

- If we insist that this is a relevant or marginal deformation of the theory, then

$$2 \leq k \leq \frac{d}{\Delta} \Rightarrow \Delta \leq \frac{d}{2}$$

- Together with unitarity this gives

$$\frac{d}{2} - 1 \leq \Delta \leq \frac{d}{2}$$

- Let us normalize \mathcal{O} such that $\langle \mathcal{O} \rangle = \mathcal{O}(N^0)$ as $N \rightarrow \infty$ and assume for concreteness that \mathcal{O} transforms in the adjoint representation so that $S \sim N^2$
- Generating function:

$$e^{-W[J]} = \int [d\phi] e^{-S[\phi] - N^2 \int d^d x J(x) \mathcal{O}(x)} \equiv e^{-N^2 w[J]}$$

with

$$\sigma(x) \equiv \langle \mathcal{O} \rangle_J = \frac{\delta w[J]}{\delta J} = \mathcal{O}(N^0)$$

- Effective action:

$$e^{-\Gamma[\sigma]} = \int [dJ] e^{-N^2 w[J] + N^2 \int d^d x J(x) \sigma(x)} \equiv e^{-N^2 \bar{\Gamma}[\sigma]}$$

with

$$J(x) = -\frac{\delta \bar{\Gamma}[\sigma]}{\delta \sigma} = \mathcal{O}(N^0)$$

- I now want to ask how do $w[J]$ and $\bar{\Gamma}[\sigma]$ transform under the multi-trace deformation

$$S[\phi] \rightarrow S_f[\phi] = S[\phi] + N^2 \int d^d x f(\mathcal{O})$$

- The answer in general depends on the details of the theory and the deformation, but large-N factorization leads to a universal result in the large-N limit...

$$\begin{aligned}
e^{-N^2 w_f[J_f]} &= \int [d\phi] e^{-S[\phi] - N^2 \int d^d x (J_f \mathcal{O} + f(\mathcal{O}))} \\
&= \int [d\phi] e^{-S[\phi] - N^2 \int d^d x (J \mathcal{O} + f(\mathcal{O}) - f'(\sigma) \mathcal{O})} \\
&\stackrel{N \rightarrow \infty}{\approx} e^{-N^2 w[J]} e^{-N^2 \int d^d x (f(\sigma) - \sigma f'(\sigma))}
\end{aligned}$$

$$J \equiv J_f + f'(\sigma)$$

$$\begin{aligned}
e^{-N^2 \bar{\Gamma}_f[\sigma]} &= \int [dJ_f] e^{-N^2 w_f[J_f] + N^2 \int d^d x J_f \sigma} \\
&\stackrel{N \rightarrow \infty}{\approx} \int [dJ] e^{-N^2 w[J]} e^{-N^2 \int d^d x (f(\sigma) - \sigma f'(\sigma))} e^{N^2 \int d^d x (J - f'(\sigma)) \sigma} \\
&= e^{-N^2 \bar{\Gamma}[\sigma] - N^2 \int d^d x f(\sigma)}
\end{aligned}$$

$$[dJ_f] = [dJ]$$

undeformed	deformed
J	$J_f = J - f'(\sigma)$
σ	$\sigma_f = \sigma$
$w[J]$	$w_f[J_f] = w[J] + \int d^d x (f(\sigma) - \sigma f'(\sigma)) _{\sigma = \frac{\delta w}{\delta J}}$
$\bar{\Gamma}[\sigma]$	$\bar{\Gamma}_f[\sigma] = \bar{\Gamma}[\sigma] + \int d^d x f(\sigma)$

- Let us now put the theory on a general background metric $g_{(0)ij}$ so that

$$\begin{aligned} w_f[J_f] &= w[J] + \int d^d x \sqrt{-g_{(0)}} (f(\sigma) - \sigma f'(\sigma)) \\ &= w[J_f] + \int d^d x \sqrt{-g_{(0)}} f(\sigma) \end{aligned}$$

- It follows that the stress tensor

$$\langle T^{ij} \rangle = - \frac{2}{\sqrt{-g_{(0)}}} \frac{\delta w}{\delta g_{(0)ij}}$$

transforms as

$$\langle T^{ij} \rangle[J, g_{(0)}] \rightarrow \langle T^{ij} \rangle_f[J_f, g_{(0)}] = \langle T^{ij} \rangle[J_f, g_{(0)}] - f(\sigma) g_{(0)}{}^{ij}$$

Outline

- 1 Multi-trace deformations in the large- N limit
- 2 The AdS Dirichlet problem & holographic renormalization**
- 3 Neumann or mixed scalar boundary conditions
- 4 An example
- 5 Concluding remarks

The Dirichlet problem for AdS gravity

- Consider first the Dirichlet boundary value problem for asymptotically locally AdS solutions of the action

$$S = \frac{1}{2\kappa^2} \int d^{d+1}x \sqrt{-g} \left(R - \frac{1}{4} Z(\varphi) F_{\mu\nu} F^{\mu\nu} - \frac{1}{2} (\partial\varphi)^2 - V(\varphi) \right)$$

$$V(\varphi) = -\frac{d(d-1)}{\ell^2} + \frac{1}{2} m^2 \varphi^2 + \dots$$

- Adding the Gibbons-Hawking term and the boundary counterterms, a generic variation of the total action takes the form

$$\delta(S + S_{GH} + S_{ct}) = \int d^{d+1}x (eoms) \delta\psi + \int d^d x \sqrt{-g_{(0)}} \left(\hat{\pi}_{(d)}^{ij} \delta g_{(0)ij} + \hat{\pi}_{(d)}^i \delta A_{(0)i} + \hat{\pi}_{(\Delta_+)} \delta\varphi_- \right)$$

where $\hat{\pi}_{(d)}^{ij}$, $\hat{\pi}_{(d)}^i$, $\hat{\pi}_{(\Delta_+)}$ are the renormalized radial momenta.

- Note that the counterterms are essential for the variational problem on the conformal boundary to be well posed [I.P, Skenderis '05]
- On a conformal boundary the bulk fields only induce a conformal class of boundary fields,

$$[g_{(0)}, A_{(0)}, \varphi_-]$$

with

$$(g_{(0)}, A_{(0)}, \varphi_-) \sim (e^{2\sigma(x)} g_{(0)}, A_{(0)}, e^{-(d-\Delta_+)\sigma(x)} \varphi_-)$$

and so the Dirichlet problem only makes sense in terms of conformal classes – not representatives! This is a sharp difference with the Dirichlet problem on a finite cut-off for which the Gibbons-Hawking term suffices.

- In the absence of a conformal anomaly

$$\int d^d x \sqrt{-g_{(0)}} \left(\hat{\pi}_{(d)}^{ij} \delta g_{(0)ij} + \hat{\pi}_{(d)}^i \delta A_{(0)i} + \hat{\pi}_{(\Delta_+)} \delta \varphi_- \right)$$

is a *class function*. Rendering this boundary term a class function is the most fundamental property of the boundary counterterms. Finiteness of the on-shell action follows as a consequence.

A comment on scheme dependence

- The local boundary counterterms are unambiguous except for the possibility of adding *finite* local boundary terms, which is interpreted as the renormalization scheme dependence in the dual theory.
- However, defining the counterterms through the variational problem imposes a severe constraint on the admissible finite local boundary terms that is not visible through the mere requirement of canceling the UV divergences:

Any finite local counterterms must be class functions, i.e. conformal invariants.

- Might have implications for supersymmetric scheme on curved boundaries [[Assel, Cassani, Di Pietro, Komargodski, Lorenzen, Martelli '15](#)]

- The AdS/CFT dictionary identifies the renormalized radial canonical momenta $\widehat{\pi}_{(d)}^{ij}$, $\widehat{\pi}_{(d)}^i$, $\widehat{\pi}_{(\Delta_+)}$ with the one-point functions of the dual operators, namely

$$\langle T^{ij} \rangle = 2\widehat{\pi}_{(d)}^{ij} \quad \langle \mathcal{J}^i \rangle = -\widehat{\pi}_{(d)}^i \quad \langle \mathcal{O}_{\Delta_+} \rangle = -\widehat{\pi}_{(\Delta_+)}$$

- Using the transformation of the sources under Penrose-Brown-Henneaux bulk diffeomorphisms one finds that these one-point functions satisfy the Ward identities

$$2D_{(0)i}\widehat{\pi}_{(d)j}^i + \widehat{\pi}_{(d)}^i F_{(0)ij} - \widehat{\pi}_{(\Delta_+)} \partial_j \varphi_- = 0$$

$$D_{(0)i}\widehat{\pi}_{(d)}^i = 0$$

$$2\widehat{\pi}_{(d)i}^i - (d - \Delta_+)\widehat{\pi}_{(\Delta_+)}\varphi_- = \mathcal{A}[g_{(0)}, A_{(0)}, \varphi_-]$$

where $\mathcal{A}[g_{(0)}, A_{(0)}, \varphi_-]$ is the trace anomaly.

- These Ward identities imply that the quantities

$$\mathcal{Q}[\xi] := \int_{\partial\mathcal{M}\cap C} (2\widehat{\pi}_{(d)j}^i + \widehat{\pi}_{(d)}^i A_{(0)j}) \xi^j = \int_{\partial\mathcal{M}\cap C} (\langle T_j^i \rangle - \langle \mathcal{J}^i \rangle A_{(0)j}) \xi^j$$

are independent of the boundary Cauchy surface C iff [I.P., Skedneris '05]:

- 1 ξ^i is a boundary conformal Killing vector, i.e.

$$\mathcal{L}_\xi g_{(0)ij} = \frac{2}{d} (D_{(0)k} \xi^k) g_{(0)ij} \quad \mathcal{L}_\xi \varphi_- = -\frac{d - \Delta_+}{d} (D_{(0)k} \xi^k) \varphi_-, \quad \mathcal{L}_\xi A_{(0)i} = 0$$

and the trace anomaly is *numerically zero*, or

- 2 ξ^i is a boundary Killing vector, i.e.

$$\mathcal{L}_\xi g_{(0)ij} = 0 \quad \mathcal{L}_\xi \varphi_- = 0, \quad \mathcal{L}_\xi A_{(0)i} = 0$$

and the trace anomaly is *numerically non-zero*.

- These charges agree with the Wald Hamiltonians and the Ashtekar-Magnon-Das mass, in a particular choice of scheme [I.P., Skenderis '05].

Smarr formula and the first law

- These conserved charges satisfy the Smarr formula

$$I = \beta G(T, \Omega_i, \Phi) = \beta (M - TS - \Omega_i J_i - \Phi Q)$$

and the first law

$$\delta M = T\delta S + \Omega_i \delta J_i + \Phi \delta Q$$

for any asymptotically locally AdS black hole with Dirichlet boundary conditions on all fields [I.P., Skenderis '05].

- The only scheme-dependent observables are the free energy I and the mass M , but the scheme dependence cancels in the Smarr formula. The scheme dependence does not affect the first law since the local finite counterterms are class functions.
- In the presence of a conformal anomaly

$$\delta_\sigma I = \beta \delta_\sigma M = \int d^d x \sqrt{-g_{(0)}} \mathcal{A} \delta \sigma$$

and so the variations in the first law must ensure that the conformal representative is held fixed (important e.g. for the Kerr-AdS₅ black hole)

Outline

- 1 Multi-trace deformations in the large- N limit
- 2 The AdS Dirichlet problem & holographic renormalization
- 3 Neumann or mixed scalar boundary conditions
- 4 An example
- 5 Concluding remarks

Neumann and mixed boundary conditions

- Let us parameterize the general asymptotic form of scalar field φ in asymptotically locally AdS_{d+1} as (except when BF bound is saturated)

$$\varphi = \varphi_-(x)e^{-\Delta_- r} + \dots + \varphi_+(x)e^{-\Delta_+ r} + \dots$$

where

$$\Delta_- + \Delta_+ = d, \quad m^2 \ell^2 = -\Delta_- \Delta_+$$

- For scalars with mass squared in the window

$$-\left(\frac{d}{2}\right)^2 \leq m^2 \leq -\left(\frac{d}{2}\right)^2 + 1$$

both modes φ_- and φ_+ are normalizable [Breitenlohner, Freedman '82; Balasubramanian, Kraus, Lawrence '98; Klebanov, Witten '99] and so any desirable combination of these modes can be held fixed in the variational problem.

- Note that these bounds on the mass squared are equivalent to

$$\frac{d}{2} - 1 \leq \Delta_- \leq \frac{d}{2}$$

i.e. Δ_- must satisfy the unitarity bound.

Additional boundary terms

- In order to modify the boundary conditions we start with the *renormalized* action corresponding to Dirichlet boundary conditions

$$S + S_{GH} + S_{ct}$$

and we add an extra *finite* boundary term

$$S_J[\varphi_-, \hat{\pi}_{(\Delta_+)}]$$

such that

$$\delta(S + S_{GH} + S_{ct} + S_J) \sim \int d^d x \sqrt{-g_{(0)}} B(\varphi_-, \hat{\pi}_{(\Delta_+)}) \delta J$$

where

$$J(\varphi_-, \hat{\pi}_{(\Delta_+)})$$

is any function of the scalar modes that we decide to keep fixed in the variational problem.

- For Dirichlet, Neumann and Mixed boundary conditions the additional boundary term S_J respectively takes the form:

$J(\varphi_-, \hat{\pi}(\Delta_+))$	$S_J[\varphi_-, \hat{\pi}(\Delta_+)]$
$J_+ = \varphi_-$ $J_- = \hat{\pi}(\Delta_+)$ $J_{f_-} = \hat{\pi}(\Delta_+) - f'(\varphi_-)$	$S_+ = 0$ $S_- = - \int d^d x \sqrt{-g(0)} \varphi_- \hat{\pi}(\Delta_+)$ $S_{f_-} = S_- - \int d^d x \sqrt{-g(0)} (f(\varphi_-) - \varphi_- f'(\varphi_-))$

- Note that it is the renormalized canonical momentum

$$\hat{\pi}(\Delta_+) \sim (\Delta_+ - \Delta_-) \ell^{-1} \varphi_+ + c(\varphi_-)$$

that enters these boundary terms – not φ_+ .

Holographic dictionary

	Dirichlet	Neumann	Mixed
J	$J_+ \equiv \varphi_-$	$J_- \equiv \widehat{\pi}_{(\Delta_+)}$	$J_{f_-} \equiv \widehat{\pi}_{(\Delta_+)} - f'(\varphi_-)$
σ	$-\widehat{\pi}_{(\Delta_+)}$	φ_-	φ_-
$W[J]$	$-S_D[J_+]$	$-S_N[J_-]$	$-S_M[J_{f_-}]$
$\Gamma[\sigma]$	$-S_N[\widehat{\pi}_{(\Delta_+)}]$	$-S_D[\varphi_-]$	$-S_D[\varphi_-] + \int d^d x \sqrt{-g_{(0)}} f(\varphi_-)$
$\langle T^{ij} \rangle$	$2\widehat{\pi}_{(d)}^{ij}$	$2\widehat{\pi}_{(d)}^{ij} - \varphi_- J_- g_{(0)}^{ij}$	$2\widehat{\pi}_{(d)}^{ij} - \left(f(\varphi_-) + \varphi_- J_{f_-} \right) g_{(0)}^{ij}$

- Comparing these quantities with the large- N transformation of the generating functional, effective action and stress tensor under a multi-trace transformation one concludes that:

Imposing Mixed boundary conditions corresponds to deforming the theory with Neumann boundary conditions with the multi-trace deformation $f(\varphi_-)$ [Witten '01].

- Recall that the condition for such a deformation to be relevant or marginal is

$$\frac{d}{2} - 1 \leq \Delta \leq \frac{d}{2}$$

which is only possible in the Δ_- quantization. Moreover, this condition on Δ_- coincides with the restriction on the scalar mass in order for the Δ_- quantization to be admissible.

Ward identities in the deformed theory

- Using the Ward identities of the Dirichlet problem and the above transformations leads to the following Ward identities for the deformed theory:

$$D_{(0)i} \langle T_j^i \rangle - \langle \mathcal{J}^i \rangle F_{(0)ij} + \langle \mathcal{O}_{\Delta_-} \rangle \partial_j J_{f_-} = 0$$

$$D_{(0)i} \langle \mathcal{J}^i \rangle = 0$$

$$\langle T_i^i \rangle + (d - \Delta_-) \langle \mathcal{O}_{\Delta_-} \rangle J_{f_-} = \mathcal{A}[g_{(0)}, A_{(0)}, \varphi_-] - d \left(f(\varphi_-) - \frac{\Delta_-}{d} \varphi_- f'(\varphi_-) \right)$$

- The first two Ward identities are identical in form to the Dirichlet ones, except that the sources and one-point functions have been transformed.
- Moreover, writing

$$f(\varphi_-) = \sum_k J_k \varphi_-^k$$

the trace Ward identity becomes

$$\langle T_i^i \rangle + (d - \Delta_-) J_{f_-} \langle \mathcal{O}_{\Delta_-} \rangle + \sum_k (d - \Delta_k) J_k \langle \mathcal{O}_{\Delta_-}^k \rangle = \mathcal{A}[g_{(0)}, A_{(0)}, \varphi_-]$$

where $\Delta_k = k\Delta_-$.

Trace anomaly

- The trace anomaly $\mathcal{A}[g_{(0)}, A_{(0)}, \varphi_-]$ is the same trace anomaly that appears in the Dirichlet problem. However, while in the Dirichlet problem φ_- is the source, in the Neumann and mixed problem it is the one-point function! So if the anomaly depends on φ_- , then there is an apparent problem.
- However, the fact that the scaling dimension Δ_- must satisfy the unitarity bound for Neumann or mixed boundary conditions to be admissible, together with the fact that the anomaly is a conformal invariant implies that the only way that φ_- can enter in the anomaly is

$$\mathcal{A}_{scalar} = \alpha \left(\partial^i \varphi_- \partial_i \varphi_- + \frac{(d-2)}{4(d-1)} R[g_{(0)}] \varphi_-^2 \right) + \beta \varphi_-^{\frac{d}{\Delta_-}}$$

where $\alpha = 0$ unless $\Delta_- = (d-2)/2$ and $\beta = 0$ unless d/Δ_- is a positive integer.

- The correct prescription to obtain the generating functional and correlation functions of the operator \mathcal{O}_{Δ_-} is to use dimensional regularization, compute the final result, and only in the end set the dimension to one of the critical dimensions for which such anomalies can potentially occur [Klebanov, Witten '99]. Since there can only be scalar contributions to the anomaly at discrete values of the dimension it follows that the conformal anomaly does not contain scalar contributions.

Conserved charges

- The Ward identities of the deformed theory suggest that conserved charges are now given by

$$\mathcal{Q}[\xi] := \int_{\partial\mathcal{M} \cap \mathcal{C}} (\langle T_j^i \rangle - \langle \mathcal{J}^i \rangle A_{(0)j}) \xi^j$$

in terms of the stress tensor in the **deformed theory**.

- Indeed, these charges are conserved provided:

- 1 ξ^i is a boundary conformal Killing vector of the **deformed sources**, i.e.

$$\mathcal{L}_\xi g_{(0)ij} = \frac{2}{d} (D_{(0)k} \xi^k) g_{(0)ij} \quad \mathcal{L}_\xi J_{f-} = -\frac{d - \Delta_-}{d} (D_{(0)k} \xi^k) J_{f-}, \quad \mathcal{L}_\xi A_{(0)i} = 0$$

the trace anomaly is *numerically zero*, **and** $f(\varphi_-) = \lambda \varphi_-^{d/\Delta_-}$, or

- 2 ξ^i is a boundary Killing vector of the **deformed sources**, i.e.

$$\mathcal{L}_\xi g_{(0)ij} = 0 \quad \mathcal{L}_\xi J_{f-} = 0, \quad \mathcal{L}_\xi A_{(0)i} = 0$$

and either the trace anomaly is numerically non-zero or $f(\varphi_-)$ is not marginal, or both.

Smarr formula and the first law

- These conserved charges again satisfy the Smarr formula

$$I = \beta G(T, \Omega_i, \Phi) = \beta (M - TS - \Omega_i J_i - \Phi Q)$$

and the first law

$$\delta M = T\delta S + \Omega_i \delta J_i + \Phi \delta Q$$

for any asymptotically locally AdS black hole with Dirichlet/Neumann/Mixed boundary conditions [I.P., Skenderis '15].

- All charges must be computed with the stress tensor of the deformed theory.
- The variations in the first law must keep fixed the multi-trace couplings in $f(\varphi_-)$, in addition to the single trace sources.
- AMD mass does not give the correct result for mixed boundary conditions!
- Wald Hamiltonians give the correct result provided the variations respect the boundary conditions.

Outline

- 1 Multi-trace deformations in the large- N limit
- 2 The AdS Dirichlet problem & holographic renormalization
- 3 Neumann or mixed scalar boundary conditions
- 4 An example
- 5 Concluding remarks

An example

- A black hole that was recently presented as an example for the necessity of scalar charges can be found in 3 dimensions ($d = 2$) for the scalar potential [Wen '15]

$$V(\varphi) = g^2 (\cosh \psi)^{4\mu} (\mu \tanh^2 \psi - 2) = -g^2 \left(2 + \frac{3}{8} \varphi^2 + \frac{1}{32} \varphi^4 + \mathcal{O}(\varphi^5) \right)$$

where $g = 1/\ell$, $\psi = \varphi/(2\sqrt{2\mu})$, and

$$-1 < m^2 \ell^2 = -3/4 < 0, \quad \Delta_- = 1/2, \quad \Delta_+ = 3/2$$

- The black hole solution takes the form

$$\varphi(r) = 2\sqrt{2\mu} \operatorname{Arctanh} \left(\frac{1}{\sqrt{1+r/q}} \right),$$
$$ds^2 = -g^2 r^2 dt^2 + \frac{dr^2}{g^2 r^2 (1+q/r)^{2\mu}} + r^2 d\theta^2$$

where q is the only parameter of the solution.

- The boundary counterterms are

$$S_{ct} = -\frac{1}{\kappa^2} \int d^d x \sqrt{-\gamma} \left(\frac{1}{\ell} + \frac{1}{8\ell} \varphi^2 \right)$$

- This leads to the renormalized metric momentum

$$\hat{\pi}_{(2)ij} = \frac{1}{2\kappa^2} \frac{g}{6} \mu(3\mu - 1) q^2 \eta_{ij}$$

- Moreover, in Fefferman-Graham coordinates

$$\phi = 2\sqrt{2\mu q} e^{-g\bar{r}/2} + \frac{(3\mu - 1)}{48\mu} 8(2\mu q)^{3/2} e^{-3g\bar{r}/2} + \mathcal{O}(e^{-5g\bar{r}/2})$$

and so

$$\hat{\pi}_{(\Delta_+)} = \frac{g}{2\kappa^2} (\Delta_+ - \Delta_-) \varphi_+ = f'(\varphi_-) = \frac{g}{2\kappa^2} \frac{(3\mu - 1)}{48\mu} \varphi_-^3 \Rightarrow$$

$$f(\varphi_-) = \frac{g}{2\kappa^2} \frac{(3\mu - 1)}{192\mu} \varphi_-^4 = \frac{g}{2\kappa^2} \frac{1}{3} \mu(3\mu - 1) q^2$$

- The stress tensor vanishes identically:

$$\begin{aligned}\langle T^{ij} \rangle &= 2\hat{\pi}_{(2)}^{ij} - g\eta^{ij}f(\varphi_-) \\ &= 2 \times \frac{1}{2\kappa^2} \frac{g}{6} \mu(3\mu - 1)q^2\eta^{ij} - \frac{g}{2\kappa^2} \frac{1}{3} \mu(3\mu - 1)q^2\eta^{ij} = 0\end{aligned}$$

and so

$$M = 0$$

- This agrees with the Wald mass and is consistent with the first law

$$\delta M = T\delta S$$

since the entropy vanishes identically too.

Outline

- 1 Multi-trace deformations in the large- N limit
- 2 The AdS Dirichlet problem & holographic renormalization
- 3 Neumann or mixed scalar boundary conditions
- 4 An example
- 5 Concluding remarks**

Concluding remarks

- The AdS/CFT dictionary provides a clear prescription for formulating the variational problem for asymptotically AdS black holes with scalar fields satisfying mixed boundary conditions.
- This leads to conserved charges that are compatible with the first law in its standard form.
- No scalar charges!