

A twisted index for three dimensional gauge theories

Alberto Zaffaroni

Università di Milano-Bicocca

Gauge/Gravity Duality 2015, GGI

Outline

- ▶ Motivations
- ▶ The twisted index
- ▶ Examples and generalizations
- ▶ Conclusions

Outline

- ▶ Motivations
- ▶ The twisted index
- ▶ Examples and generalizations
- ▶ Conclusions

based on

F. Benini, A.Z. 1504.xxxxx

Motivations

Lot of recent activity in the study of supersymmetric and superconformal theories in various dimensions in closely related contexts

- ▶ Progresses in the evaluation of exact quantum observables.
- ▶ Related to localization and the study of supersymmetry in curved space.

[Pestun + Festuccia-Seiberg + many . . .]

Motivations

Here we consider a very simple example, a 3d gauge theory on $S^2 \times S^1$ where susy is preserved by a twist on S^2

$$(\nabla_\mu - iA_\mu^R)\epsilon \equiv \partial_\mu \epsilon = 0, \quad \int_{S^2} F^R = 1$$

The result becomes interesting when supersymmetric backgrounds for the flavor symmetry multiplets (A_μ^F, σ^F, D^F) are turned on:

$$u^F = A_t^F + i\sigma^F, \quad q^F = \int_{S^2} F^F = iD^F$$

and the path integral becomes a function of a set of magnetic charges q^F and chemical potentials u^F . We can also add a refinement for angular momentum.

Motivations

Notice: we are not computing the superconformal index of the 3d gauge theory.

It is rather a **twisted index**: a trace over the Hilbert space \mathcal{H} of states on a sphere in the presence of a magnetic background for the R and the global symmetries,

$$\mathrm{Tr}_{\mathcal{H}} \left((-1)^F e^{iJ_F A^F} e^{-\beta H} \right)$$

$$Q^2 = H - \sigma^F J_F$$

holomorphic in u^F

where J_F is the generator of the global symmetry.

Motivations

The original motivation for this work comes holography.

CFTs on curved space-times described by dual regular asymptotically AdS backgrounds

$$ds_4^2 = \frac{dr^2}{r^2} + (r^2 ds_{M_3}^2 + O(r)) \quad A = A_{M_3} + O(1/r)$$

Classifications of M_3 supersymmetric backgrounds (transverse holomorphic foliations)

[Klare-Tomasiello-AZ; Closset-Dumitrescu-Festuccia-Komargodski]

Motivations

Twisted $M_3 = S^2 \times S^1$ leads to **1/4 BPS** asymptotically AdS_4 static black holes

- ▶ solutions asymptotic to *magnetic AdS_4* and with horizon $\text{AdS}_2 \times S^2$
- ▶ Characterized by a collection of magnetic charges $\int_{S^2} F$
- ▶ preserving supersymmetry via a twist

$$(\nabla_\mu - iA_\mu)\epsilon = \partial_\mu \epsilon \quad \implies \quad \epsilon = \text{const}$$

Various solutions with regular horizons, some embeddable in $\text{AdS}_4 \times S^7$.


[Cacciatori, Klemm; Gnechchi, Dall'agata; Hristov, Vandoren];

Motivations

CFTs on curved space-times described by dual regular asymptotically AdS backgrounds

$$ds_{d+1}^2 = \frac{dr^2}{r^2} + (r^2 ds_{M_d}^2 + O(r)) \quad A = A_{M_d} + O(1/r)$$

[A.Z. with Benini, Hristov, Tomasiello]



AdS₄

AdS₂ × S₂

Entropy of black holes
Counting of microstates

Partition function of twisted
3d CFT on S₂ × S₁

QM fixed point

The background

Consider an $\mathcal{N} = 2$ gauge theory on $S^2 \times S^1$

$$ds^2 = R^2(d\theta^2 + \sin^2 \theta d\varphi^2) + \beta^2 dt^2$$

with a background for the R-symmetry proportional to the spin connection:

$$A^R = -\frac{1}{2} \cos \theta d\varphi = -\frac{1}{2} \omega^{12}$$

so that the Killing spinor equation

$$D_\mu \epsilon = \partial_\mu \epsilon + \frac{1}{4} \omega_\mu^{ab} \gamma_{ab} \epsilon - i A_\mu^R \epsilon = 0 \quad \implies \quad \epsilon = \text{const}$$

The partition function

The path integral for an $\mathcal{N} = 2$ gauge theory on $S^2 \times S^1$ with gauge group G localizes on a set of BPS configurations specified by data in the vector multiplets

$$V = (A_\mu, \sigma, \lambda, \lambda^\dagger, D)$$

- ▶ A magnetic flux on S^2 , $\mathfrak{m} = \frac{1}{2\pi} \int_{S^2} F$ in the co-root lattice
- ▶ A Wilson line A_t along S^1
- ▶ The vacuum expectation value σ of the real scalar

Up to gauge transformations, the BPS manifold is

$$(u = A_t + i\sigma, \mathfrak{m}) \in \mathcal{M}_{\text{BPS}} = (H \times \mathfrak{h} \times \Gamma_{\mathfrak{h}}) / W$$

The partition function

The path integral reduces to a the saddle point around the BPS configurations

$$\sum_{\mathfrak{m} \in \Gamma_{\mathfrak{h}}} \int dud\bar{u} \mathcal{Z}^{\text{cl} + 1\text{-loop}}(u, \bar{u}, \mathfrak{m})$$

- ▶ The integrand has various singularities where chiral fields become massless
- ▶ There are fermionic zero modes

The two things nicely combine and the path integral reduces to an r -dimensional contour integral of a meromorphic form

$$\frac{1}{|W|} \sum_{\mathfrak{m} \in \Gamma_{\mathfrak{h}}} \oint_{\mathcal{C}} Z_{\text{int}}(u, \mathfrak{m})$$

The partition function

The classical and 1-loop contribution gives a meromorphic form

$$Z_{\text{int}}(u, \mathfrak{m}) = Z_{\text{class}} Z_{\text{1-loop}}$$

where

$$Z_{\text{class}}^{\text{CS}} = x^{k\mathfrak{m}}$$

$$Z_{\text{1-loop}}^{\text{chiral}} = \prod_{\rho \in \mathfrak{R}} \left[\frac{x^{\rho/2}}{1 - x^{\rho}} \right]^{\rho(\mathfrak{m}) - q + 1}$$

$$Z_{\text{1-loop}}^{\text{gauge}} = \prod_{\alpha \in G} (1 - x^{\alpha}) (i du)^r$$

The partition function

The magnetic flux on S^2 generates Landau levels. Massive bosons and fermions cancel in pairs, while zero modes

$$\begin{array}{ll}
 \rho(\mathfrak{m}) - q + 1 > 0 & \text{Fermi multiplets on } S^1 \\
 \rho(\mathfrak{m}) - q + 1 < 0 & \text{Chiral multiplets on } S^1
 \end{array}$$

$$Z_{1\text{-loop}}^{\text{chiral}} = \prod_{\rho \in \mathfrak{A}} \left[\frac{x^{\rho/2}}{1 - x^\rho} \right]^{\rho(\mathfrak{m}) - q + 1}$$

The contour

$Z_{\text{int}}(u, \mathfrak{m})$ has pole singularities at

- ▶ along the hyperplanes $x^\rho = e^{i\rho(u)} = \mathbb{1}_G$ determined by the chiral fields
- ▶ at the boundaries of $H \times \mathfrak{h}$ ($\text{Im}(u) = \pm\infty$, $x = e^{iu} = 0, \infty$)

Supersymmetric localization selects a particular contour of integration C and picks some of the residues of the form $Z_{\text{int}}(u, \mathfrak{m})$.

The contour

Consider a $U(1)$ theory with chiral fields with charges Q_i . We can use the prescription: sum the residues

- ▶ at the poles of fields with positive charge, at $x = 0$ if $k_{\text{eff}}(+\infty) < 0$ and at $x = \infty$ if $k_{\text{eff}}(-\infty) > 0$

where the effective Chern-Simons coupling is defined as

$$k_{\text{eff}}(\sigma) = k + \frac{1}{2} \sum_i Q_i^2 \text{sign}(Q_i \sigma)$$

The contour

The prescription can be written in a compact form by using the so-called Jeffrey-Kirwan residue

$$\text{JK-Res}_{y=0}(Q, \eta) \frac{dy}{y} = \theta(Q\eta) \text{sign}(Q)$$

as

$$\frac{1}{|W|} \sum_{\mathfrak{m} \in \Gamma_{\mathfrak{h}}} \left[\sum_{x_* \in \mathfrak{M}_{\text{sing}}} \text{JK-Res}(Q(x_*), \eta) Z_{\text{int}}(x; \mathfrak{m}) + \text{JK-Res}_{x=0, \infty}(Q_x, \eta) Z_{\text{int}}(x; \mathfrak{m}) \right]$$

where

$$Q_{x=0} = -k_{\text{eff}}(+\infty), \quad Q_{x=\infty} = k_{\text{eff}}(-\infty)$$

Similar to the localization of the elliptic genus for 2d theories and of the Witten index in 1d [Benini, Eager, Hori, Tachikawa; Hori, Kim, YI]

A Simple Example: $U(1)_{1/2}$ with one chiral

The theory has just a topological $U(1)_T$ symmetry: $J_\mu = \epsilon_{\mu\nu\tau} F_{\nu\tau}$

$$Z = \sum_{m \in \mathbb{Z}} \int \frac{dx}{2\pi i x} x^t (-\xi)^m x^{m/2} \left(\frac{x^{1/2}}{1-x} \right)^m = \frac{\xi}{(1-\xi)^{t+1}}$$

$$k_{\text{eff}}(\sigma) = \frac{1}{2} + \frac{1}{2} \text{sign}(\sigma) \quad \rightarrow \quad Q_{x=0} = -1, Q_{x=\infty} = 0$$

Consistent with duality with a free chiral.

	$U(1)_g$	$U(1)_T$	$U(1)_R$
X	1	0	1
\tilde{T}	0	1	0
\tilde{T}	-1	-1	0

Aharony and Giveon-Kutasov dualities

The twisted index can be used to check dualities: for example, $U(N_c)$ with $N_f = N_c$ flavors is dual to a theory of chiral fields M_{ab} , T and \tilde{T} , coupled through the superpotential $W = T\tilde{T} \det M$

$$Z_{N_f=N_c} = \left(\frac{y}{1-y^2} \right)^{(2n-1)N_c^2} \left(\frac{\xi^{\frac{1}{2}} y^{-\frac{N_c}{2}}}{1-\xi y^{-N_c}} \right)^{N_c(1-n)+t} \left(\frac{\xi^{-\frac{1}{2}} y^{-\frac{N_c}{2}}}{1-\xi^{-1} y^{-N_c}} \right)^{N_c(1-n)-t}$$

Aharony and Giveon-Kutasov dual pairs for generic (N_c, N_f) have the same partition function.

Refinement by angular momentum

Adding a fugacity $\zeta = e^{i\zeta/2}$ for the angular momentum on S^2 : the Landau zero-modes on S^2 form a representation of $SU(2)$.

$$Z_{1\text{-loop}}^{\text{chiral}} = \prod_{\rho \in \mathfrak{R}} \prod_{j=-\frac{|B|-1}{2}}^{\frac{|B|-1}{2}} \left(\frac{x^{\rho/2} \zeta^j}{1 - x^{\rho} \zeta^{2j}} \right)^{\text{sign } B}, \quad B = \rho(\mathfrak{m}) - q_{\rho} + 1$$

As noticed in other contexts: the refined partition function factorizes into the product of two vortex partition functions

$$Z = Z_{1\text{-loop}} Z_{\text{vortex}}(\zeta) Z_{\text{vortex}}(\zeta^{-1})$$

[Pasquetti;Beem-Dimofte-Pasquetti;Cecotti-Gaiotto-Vafa, . . .]

Other dimensions

We can consider other dimensions too: $(2, 2)$ theories in 2d on S^2

The BPS manifold is now $\mathfrak{M} = (\mathfrak{h} \times \mathfrak{h})/W$ and the 1-loop determinants

$$Z_{1\text{-loop}}^{\text{chiral}} = \prod_{\rho \in \mathfrak{R}} \left[\frac{1}{\rho(\sigma)} \right]^{\rho(\mathfrak{m}) - q + 1}$$

$$Z_{1\text{-loop}}^{\text{gauge}} = (-1)^{\sum_{\alpha > 0} \alpha(\mathfrak{m})} \prod_{\alpha \in G} \alpha(\sigma) (d\sigma)^r$$

Other dimensions

We are just repackaging results about the A-twist of gauged linear sigma models

For examples, for $U(1)$ with N flavors, 2d amplitudes compute the quantum cohomology of \mathbb{P}^{N-1}

$$\langle \sigma_1 \cdots \sigma_n \rangle = \sum_m \int \frac{dx}{2\pi i} \frac{1}{x^{(m+1)N}} q^m x^n = \sum_m q^m \delta_{N(m+1)-n-1,0}$$

$$\sigma^N = q$$

$$\prod_{j=1}^N (\sigma - \mu_j) = q$$

Ω - background and non abelian can be considered [see Cremonesi, Closset, Park, to appear]

Other dimensions

We can consider other dimensions too: $\mathcal{N} = 1$ theories in 4d on $S^2 \times T^2$

The BPS manifold is now $\mathfrak{M} = (H \times H)/W$ and the 1-loop determinants is elliptically generalized

$$Z_{1\text{-loop}}^{\text{chiral}} = \prod_{\rho \in \mathfrak{R}} \prod_{j=-\frac{|B|-1}{2}}^{\frac{|B|-1}{2}} \left(\frac{i\eta(q)}{\theta_1(q, x^\rho \zeta^{2j})} \right)^{\text{sign}(B)}$$

$$Z_{1\text{-loop}}^{\text{gauge, off}} = (-1)^{\sum_{\alpha > 0} \alpha(\mathfrak{m})} \prod_{\alpha \in G} \frac{\theta_1(q, x^\alpha \zeta^{|\alpha(\mathfrak{m})|})}{i\eta(q)} (du)^r$$

[also Closset-Shamir; Nishioka-Yaakov]

It can be tested against Seiberg's dualities.

Conclusions

We gave a general formula for the topologically twisted path integral of 3d $\mathcal{N} = 2$ theories.

- ▶ Higher genus $S^2 \rightarrow \Sigma$. Include Witten index
- ▶ 2d theories, Calabi-Yau's and sigma-models
- ▶ Large N limit analysis of the matrix model
- ▶ AdS_4 free-energy and entropy