

101° CONGRESSO DELLA SOCIETÀ ITALIANA DI FISICA

SEARCH FOR DARK MATTER AT LHC



21/09/2015



Marco Cipriani
Università di Roma La Sapienza
INFN Roma

Marco Cipriani




SHEDDING LIGHT ON DARK MATTER




- indirect search:

Dark matter (DM) **annihilation** into Standard Model (SM) particles




SHEDDING LIGHT ON DARK MATTER



- direct search:


DM scattering on nucleons



SHEDDING LIGHT ON DARK MATTER

- search at colliders:


DM **production** from proton-proton collisions at LHC




I will present CMS results

DARK MATTER SIGNAL AT LHC

MONOJET EVENT



exploit radiation from initial state as a trigger



look for events with high **missing transverse energy** (MET or \vec{E}_T)


SIGNAL SELECTION

→ 1 high energy jet recoiling against \cancel{E}_T

- **MET** > 200 GeV
- **Njets** ≤ 2
- **noise cleaning** on leading jet
- **jet 1** p_T > 110 GeV;
- **second jet** allowed if:
 $\Delta\phi(\text{jet1}, \text{jet2}) < 2.5$
- **photon veto**
- **lepton veto**

$$\vec{\cancel{E}}_T = - \sum_{\text{all particles}} \vec{p}_T$$

\vec{p}_T : momentum in transverse plane




SIGNAL SELECTION

→ 1 high energy jet recoiling against \cancel{E}_T

- **MET** > 200 GeV
- **Njets** ≤ 2
- **noise cleaning** on leading jet
- **jet 1** p_T > 110 GeV;
- **second jet** allowed if:
 $\Delta\phi(\text{jet1}, \text{jet2}) < 2.5$
- **photon veto**
- **lepton veto**


$$\vec{\cancel{E}}_T = - \sum_{\text{all particles}} \vec{p}_T$$

\vec{p}_T : momentum in transverse plane





MONOJET BACKGROUNDS

everything in SM that produces jets and \cancel{E}_T



signal



backgrounds

main backgrounds

- $Z(\nu\nu) + jets$ (55%)
- $W(l\nu) + jets$ (40%) (if l is lost)

estimated from data using control samples

others:


- $t\bar{t}$, single top
- QCD multijet
- diboson

estimated from MC


DATA DRIVEN BACKGROUND ESTIMATE

- select a **control sample in data**
- use **MC** to get **transfer factors** from control region (CR) to signal region (SR)

$$N_{SR}^{data} = N_{CR}^{data} \times \frac{N_{SR}^{MC}}{N_{CR}^{MC}}$$



MET defined here excluding Z or γ p_T in the sum






MET RESOLUTION AND RESPONSE


my contribution

- assess the performance of the hadronic recoil reconstruction


Use *Z(ll) + jets events*: induce MET by removing the well-measured leptons



RUN 1 RESULTS




CMS-PAS-EXO-12-055



complementarity with respect to direct searches
 collider searches are also sensitive to low DM mass

RUN 2 PERSPECTIVES



need 5 fb^{-1} at 13 TeV to be competitive with Run 1

- signal from fit to MET distribution: higher sensitivity than "cut&count"
- more control samples for background estimate

SUMMARY

- ✓ **monojet analysis for dark matter search at LHC presented**
 - simple experimental signature: 1 jet and \cancel{E}_T
- ✓ **Run 2 and data taking at 13 TeV at LHC has begun**
 - enhanced sensitivity thanks to higher energy collisions
 - with $5fb^{-1}$ we are as sensitive as previous Run 1 ($\approx 20fb^{-1}$)
- ✓ **many improvements in the analysis during Run 2**
 - more control samples to estimate the backgrounds
- ✓ **collider and direct searches complement each other**

BACKUP

COMPACT MUON SOLENOID

CMS DETECTOR

Total weight : 14,000 tonnes

Overall diameter : 15.0 m

Overall length : 28.7 m

Magnetic field : 3.8 T

STEEL RETURN YOKE
12,500 tonnes

SILICON TRACKERS

Pixel ($100 \times 150 \mu\text{m}$) $\sim 16\text{m}^2 \sim 66\text{M}$ channels

Microstrips ($80 \times 180 \mu\text{m}$) $\sim 200\text{m}^2 \sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID

Niobium titanium coil carrying $\sim 18,000\text{A}$

MUON CHAMBERS

Barrel: 250 Drift Tube, 480 Resistive Plate Chambers

Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers

PRESHOWER

Silicon strips $\sim 16\text{m}^2 \sim 137,000$ channels

FORWARD CALORIMETER


Steel + Quartz fibers $\sim 2,000$ Channels

CRYSTAL
ELECTROMAGNETIC
CALORIMETER (ECAL)


$\sim 76,000$ scintillating PbWO_4 crystals

HADRON CALORIMETER (HCAL)


Brass + Plastic scintillator $\sim 7,000$ channels



COMPACT MUON SOLENOID



COMPACT MUON SOLENOID



$$\eta = -\ln \operatorname{tg} \frac{\theta}{2}$$

η differences are Lorentz invariant for high energy particles

INTERPRETATIONS OF RUN 1 RESULTS

observed limit on cross section depends on DM mass and interaction with SM

can be translated into limit on DM-nucleon scattering cross section

different sensitivity from various experiments can lead to controversial results

