


# Higher order hadronic and leptonic contributions to the muon $g - 2$

Matthias Steinhauser

in collaboration with A. Kurz, T. Liu, P. Marquard, A. Smirnov, V. Smirnov | FCCP, September 10-12 2015


TTP KARLSRUHE



# Higher order hadronic and leptonic contributions to the muon $g - 2$

- I. NNLO hadronic contribution
- II. Electron contribution at 4 loops
- III. Tau contribution at 4 loops

# Muon $g - 2$



[Schwinger'48]



[Petermann'57;  
Sommerfeld'58]



[Laporta,Remiddi'96;  
...; Passera'06]




[Laporta'93;  
Kinoshita,Nio'06;  
Aoyama,Hayakawa,  
Kinoshita,Nio'07'08]



[Aoyama,Hayakawa,Kinoshita,Nio'12]

# Muon $g - 2$



$$0.5 \frac{\alpha}{\pi}$$



$$(-0.328 \dots + 1.094 \dots |e|) \frac{\alpha^2}{\pi^2}$$




$$(1.181 \dots + 22.868 \dots |e|) \frac{\alpha^3}{\pi^3}$$



$$(-1.910 \dots + 132.682 \dots |e|) \frac{\alpha^4}{\pi^4}$$




$$(9.168(571) + 742.18(87)|e|) \frac{\alpha^5}{\pi^5}$$



- $a_\mu(\text{exp}) - a_\mu(\text{SM}) \sim 249(87) \times 10^{-11}$
- $a_\mu(4 \text{ loop QED}) \sim 381 \times 10^{-11}$

[Aoyama, Hayakawa, Kinoshita, Nio '12]



$$0.5 \frac{\alpha}{\pi}$$



$$(-0.328 \dots + 1.094 \dots |e) \frac{\alpha^2}{\pi^2}$$



$$(1.181 \dots + 22.868 \dots |e) \frac{\alpha^3}{\pi^3}$$



$$(-1.910 \dots + 132.682 \dots |e) \frac{\alpha^4}{\pi^4}$$



$$(9.168(571) + 742.18(87)|e) \frac{\alpha^5}{\pi^5}$$



$$0.5 \frac{\alpha}{\pi}$$

$$(-0.328 \dots + 1.094 \dots |_e) \frac{\alpha^2}{\pi^2}$$

$$(1.181 \dots + 22.868 \dots |_e) \frac{\alpha^3}{\pi^3}$$

$$(-1.910 \dots + 132.682 \dots |_e) \frac{\alpha^4}{\pi^4}$$

$$(9.168(571) + 742.18(87)|_e) \frac{\alpha^5}{\pi^5}$$

- $a_\mu(\text{exp}) - a_\mu(\text{SM}) \sim 249(87) \times 10^{-11}$
- $a_\mu(4 \text{ loop QED}) \sim 381 \times 10^{-11}$

[Aoyama, Hayakawa, Kinoshita, Nio '12]

- aim: check 4-loop result
- independent calculation
- different method

# Contributions to muon $g - 2$

- **QED**:  $116\,584\,718.962(9)(19)(7)(78) \times 10^{-11}$  [Aoyama, Hayakawa, Kinoshita, Nio'12]
- **hadronic VP**:  $6949.1(37.2)(21.0) - 984(6)(4) + 12.4(0.1) \times 10^{-11}$   
[Davier, Höcker, Malaescu, Zhang'10; Hagiwara, Liao, Martin, Nomura, Teubner'11; Jegerlehner, Szafron'11;  
Benayoun, David, DelBuono, Jegerlehner'12; ...; Kurz, Marquard, Liu, Steinhauser'14]]
- **electroweak**  $154(2) \times 10^{-11}$  [Czarnecki, Krause, Marciano'96; Knecht, Peris, Perrottet, de Rafale'02;  
Czarnecki, Marciano, Vainshtein'03; Gnendiger, Stöckinger, Stöckinger-Kim'13]
- **hadronic lbl**:  $116(40) \times 10^{-11}$  [Nyffeler'09; Melnikov, Vainshtein'14; Bijmns, Prades'15; ...]

- **QED**:  $116\,584\,718.962(9)(19)(7)(78) \times 10^{-11}$  [Aoyama, Hayakawa, Kinoshita, Nio'12]
- **hadronic VP**:  $6949.1(37.2)(21.0) - 984(6)(4) + 12.4(0.1) \times 10^{-11}$   
[Davier, Höcker, Malaescu, Zhang'10; Hagiwara, Liao, Martin, Nomura, Teubner'11; Jegerlehner, Szafron'11;  
Benayoun, David, DelBuono, Jegerlehner'12; ...; Kurz, Marquard, Liu, Steinhauser'14]]
- **electroweak**  $154(2) \times 10^{-11}$  [Czarnecki, Krause, Marciano'96; Knecht, Peris, Perrottet, de Rafale'02;  
Czarnecki, Marciano, Vainshtein'03; Gnendiger, Stöckinger, Stöckinger-Kim'13]
- **hadronic lbl**:  $116(40) \times 10^{-11}$  [Nyffeler'09; Melnikov, Vainshtein'14; Bijmans, Prades'15; ...]


$g - 2$

NNLO hadronic

$$a_{\mu}^{(1)} = \frac{1}{3} \left( \frac{\alpha}{\pi} \right)^2 \int_{m_{\pi}^2}^{\infty} ds \frac{R(s)}{s} K^{(1)}(s)$$

$$R(s) = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{4\pi\alpha^2/(3s)}$$

$$K^{(1)}(s) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x)\frac{s}{m_{\mu}^2}}$$




LO: exact result [Brodsky,de Rafael'68]


NLO, NNLO: expand for  $s \gg m_{\mu}^2$

compilation of  $R(s)$  from [Nomura,Teubner]


# LO, NLO and NNLO contributions




# LO, NLO and NNLO contributions



# LO, NLO and NNLO contributions




# LO, NLO and NNLO contributions ( $\times 10^{-11}$ )




6949



-209.0



106.8




3.5



8.0





-4.1




9.1



-0.6




0.005




---

LO	6949(37.2)(21.0)
NLO	-98.4(6)(4)
NNLO	12.4(0.1)

---




---

LO lbl	116(40)
NLO lbl	3(2)

---

( $\times 10^{-11}$ )

**LO:** [Davier,Höcker,Malaescu,Zhang'10; Hagiwara,Liao,Martin,Nomura,Teubner'11; Jegerlehner,Szafron'11;

Benayoun,David,DelBuono,Jegerlehner'12]

**NLO:** [Krause'96; Hagiwara,Martin,Nomura,Teubner'03; Hagiwara,Liao,Martin,Nomura,Teubner'11; Greynat, de Rafael'12]

**NNLO:** [Kurz,Marquard,Liu,Steinhauser'14]

**LO lbl:** [Nyffeler'09; Melnikov,Vainshtein'14; Bijmans,Prades'15; ...]


**NLO lbl:** [Colangelo,Hoferichter,Nyffeler,Passera,Stoffer'14]

# $g - 2 - \text{QED}$ Electrons at 4 loops

# QED – electron loops



$$(-1.910 \dots + 132.685 \dots |e) \frac{\alpha^4}{\pi^4}$$




“light-by-light”-type




non-“light-by-light”-type

# QED – electron loops



$$(-1.910 \dots + 132.685 \dots |e) \frac{\alpha^4}{\pi^4}$$



“light-by-light”-type  $\approx 95\%$



non-“light-by-light”-type

# $n_l^2$ and $n_l^3$ terms: 2 and 3 $e^-$ loops



- gauge invariant subset

- results  $\sim c_0 + c_1 \log(m_e/m_\mu) + c_2 \log^2(m_e/m_\mu) + \dots + \frac{m_e}{m_\mu} [\dots]$

- strategy:

$$1.) m_e = 0, m_\mu^{\text{OS}}, \bar{\alpha}(\mu) \Leftrightarrow \log(\mu/m_\mu)$$



$$2.) \bar{\alpha}(\mu) \rightarrow \alpha^{\text{OS}} \Leftrightarrow \log(\mu/m_e)$$

$$\left. \begin{array}{l} 1.) \\ 2.) \end{array} \right\} \Leftrightarrow \log(m_e/m_\mu)$$

- analytic results

[Laporta'93; Aguilar,Greynat,De Rafael'08; ... Lee,Marquard,Smirnov,Smirnov,Steinhauser'13]


# “light-by-light”-type



IV(a)




IV(b)



IV(c)

- $m_e = 0$  not possible  $\Leftrightarrow$  apply methods of region [Beneke, Smirnov'98; Smirnov]

# “light-by-light”-type



- $m_e = 0$  not possible  $\Leftrightarrow$  apply methods of region [Beneke, Smirnov'98; Smirnov]

- example: 3-loop diagram

loop momenta  $k, l, r \sim m_\mu$  (“hard”) or  $k, l, r \sim m_e$  (“soft”)

e.g.:  $k$  soft,  $p^2 = m_\mu^2 \Leftrightarrow \frac{1}{m_\mu^2 - (p-k)^2} = \frac{1}{2pk - k^2} \rightarrow \frac{1}{2pk}$

$$\Leftrightarrow \left\{ \begin{array}{ll} 3\text{-loop OS integrals} & (3 \times \text{hard}) \\ 3\text{-loop “linear” integrals} & (3 \times \text{soft}) \\ 2\text{-loop OS} \times 1\text{-loop tadpole} & (2 \times \text{hard}, 1 \times \text{soft}) \\ 1\text{-loop OS} \times 2\text{-loop tadpole} & (1 \times \text{hard}, 2 \times \text{soft}) \\ 1\text{-loop OS} \times 2\text{-loop “linear”} & (1 \times \text{hard}, 2 \times \text{soft}) \end{array} \right.$$

# “light-by-light”-type

- asymptotic expansion for  $x \equiv m_e/m_\mu \ll 1$

3 loops

$$\begin{aligned} a_\mu = & a_\mu^{(0,0)} + a_\mu^{(0,1)} \ln \frac{m_\mu}{m_e} \\ & + \frac{m_e}{m_\mu} \left( a_\mu^{(1,0)} + a_\mu^{(1,1)} \ln \frac{m_\mu}{m_e} \right) \\ & + \frac{m_e^2}{m_\mu^2} \left( a_\mu^{(2,0)} + a_\mu^{(2,1)} \ln \frac{m_\mu}{m_e} + a_\mu^{(2,2)} \ln^2 \frac{m_\mu}{m_e} \right. \\ & \quad \left. + a_\mu^{(2,3)} \ln^3 \frac{m_\mu}{m_e} \right) \\ & + \dots \end{aligned}$$

$$\begin{array}{lll} \frac{m_e}{m_\mu} \approx 0.005 & \ln \frac{m_\mu}{m_e} \approx 5.332 & \\ \frac{m_e^2}{m_\mu^2} \ln^3 \frac{m_\mu}{m_e} \approx 0.004 & \frac{m_e}{m_\mu} \ln \frac{m_\mu}{m_e} \approx 0.026 & \frac{m_e}{m_\mu} \ln^2 \frac{m_\mu}{m_e} \approx 0.137 \\ & \frac{m_e^2}{m_\mu^2} \ln^4 \frac{m_\mu}{m_e} \approx 0.019 & \end{array}$$

# “light-by-light”-type

- asymptotic expansion for  $x \equiv m_e/m_\mu \ll 1$

3 loops

$$\begin{aligned} a_\mu = & a_\mu^{(0,0)} + a_\mu^{(0,1)} \ln \frac{m_\mu}{m_e} \\ & + \frac{m_e}{m_\mu} \left( a_\mu^{(1,0)} + a_\mu^{(1,1)} \ln \frac{m_\mu}{m_e} \right) \\ & + \frac{m_e^2}{m_\mu^2} \left( a_\mu^{(2,0)} + a_\mu^{(2,1)} \ln \frac{m_\mu}{m_e} + a_\mu^{(2,2)} \ln^2 \frac{m_\mu}{m_e} \right. \\ & \quad \left. + a_\mu^{(2,3)} \ln^3 \frac{m_\mu}{m_e} \right) \\ & + \dots \end{aligned}$$

$$= (20.52801865 + 0.42645778 + 0.00655695 + \dots) \left( \frac{\alpha}{\pi} \right)^3$$

$$\begin{array}{lll} \ln \frac{m_\mu}{m_e} \approx 5.332 & & \\ \frac{m_e}{m_\mu} \approx 0.005 & \frac{m_e}{m_\mu} \ln \frac{m_\mu}{m_e} \approx 0.026 & \frac{m_e}{m_\mu} \ln^2 \frac{m_\mu}{m_e} \approx 0.137 \\ \frac{m_e^2}{m_\mu^2} \ln^3 \frac{m_\mu}{m_e} \approx 0.004 & \frac{m_e^2}{m_\mu^2} \ln^4 \frac{m_\mu}{m_e} \approx 0.019 & \end{array}$$

# “light-by-light”-type

- asymptotic expansion for  $x \equiv m_e/m_\mu \ll 1$


4 loops

$$\begin{aligned} a_\mu = & a_\mu^{(0,0)} + a_\mu^{(0,1)} \ln \frac{m_\mu}{m_e} + a_\mu^{(0,2)} \ln^2 \frac{m_\mu}{m_e} \\ & + \frac{m_e}{m_\mu} \left( a_\mu^{(1,0)} + a_\mu^{(1,1)} \ln \frac{m_\mu}{m_e} + a_\mu^{(1,2)} \ln^2 \frac{m_\mu}{m_e} \right) \\ & + \frac{m_e^2}{m_\mu^2} \left( a_\mu^{(2,0)} + a_\mu^{(2,1)} \ln \frac{m_\mu}{m_e} + a_\mu^{(2,2)} \ln^2 \frac{m_\mu}{m_e} \right. \\ & \quad \left. + a_\mu^{(2,3)} \ln^3 \frac{m_\mu}{m_e} + a_\mu^{(2,4)} \ln^4 \frac{m_\mu}{m_e} \right) \\ & + \dots \end{aligned}$$

$$\begin{aligned} \frac{m_e}{m_\mu} &\approx 0.005 & \ln \frac{m_\mu}{m_e} &\approx 5.332 & \frac{m_e}{m_\mu} \ln^2 \frac{m_\mu}{m_e} &\approx 0.137 \\ \frac{m_e^2}{m_\mu^2} \ln^3 \frac{m_\mu}{m_e} &\approx 0.004 & \frac{m_e}{m_\mu} \ln \frac{m_\mu}{m_e} &\approx 0.026 & & \\ \frac{m_e^2}{m_\mu^2} \ln^4 \frac{m_\mu}{m_e} &\approx 0.019 & & & & \end{aligned}$$

- generation of amplitudes with `qgraf` [Nogueira'91]
- traces, projectors, . . . FORM [Vermaseren,...]
- asymptotic expansion: `asy` [Pak,Smirnov'10] and in-house program  
basic idea: alpha parametrization
- all 3-loop integrals are known analytically  
⇔ all 4-loop CTs known analytically
- each individual Feynman amplitude:  
⇔ analytic linear combination of master integrals
- 4-loop on-shell integrals:  
 $\approx 70 = 40$  ana/high prec. +  $30$  num [Marquard,Smirnov,Smirnov,Steinhauser'15]
- 4-loop "linear" integrals:  
 $\approx 70 = 20$  ana/high prec. +  $50$  num
- numerical integration with FIESTA [Smirnov'13]

# “light-by-light” expansion in $m_e/m_\mu$



$$\begin{aligned} & A_2^{(8),IV(a0)} \\ = & 7.5018 \pm 0.0026 + 14.8808\ell_x + 6.5797\ell_x^2 \\ & + x \left[ 6.29 \pm 0.46 - 14.6216\ell_x + 8.7729\ell_x^2 \right] \\ & + x^2 \left[ -16.81 \pm 0.43 + 30.0172\ell_x - 6.5069\ell_x^2 + 7.6489\ell_x^3 - 0.8889\ell_x^4 \right] \\ & + x^3 \left[ -48.31 \pm 0.24 - 4.8739\ell_x + 13.1595\ell_x^2 \right] \\ = & 7.5018 \pm 0.0026 - 79.3384 + 187.0352 \\ & + [0.0304 \pm 0.0022 + 0.3770 + 1.2061] \\ & + [-0.000393 \pm 0.000010 - 0.003743 - 0.004326 - 0.02711 - 0.01680] \\ & + [-0.0000055 + 0.0000029 + 0.0000423] \\ = & [115.1986 \pm 0.0026] + [1.6135 \pm 0.0022] + [-0.052378 \pm 0.000010] \\ & + [0.000040] \\ = & 116.7598 \pm 0.0034 \end{aligned}$$

$$x = m_e/m_\mu \approx 1/206.7682843$$

$$\ell_x = \log(x) \approx -5.3$$


# “light-by-light” expansion in $m_e/m_\mu$



$$\begin{aligned} & A_2^{(8),IV(b)} \\ = & 27.395 \pm 0.014 + (4.93482 \pm 0.00003)\ell_x \\ & + x [-0.81 \pm 1.22 + 59.0235\ell_x] \\ & + x^2 [142.5 \pm 7.6 + 40.6546\ell_x + 20.5582\ell_x^2 - 9.6167\ell_x^3 + 0.8333\ell_x^4] \\ & + x^3 [62.11 \pm 2.89 + 132.7421\ell_x - 40.9406\ell_x^2] \\ = & 27.395 \pm 0.014 + (-26.3105 \pm 0.0002) \\ & + [-0.0039 \pm 0.0059 - 1.5219] \\ & + [0.003334 \pm 0.0001769 - 0.005070 + 0.01367 + 0.03409 + 0.01575] \\ & + [0.000007 \pm 0. - 0.000080 - 0.000132] \\ = & [1.084 \pm 0.014] + [-1.5259 \pm 0.0059] + [0.06177 \pm 0.00018] + [-0.0002047] \\ = & -0.380 \pm 0.016 \end{aligned}$$

$$x = m_e/m_\mu \approx 1/206.7682843 \quad \ell_x = \log(x) \approx -5.3$$

# “light-by-light” results



	[Kurz,Liu et al.'15]	[Kinoshita,Nio'04; Aoyama et al.'08]
IV(a0) $ee$	$116.76 \pm 0.02$	$116.759183 \pm 0.000292$
IV(a1) $e\mu$	$2.69 \pm 0.14$	$2.697443 \pm 0.000142$
IV(a2) $\mu e$	$4.33 \pm 0.17$	$4.328885 \pm 0.000293$
IV(a)	$123.78 \pm 0.22$	$123.78551 \pm 0.00044$
IV(b)	$-0.38 \pm 0.08$	$-0.4170 \pm 0.0037$
IV(c)	$2.94 \pm 0.30$	$2.9072 \pm 0.0044$

- IV(a0): see also [Calmet,Peterman'75; Chloubert,Samuel'77]:  $117.4 \pm 0.5$
- numerical uncertainty:  $1.2 \times 10^{-11} \ll a_\mu(\text{exp}) - a_\mu(\text{SM}) \sim 249(87) \times 10^{-11}$

# Heavy-lepton contribution



- $\left(\frac{\alpha}{\pi}\right)^4 \frac{m_\mu^2}{m_\tau^2} \approx \left(\frac{\alpha}{\pi}\right)^5$
- analytic asymptotic expansion for  $m_\mu \ll m_\tau$



2 loops: [Elend'66; Passera'04'06]

3 loops: [Samuel,Li'91; Laporta,Remiddi'93; Laporta'93; Czarnecki'98; Kühn'03; Friot,Greynat,De Rafael'05; Passera'04'06]

# Graphical asymptotic expansion



- analytic result

[Kurz,Liu,Marquard,Steinhauser]

$$A_{2,\mu}^{(8)}(m_\mu/m_\tau) = \sum_n c_n \left(\frac{m_\mu}{m_\tau}\right)^n$$

	$10^2 \cdot A_{2,\mu}^{(8)}(m_\mu/m_\tau)$	
	[Kurz,Liu,Marquard,Steinhauser'13]	[Aoyama,Hayakawa,Kinoshita,Nio'12]
I(a)	0.00324281(2)	0.0032(0)
I(b) + I(c) + II(b) + II(c)	-0.6292808(6)	-0.6293(1)
I(d)	0.0367796(4)	0.0368(0)
III	4.5208986(6)	4.504(14)
II(a) + IV(d)	-2.316756(5)	-2.3197(37)
IV(a)	3.851967(3)	3.8513(11)
IV(b)	0.612661(5)	0.6106(31)
IV(c)	-1.83010(1)	-1.823(11)

- $A_{2,\mu}^{(8)}(m_\mu/m_\tau)$   
 $= 0.0421670 + 0.0003257 + 0.0000015 + \mathcal{O}\left(\frac{m_\mu}{m_\tau}\right)^8$   
 $= 0.0424942 + \mathcal{O}\left(\frac{M_\mu}{M_\tau}\right)^8$
- 0.04234(12) [Aoyama,Hayakawa,Kinoshita,Nio'12]

- 4-loop lbl-type contribution  $m_e \neq 0$   
systematic improvement of numerical accuracy possible

- 4-loop  $m_\tau$  contribution

$$10^{11} \times a_\mu \Big|_{\tau \text{ loops}} = 42.13 + 0.45 + 0.12$$

$$10^{11} \times a_\mu \Big|_{\text{univ.}} = 116\,140\,973.21 - 177\,230.51 + 1\,480.42 - 5.56 + 0.06$$

- 2 closed electron loops: known analytically for  $m_e \rightarrow 0$
- hadronic vacuum polarization contribution at NNLO



[Aoyama, Hayakawa, Kinoshita, Nio'12]

group	$A_2^{(8)}(m_\mu/m_e)$	$A_2^{(8)}(m_\mu/m_\tau)$	$A_3^{(8)}$
I(a)	7.74547 (42)	0.000032 (0)	0.003209 (0)
I(b)	7.58201 (71)	0.000252 (0)	0.002611 (0)
I(c)	1.624307 (40)	0.000737 (0)	0.001807 (0)
I(d)	-0.22982 (37)	0.000368 (0)	0
II(a)	-2.77888 (38)	-0.007329 (1)	0
II(b)	-4.55277 (30)	-0.002036 (0)	-0.009008 (1)
II(c)	-9.34180 (83)	-0.005246 (1)	-0.019642 (2)
III	10.7934 (27)	0.04504 (14)	0
IV(a)	123.78551 (44)	0.038513 (11)	0.083739 (36)
IV(b)	-0.4170 (37)	0.006106 (31)	0
IV(c)	2.9072 (44)	-0.01823 (11)	0
IV(d)	-4.43243 (58)	-0.015868 (37)	0

