

A two-Higgs-doublet model facing experimental hints

Andreas Crivellin^{1,a}, Julian Heeck^{2,b}, and Peter Stoffer^{3,c}

¹ CERN Theory Division, CH-1211 Geneva 23, Switzerland

² Service de Physique Théorique, Université Libre de Bruxelles, Boulevard du Triomphe, CP225, 1050 Brussels, Belgium

³ Helmholtz-Institut für Strahlen- und Kernphysik (Theory) and Bethe Center for Theoretical Physics, University of Bonn, D-53115 Bonn, Germany

Abstract. Physics beyond the Standard Model has so far eluded our experimental probes. Nevertheless, a number of interesting anomalies have accumulated that can be taken as hints towards new physics: BaBar, Belle, and LHCb have found deviations of approximately 3.8σ in $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$; the anomalous magnetic moment of the muon differs by about 3σ from the theoretic prediction; the branching ratio for $\tau \rightarrow \mu\nu\nu$ is about 2σ above the Standard Model expectation; and CMS and ATLAS found hints for a non-zero decay rate of $h \rightarrow \mu\tau$ at 2.6σ . Here we consider these processes within a lepton-specific two-Higgs doublet model with additional non-standard Yukawa couplings and show how (and which of) these excesses can be accommodated.

1 Introduction

This talk is based on Ref. [1], where a more detailed discussion can be found. Tests of flavor universality or flavor violation serve as a useful tool to search for physics beyond the Standard Model (SM), seeing as the SM predictions are precisely known. Some experiments have reported on deviations from the SM, which we list below.

- Lepton universality in semileptonic B decays, parametrized by the ratio

$$\mathcal{R}(D^{(*)}) \equiv \frac{\text{BR}(B \rightarrow D^{(*)}\tau\nu)}{\text{BR}(B \rightarrow D^{(*)}\ell\nu)}, \quad \ell = e, \mu, \quad (1)$$

has been studied by BaBar, Belle and LHCb. Combining their experimental values yields

$$\mathcal{R}(D)_{\text{exp}} = 0.388 \pm 0.047, \quad \mathcal{R}(D^*)_{\text{exp}} = 0.321 \pm 0.021.$$

Together, these values deviate by more than 3σ from the SM prediction [2–4]

$$\mathcal{R}(D)_{\text{SM}} = 0.297 \pm 0.017, \quad \mathcal{R}(D^*)_{\text{SM}} = 0.252 \pm 0.003.$$

One possible new-physics explanation comes in the form of a charged scalar [5–7].

- The muon’s anomalous magnetic moment $a_\mu \equiv (g - 2)_\mu/2$, as measured by the Brookhaven experiment E821 [8], deviates by 2.7σ from its SM value,

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (236 \pm 87) \times 10^{-11}. \quad (2)$$

It is not yet clear if this discrepancy is due to new physics or rather underestimated hadronic uncertainties; there are ongoing efforts to reduce the model dependence in the hadronic light-by-light estimate based on dispersion relations [9–12] or lattice QCD [13–16]. A possible interpretation in terms of new physics has been found in two-Higgs-doublet models (2HDM) [17, 18], in particular the lepton-specific 2HDM [19–22].

- For $\tau \rightarrow \mu\nu\nu$, the dominant uncertainty in the SM prediction for the branching ratio comes from the τ lifetime. Using the PDG [23] values for τ lifetime and branching ratios

$$\begin{aligned} B_\mu &\equiv \text{BR}(\tau \rightarrow \mu\bar{\nu}\nu)_{\text{exp}} = (17.41 \pm 0.04)\%, \\ B_e &\equiv \text{BR}(\tau \rightarrow e\bar{\nu}\nu)_{\text{exp}} = (17.83 \pm 0.04)\%, \end{aligned} \quad (3)$$

we can determine the deviations from the SM prediction $\Delta_\ell \equiv B_\ell/B_\ell^{\text{SM}} - 1$ [24] as

$$\Delta_\mu^{\text{PDG}} = (0.69 \pm 0.29)\%, \quad \Delta_e = (0.28 \pm 0.28)\%. \quad (4)$$

There is a deviation of about 2.4σ for the muon final state, whereas the electron channel is compatible with the SM prediction, hinting at lepton non-universality. Charged scalars of a 2HDM will modify the rate [24–26].

- Going from lepton non-universality to outright lepton flavor violation, we are drawn to the recent CMS excess of 2.4σ in $h \rightarrow \mu\tau$ [27]:

$$\text{BR}(h \rightarrow \mu\tau) = (0.84^{+0.39}_{-0.37})\%. \quad (5)$$

A similar search at ATLAS [28] finds $\text{BR}(h \rightarrow \mu\tau) = (0.77 \pm 0.62)\%$, which slightly bumps the significance to about 2.6σ . Possible explanations naturally require an extended scalar sector [18, 29–35].

^ae-mail: andreas.crivellin@cern.ch

^be-mail: julian.heeck@ulb.ac.be

^ce-mail: stoffer@hiskp.uni-bonn.de

Each anomaly individually can be accommodated in SM extensions by scalars, e.g. 2HDMs. The goal of our study is to see if all four anomalies can be explained simultaneously with a fairly minimal model.

2 Modified 2HDM-X

We will study a lepton-specific 2HDM (2HDM-X), defined by the Yukawa couplings in the Lagrangian

$$\mathcal{L}_Y = -\bar{Q}_L Y^u \tilde{\Phi}_2 u_R - \bar{Q}_L Y^d \Phi_2 d_R - \bar{L}_L Y^\ell \Phi_1 e_R + \text{h.c.}, \quad (6)$$

with additional couplings that break the type-X structure

$$\Delta \mathcal{L}_Y = -\bar{Q}_L \xi^u \tilde{\Phi}_1 u_R - \bar{Q}_L \xi^d \Phi_1 d_R - \bar{L}_L \xi^\ell \Phi_2 e_R + \text{h.c.} \quad (7)$$

The scalar interactions with fermions after electroweak breaking can be written as

$$\begin{aligned} \mathcal{L} \supset & \bar{\nu}_i \Gamma_{\nu_i \ell_j}^{H^+ LR} P_R \ell_j H^+ \\ & + \bar{u}_i \left(\Gamma_{u_i d_j}^{H^+ RL} P_L + \Gamma_{u_i d_j}^{H^+ LR} P_R \right) d_j H^+ \\ & + \sum_{H_k^0 = h, A, H} \sum_{f = u, d, \ell} \left(\bar{f}_i \Gamma_{f_i f_j}^{H_k^0 LR} P_R f_j H_k^0 \right) + \text{h.c.}, \quad (8) \end{aligned}$$

where the couplings – in the limit of large $\tan\beta$ of interest here – are given by

$$\Gamma_{q_i q_j}^{hLR} \simeq -\frac{1}{\sqrt{2}} \left(\frac{m_{q_i}}{v} \delta_{ij} \cos \alpha - \epsilon_{ij}^q \sin \alpha \right), \quad (9)$$

$$\Gamma_{q_i q_j}^{H^+ LR} \simeq -\frac{1}{\sqrt{2}} \left(\frac{m_{q_i}}{v} \delta_{ij} \sin \alpha + \epsilon_{ij}^q \cos \alpha \right), \quad (10)$$

$$\Gamma_{d_i d_j}^{ALR} \simeq -i \frac{1}{\sqrt{2}} \epsilon_{ij}^d, \quad (11)$$

$$\Gamma_{u_i u_j}^{ALR} \simeq i \frac{1}{\sqrt{2}} \epsilon_{ij}^u, \quad (12)$$

$$\Gamma_{u_i d_j}^{H^+ LR} \simeq V_{ij} \epsilon_{ij}^d, \quad (13)$$

$$\Gamma_{u_i d_j}^{H^+ RL} \simeq -\epsilon_{ij}^{u*} V_{ij}^*, \quad (14)$$

$$\Gamma_{\ell_f \ell_i}^{hLR} \simeq \frac{\sin \alpha \tan \beta}{\sqrt{2}} \left(\frac{m_{\ell_i}}{v} \delta_{fi} - \epsilon_{fi}^\ell \right), \quad (15)$$

$$\Gamma_{\ell_f \ell_i}^{H^+ LR} \simeq -\frac{\cos \alpha \tan \beta}{\sqrt{2}} \left(\frac{m_{\ell_i}}{v} \delta_{fi} - \epsilon_{fi}^\ell \right), \quad (16)$$

$$\Gamma_{\ell_f \ell_i}^{ALR} \simeq -i \frac{\tan \beta}{\sqrt{2}} \left(\frac{m_{\ell_i}}{v} \delta_{fi} - \epsilon_{fi}^\ell \right), \quad (17)$$

$$\Gamma_{\nu_f \ell_i}^{H^+ LR} \simeq \tan \beta \left(\frac{m_{\ell_i}}{v} \delta_{fi} - \epsilon_{fi}^\ell \right). \quad (18)$$

V denotes the Cabibbo–Kobayashi–Maskawa mixing matrix, $v \simeq 174$ GeV the vacuum expectation value, and the ϵ matrices parametrize the deviation from the type-X structure (ϵ^f and ξ^f are related by fermion rotations). On phenomenological grounds, we take $\epsilon^d = 0$ and both ϵ^u and ϵ^ℓ of the form

$$\epsilon^{u,\ell} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \times & \times \end{pmatrix}, \quad (19)$$

where \times denotes a non-zero entry, allowing for lepton flavor violation in the μ – τ sector.




Figure 1. Allowed regions for z and δg from Δ_μ (yellow), Δ_e (blue), and the Michel parameter η (red); see text for definitions. Darker (lighter) regions are at the 2σ (3σ) level, using PDG values [1].

3 Phenomenology

In this section we address the constraints on our model and its potential to resolve the experimental anomalies outlined above.

3.1 Tau decays $\tau \rightarrow \ell \nu \nu$

Decays of the tau are modified at tree-level by the charged scalar H^+ (including a change in the Michel parameter η) [24–26] and at loop-level through a modified $W\tau\nu$ coupling $g_{W\tau\nu} \rightarrow g_{W\tau\nu}(1 + \delta g)$ [21, 24, 25]:

$$\Delta_e \simeq 2\delta g, \quad (20)$$

$$\Delta_\mu \simeq 2\delta g + \frac{z^2}{4} - 2z \frac{m_\mu}{m_\tau}, \quad (21)$$

$$\eta = -\frac{2z}{4 + z^2}, \quad (22)$$

with the 2HDM couplings [21]

$$z \equiv \frac{v^2}{m_{H^+}^2} \Gamma_{\nu_f \tau}^{LRH^+} \Gamma_{\nu_\mu}^{LRH^+*}, \quad (23)$$

$$\delta g = \frac{\tan^2 \beta}{32\pi^2} \left| \frac{m_\tau}{v} - \epsilon_{33}^\ell \right|^2 \left[F\left(\frac{m_A^2}{m_{H^+}^2}\right) + F\left(\frac{m_H^2}{m_{H^+}^2}\right) \right]. \quad (24)$$

The experimentally allowed regions are shown in Fig. 1. In the SM we have $\delta g = z = 0$, which is disfavored by $\tau \rightarrow \mu\nu\nu$ at more than 2σ (with PDG values). In the 2HDM-X one has $\delta g \leq 0$ and $z > 0$, which makes Δ_μ even worse and puts pressure on the 2HDM-X and 2HDM-II (which has the same lepton couplings). In our modified 2HDM-X, we can however flip the sign of the tau coupling via $\epsilon_{33}^\ell > m_\tau/v$ and obtain $z < 0$, alleviating the tension in $\tau \rightarrow \mu\nu\nu$ decays.




Figure 2. Allowed regions in the m_A - m_H plane from $(g-2)_\mu$, $\tau \rightarrow \mu\gamma$ and $h \rightarrow \mu\tau$ for $\tan(\beta) = 50$ and $\epsilon_{33}^\ell = 2m_\tau/v$. For $h \rightarrow \mu\tau$ blue corresponds to $\cos(\alpha - \beta) = 0.1$ and light blue to $\cos(\alpha - \beta) = 0.2$. The allowed region for Δa_μ is maximal in the sense that we have allowed for the three possibilities $\epsilon_{32}^\ell \neq 0$, $\epsilon_{32}^\ell = \epsilon_{23}^\ell \neq 0$ and $\epsilon_{32}^\ell = -\epsilon_{23}^\ell \neq 0$ as the latter ones can give m_τ/m_μ enhanced one-loop contributions [1].

3.2 Magnetic moment a_μ and $h \rightarrow \mu\tau$

It has been shown that a light pseudoscalar in the 2HDM-X with large $\tan\beta$ can resolve the Δa_μ anomaly via its contribution in a Barr-Zee diagram [36], see for example Refs. [19–21]. As seen above, we need to flip the sign of the tau coupling to the new scalars in order to alleviate the $\tau \rightarrow \mu\nu\nu$ discrepancy. Because of this, it is the non-SM-like CP-even scalar that can resolve the Δa_μ anomaly in our model (and not the pseudoscalar). The pseudoscalar then needs to be heavier than the scalar in order not to cancel the contribution to Δa_μ (see Fig 2).

The same Barr-Zee diagrams that give the desired Δa_μ also lead to $\tau \rightarrow \mu\gamma$ decays in case $\epsilon_{32}^\ell \neq 0$ (as required for $h \rightarrow \mu\tau$). If we want to explain $h \rightarrow \mu\tau$, the $\tau \rightarrow \mu\gamma$ rate needs to be tuned to small values using $m_H \simeq m_A$, which necessarily also suppresses the contribution to Δa_μ (see Fig 2). As a result, it is possible within our modified 2HDM-X to resolve the anomalies in $\tau \rightarrow \mu\nu\nu$ and Δa_μ or to resolve $\tau \rightarrow \mu\nu\nu$ and $h \rightarrow \mu\tau$, but not both at the same time, at least without introducing more parameters.

3.3 Taonic B decays

The relevant effective Hamiltonian for the semileptonic B decays in our model is

$$\mathcal{H} = C_{SM}^{cb} \bar{c}\gamma_\mu P_L b \bar{\tau}\gamma_\mu P_L \nu_\tau + \sum_{X=L,R} C_X^{cb} \bar{c}P_X b \bar{\tau}P_L \nu_\tau, \quad (25)$$




Figure 3. Allowed regions in the C_L^{cb} - C_R^{cb} plane from $\mathcal{R}(D)$ (blue) and $\mathcal{R}(D^*)$ (yellow) for real values of $C_{L,R}^{cb}$. The lighter regions correspond to 2σ experimental uncertainties while the darker regions correspond to 1σ [1].

with SM Wilson coefficient $C_{SM}^{cb} = 4G_F V_{cb}/\sqrt{2}$ and the charged-scalar contribution

$$C_L^{cb} \simeq \frac{\tan\beta}{m_{H^\pm}^2} \epsilon_{32}^{u*} \left(\frac{m_\tau}{v} - \epsilon_{33}^* \right), \quad C_R^{cb} \simeq 0. \quad (26)$$

These Wilson coefficients affect the two ratios $\mathcal{R}(D^{(*)})$ in the following way [4, 37, 38],

$$\frac{\mathcal{R}(D)}{\mathcal{R}(D)_{SM}} = 1 + 1.5 \Re \left[\frac{C_R^{cb} + C_L^{cb}}{C_{SM}^{cb}} \right] + 1.0 \left| \frac{C_R^{cb} + C_L^{cb}}{C_{SM}^{cb}} \right|^2, \quad (27)$$

$$\frac{\mathcal{R}(D^*)}{\mathcal{R}(D^*)_{SM}} = 1 + 0.12 \Re \left[\frac{C_R^{cb} - C_L^{cb}}{C_{SM}^{cb}} \right] + 0.05 \left| \frac{C_R^{cb} - C_L^{cb}}{C_{SM}^{cb}} \right|^2, \quad (28)$$

leading to the allowed regions of Fig. 3. As can be seen, our new-physics model from Eq. (26) has the right structure to easily resolve the anomaly, e.g. with real $C_L^{cb} \simeq -1.2 |C_{SM}^{cb}|$.¹

4 Conclusions

We addressed the measured anomalies in $\mathcal{R}(D^{(*)})$ (3.8σ), a_μ ($\sim 3\sigma$), $\tau \rightarrow \mu\nu\nu$ (2.4σ), and $h \rightarrow \mu\tau$ (2.6σ) within a simple two-Higgs-doublet model. The Yukawa structure of our model is close to the lepton-specific 2HDM

¹Efficiency corrections to $\mathcal{R}(D)$ due to the BaBar detector [2] are important in the case of large contributions from $C_{R,L}^{cb}$, i.e. if one wants to explain $\mathcal{R}(D)$ with destructive interference with the SM contribution. As shown in Ref. [39], these corrections can be effectively taken into account by multiplying the quadratic term in $C_{R,L}^{cb}$ of Eq. (27) by an approximate factor of 1.5 (not included in Eq. (27)).




Figure 4. Allowed regions in the $\tan\beta$ - $v/m_\tau\epsilon_{33}^\ell$ plane from $\mathcal{R}(D^{(*)})$ and $\tau \rightarrow \mu\nu\nu$ at the 2σ level. The yellow region is allowed by $\tau \rightarrow \mu\nu\nu$ using the HFAG result for $m_H = 30$ GeV and $m_A = 200$ GeV, while the (darker) blue one is the allowed region using the PDG result. The red, orange, green, and magenta bands correspond to the allowed regions by $\mathcal{R}(D^{(*)})$ for different values of ϵ_{32}^u . The gray region is excluded by $Z \rightarrow \tau\tau$ and $\tau \rightarrow e\nu\nu$. For $m_H \simeq m_A$, the allowed regions from $\tau \rightarrow \mu\nu\nu$ would be slightly larger [1].

(type X), but with some additional Yukawa couplings involving third-generation fermions that give rise to the b - c (necessary for $\mathcal{R}(D^{(*)})$) and μ - τ transitions (relevant for $h \rightarrow \mu\tau$) as well as corrections to $\tau\tau$ couplings (important for $\tau \rightarrow \mu\nu\nu$).

The charged scalar H^+ influences $\mathcal{R}(D^{(*)})$ and $\tau \rightarrow \mu\nu\nu$ and can resolve both deviations simultaneously (see Fig. 4), as long as the sign of the tau coupling is flipped, meaning $\epsilon_{33}^\ell > m_\tau/v$. Because of this, a light scalar (not pseudoscalar) is needed to resolve the Δa_μ anomaly, which then induces the top decay $t \rightarrow Hc$ [1]. $h \rightarrow \mu\tau$ can only be explained if we give up on Δa_μ , otherwise the rate $\tau \rightarrow \mu\gamma$ would be too large.

It remains to be seen which of these anomalies stand the test of time and which are simply statistical fluctuations. The fact that some of them can be explained rather naturally within a fairly minimal model gives however hope that we are on the verge of something interesting.

Acknowledgements

We thank the organizers of the FCCP2015 for an interesting workshop. A. Crivellin is supported by a Marie Curie Intra-European Fellowship of the European Community's 7th Framework Programme under contract number PIEF-GA-2012-326948. The work of J. Heeck is funded in part by IISN and by Belgian Science Policy (IAP VII/37). P. Stoffer gratefully acknowledges financial support by the DFG (CRC 16, "Subnuclear Structure of Matter").

References

- [1] A. Crivellin, J. Heeck, P. Stoffer (2015), 1507.07567
- [2] J. Lees et al. (BABAR Collaboration), Phys.Rev.Lett. **109**, 101802 (2012), 1205.5442
- [3] J.F. Kamenik, F. Mescia, Phys.Rev. **D78**, 014003 (2008), 0802.3790
- [4] S. Fajfer, J.F. Kamenik, I. Nisandzic, Phys.Rev. **D85**, 094025 (2012), 1203.2654
- [5] A. Crivellin, C. Greub, A. Kokulu, Phys.Rev. **D86**, 054014 (2012), 1206.2634
- [6] A. Celis, M. Jung, X.Q. Li, A. Pich, JHEP **1301**, 054 (2013), 1210.8443
- [7] A. Crivellin, A. Kokulu, C. Greub, Phys.Rev. **D87**, 094031 (2013), 1303.5877
- [8] G. Bennett et al. (Muon $(g-2)$ Collaboration), Phys.Rev. **D73**, 072003 (2006), hep-ex/0602035
- [9] G. Colangelo, M. Hoferichter, M. Procura, P. Stoffer, JHEP **1409**, 091 (2014), 1402.7081
- [10] G. Colangelo, M. Hoferichter, B. Kubis, M. Procura, P. Stoffer, Phys. Lett. **B738**, 6 (2014), 1408.2517
- [11] V. Pauk, M. Vanderhaeghen, Phys. Rev. **D90**, 113012 (2014), 1409.0819
- [12] G. Colangelo, M. Hoferichter, M. Procura, P. Stoffer, JHEP **09**, 074 (2015), 1506.01386
- [13] M. Hayakawa, T. Blum, T. Izubuchi, N. Yamada, PoS **LAT2005**, 353 (2006), hep-lat/0509016
- [14] T. Blum, M. Hayakawa, T. Izubuchi, PoS **LAT-TICE2012**, 022 (2012), 1301.2607
- [15] T. Blum, S. Chowdhury, M. Hayakawa, T. Izubuchi, Phys.Rev.Lett. **114**, 012001 (2015), 1407.2923
- [16] J. Green, O. Gryniuk, G. von Hippel, H.B. Meyer, V. Pascalutsa (2015), 1507.01577
- [17] E.O. Iltan, H. Sundu, Acta Phys. Slov. **53**, 17 (2003), hep-ph/0103105
- [18] Y. Omura, E. Senaha, K. Tobe, JHEP **1505**, 028 (2015), 1502.07824
- [19] A. Broggio, E.J. Chun, M. Passera, K.M. Patel, S.K. Vempati, JHEP **1411**, 058 (2014), 1409.3199
- [20] L. Wang, X.F. Han, JHEP **1505**, 039 (2015), 1412.4874
- [21] T. Abe, R. Sato, K. Yagyu, JHEP **07**, 064 (2015), 1504.07059
- [22] E.J. Chun, Z. Kang, M. Takeuchi, Y.L.S. Tsai (2015), 1507.08067
- [23] K. Olive et al. (Particle Data Group), Chin.Phys. **C38**, 090001 (2014)
- [24] M. Krawczyk, D. Temes, Eur.Phys.J. **C44**, 435 (2005), hep-ph/0410248
- [25] W. Hollik, T. Sack, Phys.Lett. **B284**, 427 (1992)
- [26] M. Aoki, S. Kanemura, K. Tsumura, K. Yagyu, Phys.Rev. **D80**, 015017 (2009), 0902.4665
- [27] V. Khachatryan et al. (CMS), Phys. Lett. **B749**, 337 (2015), 1502.07400
- [28] G. Aad et al. (ATLAS) (2015), 1508.03372

- [29] R. Harnik, J. Kopp, J. Zupan, *JHEP* **03**, 026 (2013), 1209.1397
- [30] M.D. Campos, A.E.C. Hernández, H. Päs, E. Schumacher, *Phys.Rev.* **D91**, 116011 (2015), 1408.1652
- [31] D. Aristizabal Sierra, A. Vicente, *Phys.Rev.* **D90**, 115004 (2014), 1409.7690
- [32] J. Heeck, M. Holthausen, W. Rodejohann, Y. Shimizu, *Nucl.Phys.* **B896**, 281 (2015), 1412.3671
- [33] A. Crivellin, G. D’Ambrosio, J. Heeck, *Phys.Rev.Lett.* **114**, 151801 (2015), 1501.00993
- [34] I. Dorsner, S. Fajfer, A. Greljo, J.F. Kamenik, N. Kosnik et al., *JHEP* **1506**, 108 (2015), 1502.07784
- [35] A. Crivellin, G. D’Ambrosio, J. Heeck, *Phys.Rev.* **D91**, 075006 (2015), 1503.03477
- [36] S.M. Barr, A. Zee, *Phys. Rev. Lett.* **65**, 21 (1990), [Erratum: *Phys. Rev. Lett.*65,2920(1990)]
- [37] A. Akeroyd, S. Recksiegel, *J.Phys.G* **G29**, 2311 (2003), hep-ph/0306037
- [38] Y. Sakaki, H. Tanaka, *Phys.Rev.* **D87**, 054002 (2013), 1205.4908
- [39] S. Fajfer, J.F. Kamenik, I. Nisandzic, J. Zupan, *Phys.Rev.Lett.* **109**, 161801 (2012), 1206.1872