



Results from the search for neutrinoless double beta decay of ^{130}Te with CUORE-0

Lucia Canonica
INFN-LNGS

for the CUORE Collaboration

LNGS, April 9th 2015



The CUORE collaboration





The CUORE Collaborators

C. Alduino¹, K. Alfonso², D. R. Artusa^{1,3}, F. T. Avignone III¹, O. Azzolini⁴, M. Balata³, T. I. Banks^{5,6}, G. Bari⁷, J.W. Beeman⁸, F. Bellini^{9,10}, A. Bersani¹¹, M. Biassoni^{12,13}, C. Brofferio^{12,13}, C. Bucci³, A. Camacho⁴, A. Caminata¹¹, L. Canonica³, X. G. Cao¹⁴, S. Capelli^{12,13}, L. Cappelli^{3,15}, L. Carbone¹³, L. Cardani^{9,10}, N. Casali^{3,16}, L. Cassina^{12,13}, D. Chiesa^{12,13}, N. Chott¹, M. Clemenza^{12,13}, S. Copello¹⁷, C. Cosmelli^{9,10}, O. Cremonesi¹³, R. J. Creswick¹, J. S. Cushman¹⁸, I. Dafinei¹⁰, A. Dally¹⁹, S. Dell'Oro^{3,20}, M. M. Deninno⁷, S. Di Domizio^{17,11}, M. L. Di Vacri^{3,16}, A. Drobizhev^{5,6}, L. Ejzak¹⁹, D. Q. Fang¹⁴, M. Faverzani^{12,13}, G. Fernandes^{17,11}, E. Ferri^{12,13}, F. Ferroni^{9,10}, E. Fiorini^{13,12}, M. A. Franceschi²¹, S. J. Freedman^{6,5}, B. K. Fujikawa⁶, A. Giachero^{12,13}, L. Gironi^{12,13}, A. Giuliani²², P. Gorla³, C. Gotti^{12,13}, T. D. Gutierrez²³, E. E. Haller^{8,24}, K. Han^{18,6}, E. Hansen^{25,2}, K. M. Heeger¹⁸, R. Hennings-Yeomans^{5,6}, K. P. Hickerson², H. Z. Huang², R. Kadel²⁶, G. Keppel⁴, Yu. G. Kolomensky^{5,26}, C. Ligi²¹, K. E. Lim¹⁸, X. Liu², Y. G. Ma¹⁴, M. Maino^{12,13}, M. Martinez²⁷, R. H. Maruyama¹⁸, Y. Mei⁶, N. Moggi^{28,7}, S. Morganti¹⁰, T. Napolitano²¹, S. Nisi³, C. Nones²⁹, E. B. Norman^{30,31}, A. Nucciotti^{12,13}, T. O'Donnell^{5,6}, F. Orio¹⁰, D. Orlandi³, J. L. Ouellet^{5,6}, C. E. Pagliarone^{3,15}, M. Pallavicini^{17,11}, V. Palmieri⁴, L. Pattavina³, M. Pavan^{12,13}, G. Pessina¹³, V. Pettinacci¹⁰, G. Piperno^{9,10}, C. Pira⁴, S. Pirro³, S. Pozzi^{12,13}, E. Previtali¹³, C. Rosenfeld¹, C. Rusconi¹³, E. Sala^{12,13}, S. Sangiorgio³⁰, D. Santone^{3,16}, N. D. Scielzo³⁰, M. Sisti^{12,13}, A. R. Smith⁶, L. Taffarello³², M. Tenconi²², F. Terranova^{12,13}, C. Tomei¹⁰, S. Trentalange², G. Ventura^{33,34}, M. Vignati¹⁰, S. L. Wagaarachchi^{5,6}, B. S. Wang^{30,31}, H. W. Wang¹⁴, L. Wielgus¹⁹, J. Wilson¹, L. A. Winslow²⁵, T. Wise^{18,19}, A. Woodcraft³⁵, L. Zanotti^{12,13}, C. Zarra³, G. Q. Zhang¹⁴, B. X. Zhu², S. Zucchelli^{36,7}

- 19 groups
 - 8 Italy
 - 6 USA
 - 5 associate groups

- 157 collaborators
 - 120 researchers/authors
 - Italy: 71
 - USA: 38
 - Associated Institutions: 11

¹Department of Physics and Astronomy, University of South Carolina, Columbia, SC 29208 - USA

²Department of Physics and Astronomy, University of California, Los Angeles, CA 90095 - USA

³INFN - Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila) I-67010 - Italy

⁴INFN - Laboratori Nazionali di Legnaro, Legnaro (Padova) I-35020 - Italy

⁵Department of Physics, University of California, Berkeley, CA 94720 - USA

⁶Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720 - USA

⁷INFN - Sezione di Bologna, Bologna I-40127 - Italy

⁸Materials Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720 - USA

⁹Dipartimento di Fisica, Sapienza Università di Roma, Roma I-00185 - Italy

¹⁰INFN - Sezione di Roma, Roma I-00185 - Italy

¹¹INFN - Sezione di Genova, Genova I-16146 - Italy

¹²Dipartimento di Fisica, Università di Milano-Bicocca, Milano I-20126 - Italy

¹³INFN - Sezione di Milano Bicocca, Milano I-20126 - Italy

¹⁴Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800 - China

¹⁵Dipartimento di Ingegneria Civile e Meccanica, Università degli Studi di Cassino e del Lazio Meridionale, Cassino I-03043 - Italy

¹⁶Dipartimento di Scienze Fisiche e Chimiche, Università dell'Aquila, L'Aquila I-67100 - Italy

¹⁷Dipartimento di Fisica, Università di Genova, Genova I-16146 - Italy

¹⁸Department of Physics, Yale University, New Haven, CT 06520 - USA

¹⁹Department of Physics, University of Wisconsin, Madison, WI 53706 - USA

²⁰INFN - Gran Sasso Science Institute, L'Aquila I-67100 - Italy

²¹INFN - Laboratori Nazionali di Frascati, Frascati (Roma) I-00044 - Italy

²²Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, 91405 Orsay Campus - France

²³Physics Department, California Polytechnic State University, San Luis Obispo, CA 93407 - USA

²⁴Department of Materials Science and Engineering, University of California, Berkeley, CA 94720 - USA

²⁵Massachusetts Institute of Technology, Cambridge, MA 02139 - USA

²⁶Physics Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720 - USA

²⁷Laboratorio de Física Nuclear y Astroparticulas, Universidad de Zaragoza, Zaragoza 50009 - Spain

²⁸Dipartimento di Scienze per la Qualità della Vita, Alma Mater Studiorum - Università di Bologna, Bologna I-47921 - Italy

²⁹Service de Physique des Particules, CEA / Saclay, 91191 Gif-sur-Yvette - France

³⁰Lawrence Livermore National Laboratory, Livermore, CA 94550 - USA

³¹Department of Nuclear Engineering, University of California, Berkeley, CA 94720 - USA

³²INFN - Sezione di Padova, Padova I-35131 - Italy

³³Dipartimento di Fisica, Università di Firenze, Firenze I-50125 - Italy

³⁴INFN - Sezione di Firenze, Firenze I-50125 - Italy

³⁵SUPA, Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ - UK

³⁶Dipartimento di Fisica e Astronomia, Alma Mater Studiorum - Università di Bologna, Bologna I-40127 - Italy



Outline

- Double beta decay physics
- Thermal detectors
- History of ^{130}Te double beta decay experiments
- CUORE-0 results
 - Detector performance
 - Neutrinoless double beta decay analysis



Outline


- **Double beta decay physics**
- Thermal detectors
- History of ^{130}Te double beta decay experiments
- CUORE-0 results
 - Detector performance
 - Neutrinoless double beta decay analysis



Double beta decay

Double beta decay is a very rare nuclear decay $(N, Z) \rightarrow (N-2, Z+2)$


$$(A, Z) \rightarrow (A, Z + 2) + 2e^- + 2\bar{\nu}$$



2νDBD:

- 2nd order process allowed in SM
- observed in several nuclei
- $\tau \sim 10^{19-21} \text{ y}$


$$(A, Z) \rightarrow (A, Z + 2) + 2e^-$$



0νDBD:

- Lepton number violation $\Delta L=2$
- exists for Majorana neutrinos
- $\nu = \bar{\nu}$
- $\tau > 10^{24-25} \text{ y}$

$\beta\beta$ summed e^- energy spectrum





$0\nu\text{DBD}$ and neutrino mass

$$T_{1/2}^{0\nu} = \frac{m_e^2}{G_{0\nu} \cdot M_{nucl}^2 \cdot m_{\beta\beta}^2}$$


$G_{0\nu}$: Phase space integral $\sim Q^5$

M_{nucl} : Nuclear Matrix Elements

$m_{\beta\beta}$: effective neutrino mass

$$m_{\beta\beta} = \left| \sum_i m_{\nu_i} U_{ei}^2 \right|$$

- The observation of $0\nu\text{DBD}$:
 - proof of the Majorana nature of neutrino
 - constraints on neutrino mass hierarchy and scale




IBM-2 Phys. Rev. C 91, 034304 (2015)
 QRPA-TU Phys. Rev. C 87, 045501 (2013)
 pnQRPA Phys. Rev. C 91, 024613 (2015)
 ISM Nucl. Phys. A 818, 139 (2009)
 EDF Phys. Rev. Lett. 105, 252503 (2010)



Sensitivity

- Half-life corresponding to the minimum number of detectable signal events above background at a given C.L.






Outline

- Double beta decay physics
- **Thermal detectors**
- History of ^{130}Te double beta decay experiments
- CUORE-0 results
 - Detector performance
 - Neutrinoless double beta decay analysis


Thermal detectors

The working principle is very simple:



The energy deposited by a particle interaction in the absorber is converted to a measurable temperature variation.




- wide choice of detector materials
- source embedded in the detector
- excellent energy resolution






^{130}Te for $0\nu\text{DBD}$

- Q-value (2528 keV)
- Highest natural isotopic abundance (~34%)
- Favourable calculation of NME








The CUORE bolometers



Sensitive
thermometers:
NTD thermistor



- natTeO₂ crystals
- NTD-Ge thermistor ($R \sim 50 M\Omega$)
$$R(T) = R_0 \exp \left[\frac{T_0}{T} \right]^{1/2}$$
- Resolution @0vDBD energy (2528 keV):
 $\Delta E = 5-7 \text{ keV FWHM}$






Outline


- Double beta decay physics
- Thermal detectors
- **History of ^{130}Te double beta decay experiments**
- CUORE-0 results
 - Detector performance
 - Neutrinoless double beta decay analysis

TeO₂ arrays





The CUORE challenge



- 19 towers
- 13 floors
- 4 crystals

- Tightly packed array of 988 bolometric detectors.
- $M = 741$ kg of TeO_2 (206 kg ^{130}Te) to look for $0\nu\text{DBD}$ of ^{130}Te .

Energy resolution @ ROI: 5 keV

Background goal: 0.01 c/(keV kg y)

Sensitivity 90% C.L. (5 y):

$$T_{1/2} = 9.5 \times 10^{25} \text{ y}$$

$$m_{\beta\beta} = 50\text{-}130 \text{ meV}$$

IBM-2 Phys. Rev. C 91, 034304 (2015)
QRPA-TU Phys. Rev. C 87, 045501 (2013)
pnQRPA Phys. Rev. C 91, 024613 (2015)
ISM Nucl. Phys. A 818, 139 (2009)
EDF Phys. Rev. Lett. 105, 252503 (2010)




Cuoricino

- 62 TeO₂ crystals, for a total mass of 40.7 kg.
- 19.75 kg · yr of ¹³⁰Te
- Run between 2003 and 2008


$$T_{1/2} > 2.8 \cdot 10^{24} y$$
$$m_{\beta\beta} < 0.3 \div 0.7 eV$$

Astropart. Phys. (2011),
doi:10.1016/j.astropartphys.2011.02.002

Cuoricino final energy spectrum




The Cuoricino background



Background @ Q-value
 $0\nu\text{DBD}$: 0.15 c/(keV · kg · y)

From MC simulations:

- ^{232}Th contaminations in cryostat shield: (~30%)
- Degraded alphas from crystal surfaces (~10%)
- Degraded alphas from Cu holders surfaces (~60%)







From Cuoricino to CUORE

- **Background suppression**

- New (lighter) detector design structure
- Reduced overall detector surfaces by a factor ~ 2
- New surface cleaning technique
- Strict production protocols for TeO_2 surface contamination
- Minimization of Rn exposure (Glove Box assembly)
- Strict material selection (e.g. raw materials)






CUORE-0

CUORE-0 is the **first tower** produced out of the CUORE assembly line.

- 52 TeO_2 $5 \times 5 \times 5 \text{ cm}^3$ crystals ($\sim 750 \text{ g}$ each)
- 13 floors of 4 crystals each
- total detector mass: 39 kg TeO_2 (10.9 kg of ^{130}Te)


CUORE-0 has been taking data since March 2013 in the 25 year old Cuoricino cryostat.

- **Proof of concept** of CUORE detector in all stages
- Test and debug of the CUORE tower assembly line
- Test of the CUORE **DAQ and analysis framework**
- Extend the physics reach beyond Cuoricino while CUORE is being assembled









Tower assembly







Sensors coupling



- The detector performance (e.g. energy resolution) are driven by the sensor-to-crystal coupling (glue spots).
- Features:
 - new semi-automatic system
 - highly-reproducible
 - minimize radioactive recontamination.



Tower assembly







CUORE-0

51/52 NTD connected

51/52 heaters connected

Tower assembly



CUORE-0

51/52 NTD connected

51/52 heaters connected

CUORE

988/988 NTD connected

988/988 heaters connected

Tower installation





**From the CUORE
assembly clean room,
to the Cuoricino
dilution refrigerator**





Experimental setup

- Same cryostat as Cuoricino:
 - inner shield: 1 cm of Roman Lead ($A < 4$ mBq/kg).
 - External shield: 20 cm of Modern Lead.
 - nitrogen flushing
- **gamma background from cryostat shields not expected to change** (test of alpha background)






Outline


- Double beta decay physics
- Thermal detectors
- History of ^{130}Te double beta decay experiments
- **CUORE-0 results**
 - **Detector performance**
 - Neutrinoless double beta decay analysis



Data taking



1st Campaign




2nd Campaign

- Detector assembled in Spring 2012.
- First successful cooldown in March 2013.
- One heater connection lost during the cooldown
 - 51/52 NTD connected
 - 50/52 heater connected
- 2-3 days per months are devoted to ^{232}Th calibrations
- Time between calibrations was devoted to physics data taking, and used for 0νDBD decay search

Calibration data taking
Physics data taking



Exposure overview




- Acquired statistic for 0νDBD decay search:
 - 35.2 kg·yr TeO_2
 - 9.8 kg·yr ^{130}Te



Detector stability


- We measure the resistance of each bolometer daily, to monitor the detector stability over time

CUORE-0 Preliminary



- The resistance of the bolometers are within a factor of 3

CUORE-0 Preliminary



- The bolometers are stable within ~3% over a month timescale



Detector uniformity


- One of the main goal of CUORE-0 was to verify the improvements and the level of reproducibility in the bolometric performance achieved with the new CUORE assembly line.
- We evaluated the distribution of the thermistors temperatures once the detector has been cooled to base temperature and we compared to the Cuoricino one.

$$R(T) = R_0 \exp \left[\frac{T_0}{T} \right]^{1/2}$$



$$T(R) = T_0 \ln^2 \left[\frac{R}{R_0} \right]$$


The narrower distribution of CUORE-0 temperatures compared to Cuoricino shows the **improvement in the reproducibility of the detector construction**





Calibration spectra

CUORE-0 total calibration energy spectrum




Apr-02-2015

- We calibrate the detector using two thoriated tungsten wires source placed in between the outermost cryostat shield and the external lead shield.




Energy resolution

Total fit on the 2615 keV line



Distribution of energy resolution @ 2615 keV



Physics-exposure-weighted harmonic mean


The 5 keV CUORE goal has been reached

	Average FWHM [keV]	RMS of FWHM [keV]
Cuoricino	5.8	2.1
CUORE-0	4.9	2.9



Background reduction

Comparison of the total background spectrum in CUORE-0 and Cuoricino




- ^{238}U and ^{232}Th α lines reduced thanks to the new detector surface treatment.
- ^{238}U γ lines reduced by a factor 2 (better radon control)
- ^{232}Th γ lines not reduced (originate from the cryostat).

Dedicated paper on background model is in preparation



Alpha background rate



	2.7-3.9 MeV	eff
CUORE-0	0.016 ± 0.001	81 ± 1
Cuoricino	0.110 ± 0.001	83 ± 1

- x6 reduction in the alpha continuum region



Outline

- Double beta decay physics
- Thermal detectors
- History of ^{130}Te double beta decay experiments
- **CUORE-0 results**
 - Detector performance
 - **Neutrinoless double beta decay analysis**




Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
- Pulse filtering
- Thermal Gain Stabilization (TGS)
- Energy calibration
- Particle event selection
- Energy spectrum



Analysis technique

- Acquisition of triggered signals



- Each thermistor voltage is continuously sampled at 125Hz
- Once triggered, a 5 sec window is selected for further study of the waveform



The pre trigger voltage is a good proxy for the bolometer temperature before the event

The following 4 sec are analysed to determine the pulse amplitude and to study the pulse shape parameters



Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
 - Raw waveform parameters: baseline, baseline RMS, raw pulse amplitude
- rejection of noisy time intervals






Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
- Pulse filtering
 - **Optimal Filter**: we require that the waveform is consistent with an average reference waveform template. We can optimise energy resolution by exploiting differences in the frequency characteristic of signal and noise events.

- **Decorrelated Optimal Filter**: reduces the correlated noise between adjacent crystals in the array.

Signal and noise power spectrum





New technique developed for CUORE-0 analysis



Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
- Pulse filtering
- Thermal Gain Stabilization (TGS)
 - We need to correct the filtered pulse amplitude for small changes in the energy-to-amplitude response of the bolometer:
 - **heater-TGS**: uses as input the mono-energetic heater pulse
 - **calibration-TGS**: uses the 2.6 MeV line from calibration runs, to correct for the electronic parameters that can affect the bolometers response (drift in amplifier gain or DC offset).




New technique developed for CUORE-0 analysis. We were able to recover the two channels without active heater



Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
- Pulse filtering
- Thermal Gain Stabilization (TGS)
- Energy calibration
 - Each stabilised amplitude is fitted with a gaussian peak and a polynomial function in the region of the peak.
 - We fit a quadratic function with zero intercept to the stabilized pulse amplitude vs know energy to determine the calibration function




$$E(StabAmpl) = a \cdot StabAmpl + b \cdot StabAmpl^2$$





Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
- Pulse filtering
- Thermal Gain Stabilization (TGS)
- Energy calibration
- Particle event selection
 - **Cuts on the Pulse Shape parameters**
 - **Anticoincidence cut**




Exclude multi-site events





Analysis technique

- Acquisition of triggered signals
- Data preprocessing: estimation of raw parameters
- Pulse filtering
- Thermal Gain Stabilization (TGS)
- Energy calibration
 - For each channel and dataset we choose the best performing energy estimator to draw the energy spectrum
- Particle event selection
- Energy spectrum





Efficiency of event selection


- Remove events from periods of low-quality data (total exposure reduced of 7%).

CUORE-0 Preliminary	efficiency [%]	error [%]
Trigger	98,529	0,004
Pile-up and PSA	93,7	0,7
Event containment	88,4	0,09
Accidental coincidence	99,64	0,10

- The total selection efficiency is: **(81,3 ± 0,6)%**



Blinded spectrum



- To blind our data we randomly move a blinded fraction of events within ± 10 keV of the 2615 keV γ -ray peak with events within ± 10 keV of the $0\nu\text{DBD}$ Q-value.

- *The blinding algorithm produces an artificial peak around the $0\nu\text{DBD}$ Q-value and blinds the real $0\nu\text{DBD}$ rate of ^{130}Te .*
- *This method of blinding the data preserves the integrity of the possible $0\nu\text{DBD}$ events while maintaining the spectral characteristics with measured energy resolution and introducing no discontinuities in the spectrum.*




Before the unblinding

- We studied, discussed and froze the analysis methods prior to unblinding:
 - Exposure to be collected
 - Energy reconstruction approach
 - Pulse shape cuts
 - Efficiency calculation approach
 - Detector response model
 - Fitting function in the region of interest




Unblinding

- We unblinded our data in of February 2015, once we surpassed the Cuoricino equivalent sensitivity.






Unblinded spectrum





Unblinded spectrum





Fit in the ROI

- We determined the yield of $0\nu\text{DBD}$ events by performing a simultaneous UEMML fit in the energy region 2470-2570 keV
- The fit has 3 components:
 - a posited peak at the Q-value of ^{130}Te
 - a peak at 2507 keV, attributed to the double gamma events from ^{60}Co in the nearby copper
 - a smooth continuum background, attributed to multi scatter Compton events from ^{208}Tl and surface alpha events





Fit in the ROI

- The best fit value of the $0\nu\text{DBD}$ decay rate is

$$\Gamma_{0\nu} = 0.01 \pm 0.12 \text{ (stat.)} \pm 0.01 \text{ (syst.)} \times 10^{-24} \text{ yr}^{-1}$$

- The background index in the ROI is:


$$0.058 \pm 0.004 \text{ (stat.)} \pm 0.002 \text{ (syst.)} \text{ c/keV/kg/yr}$$



- We set a 90% C.L. Bayesian lower limit of: $T_{1/2} > 2.7 \times 10^{24} \text{ yr}$.




Statistical fluctuations



- We evaluated the statistical properties of the data (event excess above $Q_{\beta\beta}$, dips below and above $Q_{\beta\beta}$).

- A Kolmogorov-Smirnov test shows the data is consistent with the null hypothesis (i.e., the best-fit model but with $\Gamma_{0\nu}$ fixed to zero).
- We compared the value of the binned χ^2 with the distribution from a large set of Toy MC. The 90% of such experiments return a value of $\chi^2 > 43.9$.





Systematics errors


- For each systematic, we run toy MC experiments to evaluate bias on fitted 0νDBD decay rate. We parametrize the bias as p_0 (additive) + $p_1 \cdot \Gamma_{0\nu}$ (scaling)
- **Signal Lineshape**: used variety of different line shapes to model signal.
- **Energy resolution and scale**: we varied the resolution up to 10% of calibration-derived values and we propagated the uncertainties at the Q-value energy.
- **Fit bias**: We evaluated the bias on fitted 0νDBD decay rate as a function of a simulated rate.
- **Bkg function**: we treated the choice of 0-, 1-, or 2-order polynomial for the continuum background as a discrete nuisance parameter.

	Additive (10^{-24} y^{-1})	Scaling (%)
Lineshape	0.007	1.3
Energy resolution	0.006	2.3
Fit bias	0.006	0.15
Energy scale	0.005	0.4
Bkg function	0.004	0.8
Signal normalization	0.7%	




Cuoricino combination

- We combine the CUORE-0 result with the existing 19.75 kg · yr of ^{130}Te exposure from Cuoricino
- The combined 90% C.L. limit is $\mathbf{T_{0\nu} > 4.0 \times 10^{24} \text{ yr.}}$





Limit on $m_{\beta\beta}$



We interpret our combined Bayesian half-life result as a limit on the effective Majorana neutrino mass:

$$m_{\beta\beta} < (270-650) \text{ meV}$$

IBM-2 Phys. Rev. C 91, 034304 (2015)
 QRPA-TU Phys. Rev. C 87, 045501 (2013)
 pnQRPA Phys. Rev. C 91, 024613 (2015)
 ISM Nucl. Phys. A 818, 139 (2009)
 EDF Phys. Rev. Lett. 105, 252503 (2010)



Conclusions

- TeO₂ bolometers offer a well-established, competitive technique in the search for 0νDBD decay

- CUORE-0

- Achieved its energy resolution and background level objectives, surpassing the Cuoricino sensitivity in ~ half the time.
- Did not find evidence of 0νDBD decay.
- Indicated CUORE sensitivity goal is within reach.
- We are going to post the paper to arXiv and to submit it to PRL. We have two more papers in preparation (detector and background).

- CUORE:

- Assembly of the 19 CUORE towers is complete.
- Commissioning of the cryogenic system and experimental infrastructure is in progress
- Plan to start operations by end of 2015.