

Hagedorn spectrum in pure Yang-Mills theories on the lattice¹

Michele Caselle, Alessandro Nada *, Marco Panero

Università degli Studi di Torino

XVI Workshop on Statistical Mechanics and Nonperturbative Field Theory, Bari, 9-11
December 2015



* Speaker

¹M. Caselle, A. Nada, M. Panero, JHEP 1507 (2015) 143

One of the main features of $SU(N)$ non-abelian gauge theories is the existence of a deconfinement phase transition, i.e. a temperature above which gluons are “deconfined”. Our goal is to study the **thermodynamics of pure gauge theories in the confining phase when approaching the deconfinement transition from below**.

In the confining phase the **only** degrees of freedom of the theory without quarks are the **glueballs**: looking at the thermodynamics in the confining phase we have a tool to **explore the glueball spectrum of the theory**.

Our main result is that the thermodynamics of the model can only be described assuming a **string-like** description of glueballs (and thus a Hagedorn spectrum). The fine details of the spectrum agree remarkably well with the predictions of the Nambu-Goto effective string. **This turns out to be an highly non trivial test of the effective string picture of confinement.**

This analysis was performed in the 3+1 dimensional $SU(3)$ model in the pioneering work of Meyer¹. Now, using high precision lattice data for $SU(3)$ ² and a new set of $SU(2)$ data on (3+1) dimensions³, we are in the position to **refine the effective string analysis** and **test its predictive power**. The present results confirm our previous findings⁴ for (2+1) dimensional $SU(N)$ theories (with $N = 2, 3, 4, 5, 6$).

¹H. Meyer, *High-Precision Thermodynamics and Hagedorn Density of States*, 2009


²Sz. Borsanyi *et al.*, *Precision $SU(3)$ lattice thermodynamics for a large temperature range*, 2012

³M. Caselle, A. Nada, M. Panero, *Hagedorn spectrum and thermodynamics of $SU(N)$ Yang-Mills theories*, arXiv:1505.01106

⁴M. Caselle *et al.*, *Thermodynamics of $SU(N)$ Yang-Mills theories in 2+1 dimensions I - The confining phase*, 2011

Thermodynamic quantities

On a $N_t \times N_s^3$ lattice the volume is $V = (aN_s)^3$ (where a is the lattice spacing), while the temperature is determined by the inverse of the temporal extent (with periodic boundary conditions): $T = (aN_t)^{-1}$.



The thermodynamic quantities taken into account will be:

- the **pressure** p , that in the thermodynamic limit (i.e. for large and homogenous systems) can be written as

$$p \simeq \frac{T}{V} \log Z(T, V)$$

- the **trace of the energy-momentum tensor** Δ , that in units of T^4 is

$$\frac{\Delta}{T^4} = \frac{\epsilon - 3p}{T^4} = T \frac{\partial}{\partial T} \left(\frac{p}{T^4} \right).$$

Energy density $\epsilon = \Delta + 3p$ and entropy density $s = \frac{\epsilon + p}{T} = \frac{\Delta + 4p}{T}$ can be easily calculated.

The **pressure** can be estimated by the means of the so-called “integral method”¹:

$$p(T) \simeq \frac{T}{V} \log Z(T, V) = \frac{1}{a^4} \frac{1}{N_t N_s^3} \int_0^{\beta(T)} d\beta' \frac{\partial \log Z}{\partial \beta'}.$$

It can be written (relative to its $T = 0$ vacuum contribution) as

$$\frac{p(T)}{T^4} = -N_t^4 \int_0^{\beta} d\beta' [3(P_\sigma + P_\tau) - 6P_0]$$

where P_σ and P_τ are the expectation values of spacelike and timelike plaquettes respectively and P_0 is the expectation value at zero T .

The **trace of energy-momentum tensor** (also called **trace anomaly**) is simply

$$\frac{\Delta(T)}{T^4} = T \frac{\partial}{\partial T} \left(\frac{p}{T^4} \right) = -N_t^4 T \frac{\partial \beta}{\partial T} [3(P_\sigma + P_\tau) - 6P_0].$$

¹J. Engels *et al.*, *Nonperturbative thermodynamics of SU(N) gauge theories*, 1990

The SU(2) model is a perfect laboratory to test these results.

- It is easy to simulate: very precise results may be obtained with a reasonable amount of computing power.
- The masses of several states of the glueball spectrum are known with remarkable accuracy.
- The deconfinement transition is of second order and thus it is expected to coincide with the Hagedorn temperature, i.e. $T_H \equiv T_c$.
- The infrared physics of the model is very similar to that of the SU(3) theory, with one important difference: the representations of SU(2) are (pseudo)real and thus **only $C = 1$ glueballs exist**. Thus the glueball exponential spectrum contains only **half** of the states with respect to SU(3).

The SU(2) model is a perfect laboratory to test these results.

- It is easy to simulate: very precise results may be obtained with a reasonable amount of computing power.
- The masses of several states of the glueball spectrum are known with remarkable accuracy.
- The deconfinement transition is of second order and thus it is expected to coincide with the Hagedorn temperature, i.e. $T_H \equiv T_c$.
- The infrared physics of the model is very similar to that of the SU(3) theory, with one important difference: the representations of SU(2) are (pseudo)real and thus **only $C = 1$ glueballs exist**. Thus the glueball exponential spectrum contains only **half** of the states with respect to SU(3).

The SU(2) model is a perfect laboratory to test these results.

- It is easy to simulate: very precise results may be obtained with a reasonable amount of computing power.
- The masses of several states of the glueball spectrum are known with remarkable accuracy.
- The deconfinement transition is of second order and thus it is expected to coincide with the Hagedorn temperature, i.e. $T_H \equiv T_c$.
- The infrared physics of the model is very similar to that of the SU(3) theory, with one important difference: the representations of SU(2) are (pseudo)real and thus **only $C = 1$ glueballs exist**. Thus the glueball exponential spectrum contains only **half** of the states with respect to SU(3).

The SU(2) model is a perfect laboratory to test these results.

- It is easy to simulate: very precise results may be obtained with a reasonable amount of computing power.
- The masses of several states of the glueball spectrum are known with remarkable accuracy.
- The deconfinement transition is of second order and thus it is expected to coincide with the Hagedorn temperature, i.e. $T_H \equiv T_c$.
- The infrared physics of the model is very similar to that of the SU(3) theory, with one important difference: the representations of SU(2) are (pseudo)real and thus **only $C = 1$ glueballs exist**. Thus the glueball exponential spectrum contains only **half** of the states with respect to SU(3).

The SU(2) scale setting is fixed by calculating the string tension via the computation of **Polyakov loop correlators** with the multilevel algorithm.

The interquark potential is extracted using

$$V = -\frac{1}{N_t} \log \langle PP \rangle$$


and the string tension is calculated by fitting V for different values of r using

$$V = \sigma r + V_0 - \frac{\pi}{12r}.$$

The values of the string tension are interpolated by a fit to

$$\log(\sigma a^2) = \sum_{j=0}^{n_{\text{par}}-1} a_j (\beta - \beta_0)^j \quad \text{with } \beta_0 = 2.40 \text{ and } n_{\text{par}} = 4$$

Results are presented below along with older data¹.




¹B. Lucini, M. Teper, U. Wenger, *The high temperature phase transition in SU(N) gauge theories*, 2003

N_s^4 at $T = 0$	$N_s^3 \times N_t$ at $T \neq 0$	n_β	β -range	n_{conf}
32^4	$60^3 \times 5$	17	[2.25, 2.3725]	1.5×10^5
40^4	$72^3 \times 6$	25	[2.3059, 2.431]	1.5×10^5
40^4	$72^3 \times 8$	12	[2.439, 2.5124]	10^5

The first two columns show the lattice sizes (in units of the lattice spacing a) for the $T = 0$ and finite-temperature simulations, respectively. In the third column, n_β denotes the number of β -values simulated within the β -range indicated in the fourth column. Finally, in the fifth column we report the cardinality n_{conf} of the configuration set for the $T = 0$ and finite- T simulations.

SU(2): trace of energy-momentum tensor



Despite the small values of N_t the data scale reasonably well.

The behaviour of the system is supposed to be dominated by a **gas of non-interacting glueballs**. The prediction of an **ideal relativistic Bose gas** can be used to describe the thermodynamics of such gas. Its partition function for 3 spatial dimensions is

$$\log Z = (2J + 1) \frac{2V}{T} \left(\frac{m^2}{2\pi} \right)^2 \sum_{k=1}^{\infty} \left(\frac{T}{km} \right)^2 K_2 \left(k \frac{m}{T} \right)$$


where m is the mass of the glueball, J is its spin and K_2 is the modified Bessel function of the second kind of index 2.

Observables such as Δ and p thus can be easily derived:

$$p = \frac{T}{V} \log Z = 2(2J + 1) \left(\frac{m^2}{2\pi} \right)^2 \sum_{k=1}^{\infty} \left(\frac{T}{km} \right)^2 K_2 \left(k \frac{m}{T} \right)$$


$$\Delta = \epsilon - 3p = 2(2J + 1) \left(\frac{m^2}{2\pi} \right)^2 \sum_{k=1}^{\infty} \left(\frac{T}{km} \right) K_1 \left(k \frac{m}{T} \right)$$

SU(2): trace of energy-momentum tensor



Plot of the contribution of the lowest glueball state 0^{++} compared with the data.

SU(2): trace of energy-momentum tensor



The contribution of all SU(2) glueball states with mass $m < 2m_{0^{++}}$.

- Usually the thermodynamics of the system is saturated by the first state (or, in some cases, the few lowest states) of the spectrum due to the exponential dependence on the mass.
- The large gap between the $m_{0^{++}}$ and the $m < 2m_{0^{++}}$ curves and those between them and the data show that the spectrum must be of the **Hagedorn** type, i.e. that **the number of states increases exponentially** with the mass.
- A Hagedorn spectrum is typically the signature of a string-like origin of the spectrum.
- The thermal behaviour of the model in the confining phase is thus a perfect laboratory to study **the nature of this spectrum and of the underlying string model**.
Let us see the consequences of this assumption.

- Usually the thermodynamics of the system is saturated by the first state (or, in some cases, the few lowest states) of the spectrum due to the exponential dependence on the mass.
- The large gap between the $m_{0^{++}}$ and the $m < 2m_{0^{++}}$ curves and those between them and the data show that the spectrum must be of the **Hagedorn** type, i.e. that **the number of states increases exponentially** with the mass.
- A Hagedorn spectrum is typically the signature of a string-like origin of the spectrum.
- The thermal behaviour of the model in the confining phase is thus a perfect laboratory to study **the nature of this spectrum and of the underlying string model**.
Let us see the consequences of this assumption.

- Usually the thermodynamics of the system is saturated by the first state (or, in some cases, the few lowest states) of the spectrum due to the exponential dependence on the mass.
- The large gap between the $m_{0^{++}}$ and the $m < 2m_{0^{++}}$ curves and those between them and the data show that the spectrum must be of the **Hagedorn** type, i.e. that **the number of states increases exponentially** with the mass.
- A Hagedorn spectrum is typically the signature of a string-like origin of the spectrum.
- The thermal behaviour of the model in the confining phase is thus a perfect laboratory to study **the nature of this spectrum and of the underlying string model**.
Let us see the consequences of this assumption.

- Usually the thermodynamics of the system is saturated by the first state (or, in some cases, the few lowest states) of the spectrum due to the exponential dependence on the mass.
- The large gap between the $m_{0^{++}}$ and the $m < 2m_{0^{++}}$ curves and those between them and the data show that the spectrum must be of the **Hagedorn** type, i.e. that **the number of states increases exponentially** with the mass.
- A Hagedorn spectrum is typically the signature of a string-like origin of the spectrum.
- The thermal behaviour of the model in the confining phase is thus a perfect laboratory to study **the nature of this spectrum and of the underlying string model**.
Let us see the consequences of this assumption.

A **closed string model** for the full glueball spectrum that follows the original work of Isgur and Paton¹² can be introduced to account for the values of thermodynamic quantities near the transition. In the closed-string approach glueballs are described in the limit of large masses as “**rings of glue**”, that is **closed tubes of flux modelled by closed bosonic string states**.

The **mass spectrum** of a closed strings gas in D spacetime dimensions is given by

$$m^2 = 4\pi\sigma \left(n_L + n_R - \frac{D-2}{12} \right)$$

where $n_L = n_R = n$ are the total contribution of left- and right-moving phonons on the string.

¹N. Isgur and J. Paton, *A Flux Tube Model for Hadrons in QCD*, 1985

²R. Johnson and M. Teper, *String models of glueballs and the spectrum of $SU(N)$ gauge theories in $(2+1)$ -dimensions*, 2002

Every glueball state corresponds to a given phonon configuration, but associated to each fixed n there are multiple different states whose number is given by $\pi(n)$, i.e. the **partitions** of n .

The **density of states** $\rho(n)$ is expressed through the square of $\pi(n)$

$$\rho(n) = \pi(n_L)\pi(n_R) = \pi(n)^2 \simeq 12(D-2)^{\frac{D-1}{2}} \left(\frac{1}{24n}\right)^{\frac{D+1}{2}} \exp\left(2\pi\sqrt{\frac{2(D-2)n}{3}}\right).$$

The spectral density as a function of the mass (i.e. $\hat{\rho}(m)dm = \rho(n)dn$) can be expressed as

$$\hat{\rho}(m) = \frac{(D-2)^{D-1}}{m} \left(\frac{\pi T_H}{3m} \right)^{D-1} e^{m/T_H}$$

where the **Hagedorn temperature**¹ is defined as

$$T_H = \sqrt{\frac{3\sigma}{\pi(D-2)}}.$$


Finally, the spectral density is used to account for all the states above the mass threshold $2m_{0++}$:


$$\Delta = \sum_{m < 2m_{0++}} (2J+1)\Delta(m, T) + \int_{2m_{0++}}^{\infty} dm \hat{\rho}(m) \Delta(m, T)$$

¹R. Hagedorn, Nuovo Cim. Suppl. 3, 147 (1965)

SU(2): trace of energy-momentum tensor

For $N = 2$ we have that $T_H = T_c$, as the deconfinement transition is 2nd order





The SU(3) case was studied for the first time in 2009 in the pioneering work of Meyer¹. Now, using high precision lattice data² we are in the position to test the Hagedorn behaviour in a very stringent way.

With respect to SU(2)

- SU(3) has **complex** representations, thus glueballs have both $C = +1/ - 1$ and the spectrum contains **approximately twice** the number of glueballs than in the SU(2) case.
- SU(3) has a **first order** deconfining transition, so $T_c < T_H$.

In the effective string framework we can safely fix T_H at the expected Nambu-Goto value, i.e. $T_H = \sqrt{3\sigma/2\pi} \simeq 0.691\sqrt{\sigma}$. Lorentz invariance of the effective string tells us that this should be a very good approximation of the exact result.


The relation between T_H and T_c is:

$$\frac{T_H}{T_c} = 1.098$$


¹H. Meyer, *High-Precision Thermodynamics and Hagedorn Density of States*, 2009

²Sz. Borsanyi et al., *Precision SU(3) lattice thermodynamics for a large temperature range*, 2012

SU(3): trace of energy-momentum tensor



Also in this case the $m < 2m_{0^{++}}$ sector of the glueball spectrum is not enough to fit the behaviour of Δ/T^4 , while including the whole Hagedorn spectrum we find again a remarkable agreement (with no free parameter!)



The doubling of the Hagedorn spectrum is clearly visible in the data!

- The thermodynamics of $SU(2)$ and $SU(3)$ Yang-Mills theories in $D = (3 + 1)$ is well described by a gas of **non-interacting** glueballs.
- The agreement is obtained only assuming a **Hagedorn spectrum** for the glueballs.
- The fine details of the spectrum, in particular the Hagedorn temperature, agree well with the predictions of the **Nambu-Goto effective string**.
- The results agree with previous findings¹ in $D = (2 + 1)$ $SU(N)$ Yang Mills theories with $N = 2, 3, 4, 5, 6$.


¹M. Caselle *et al.*, *Thermodynamics of $SU(N)$ Yang-Mills theories in 2+1 dimensions I - The confining phase*, 2011

The same picture is confirmed by a study performed a few years ago¹ in (2+1) dimensional SU(N) Yang-Mills theories for $N = 2, 3, 4, 6$. Also in this case:

- a Hagedorn spectrum was mandatory to fit the thermodynamic data
- there was a jump between the SU(2) and the SU($N > 2$) case due to the doubling of the spectrum
- we had to fix the Hagedorn temperature to the Nambu-Goto value which, due to the different number of transverse degrees of freedom is different from the (3+1) dimensional one: $T_H = \sqrt{3\sigma/\pi} \simeq 0.977\sqrt{\sigma}$

¹M. Caselle et al., *Thermodynamics of SU(N) Yang-Mills theories in 2+1 dimensions I - The confining phase*, 2011

SU(N) Yang-Mills theories in (2 + 1) dimensions



For $SU(N)$ pure gauge theories on the lattice the dynamics is described by the standard Wilson action

$$S_W = \beta \sum_{p=sp, tp} \left(1 - \frac{1}{N} \text{ReTr} U_p\right)$$


where U_P is the product of four U_μ $SU(N)$ variables on the space-like or time-like plaquette P and $\beta = \frac{2N}{g^2}$.

The partition function is

$$Z = \int \prod_{x, \mu} dU_\mu(x) e^{-S_W}$$

the expectation value of an observable A

$$\langle A \rangle = \frac{1}{Z} \int \prod_{n, \mu} dU_\mu(n) A(U_\mu(n)) e^{-S_W}$$



β	r_{\min}/a	σa^2	aV_0	χ_{red}^2
2.27	2.889	0.157(8)	0.626(14)	0.6
2.30	2.889	0.131(4)	0.627(30)	0.1
2.32	3.922	0.115(6)	0.627(32)	2.3
2.35	3.922	0.095(3)	0.623(20)	0.2
2.37	3.922	0.083(3)	0.621(18)	1.0
2.40	4.942	0.068(1)	0.617(10)	1.4
2.42	4.942	0.0593(4)	0.613(5)	0.1
2.45	4.942	0.0482(2)	0.608(4)	0.4
2.47	4.942	0.0420(4)	0.604(5)	0.3
2.50	5.954	0.0341(2)	0.599(2)	0.1
2.55	6.963	0.0243(13)	0.587(11)	0.2
2.60	7.967	0.0175(16)	0.575(16)	0.3

Results for the string tension in units of the inverse squared lattice spacing at different values of the Wilson action parameter β (first column). V was extracted from Polyakov loop correlators on lattices of temporal extent $L_t = 32a$.

