

Spectroscopic study on ³⁹Ca using the ⁴⁰Ca(d,t)³⁹Ca reaction for classical nova endpoint nucleosynthesis

J. Liang^a, A.A. Chen^a, M. Anger^c, S. Bishop^c, T. Faestermann^c, C. Fry^b, R. Hertenberger^d, A. Psaltis^a, D. Seiler^c, P. Tiwari^b, H-F. Wirth^d, C. Wrede^b

^aDepartment of Physics and Astronomy, McMaster University, Hamilton, Ontario L8S 4M1, Canada, ^bDepartment of Physics and Astronomy, National Superconducting Cyclotron Laboratory, Michigan State University, 640 S. Shaw Ln., East Lansing, MI 48864, USA ^cDepartment of Physics, Technical University Munich, James-Franck-Str. 1, 685748 Garching, Germany ^dFaculty of Physics, Ludwig-Maximilians-Universität München, Geschwister-Scholl-Platz 1, 80539 Munich, Germany

In classical nova nucleosythesis repeated proton capture reactions and beta-decays produce proton-rich isotopes and the endpoint of this nucleosynthesis typically occurs in nuclei close to A \sim 40. There is currently a discrepancy between the observed and predicted isotopic abundances in this mass region. One particular reaction, $^{38}\text{K}(\text{p},\gamma)^{39}\text{Ca}$ is important in this regard. Nova simulations show that this reaction can alter the isotopic abundances of ^{38}Ar , ^{39}Ar , and ^{40}Ca significantly when the reaction rate is varied by its maximum uncertainty. Thus, it is important to constrain uncertainties of this reaction rate to accurately predict isotopic abundances.

Although a recent direct measurement has reduced the reaction rate uncertainty, further work is needed to constrain this reaction rate. Specifically, additional measurements to precisely probe the low energy resonances within the Gamow window. To that end, I will present the preliminary results measuring these astrophysically important levels in ³⁹Ca using the reaction ⁴⁰Ca(d,t)³⁹Ca. The experiment was carried out at the Maier-Leibnitz-Laboratory (MLL) using the 14 MV MP-Tandem accelerator and Quadrupole 3-Dipole (Q3D) magnetic spectrograph.