



**BABAR**  
TM and © Nelvana, All Rights Reserved




# *Results on Charmonium-like States at BaBar*

Valentina Santoro

INFN Ferrara

Representing the BaBar Collaboration

QCD@ Work , June 18-21 , Lecce, Italy



- **Introduction and Overview**
- **Search for the  $Z_1(4050)^+$  and  $Z_2(4250)^+$  states in  $\bar{B}^0 \rightarrow \chi_{c1} K^- \pi^+$  and  $B^+ \rightarrow \chi_{c1} K_s^0 \pi^+$** 

**PRD 85, 052003 (2012)**
- **Study of the  $J/\psi\omega$  final state in two-photon collisions**
- **Search for resonances decaying to  $\eta_c(1S) \pi^+ \pi^-$** 


**arXiv:1206.2008**
- **Study of the reaction  $e^+ e^- \rightarrow J/\psi \pi^+ \pi^-$  via initial state radiation**

**arXiv:1204.2158**
- **Study of the reaction  $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^-$  via initial state radiation**

# Charmonium spectrum

(June 2012)

Eur. Phys.J.C71, 1534 (2011)



- Charmonium properties are well understood up to  $\psi(3770)$  (i.e. about the  $D\bar{D}$  threshold)
  - Many unexpected states above the  $D\bar{D}$  threshold.
  - Several exotic hypotheses on their nature : e.g. tetraquarks, hadronic molecules, hybrids ...
- To identify exotics:
- Measure  $J^{PC}$  that is forbidden for charmonium:  $0^{+-}$ ,  $1^{-+}$ ,  $2^{+-}$
  - Observed a narrow width above  $D\bar{D}$  threshold
  - Observed a  $c\bar{c}$ -like state with charge and/ or strangeness

# Charmonium-like States


**Over 16 new resonances only 5 have been seen by different experiments**

OK = confirmed by different experiments  
 N.C. = Seen by only one experiment not by other  
 N.Y.C = Not yet confirmed


| State                              | m(MeV/c <sup>2</sup> ) | $\Gamma$ (MeV)         | $J^{PC}$                         | Process (mode)   | Experiment         | Year | Status |
|------------------------------------|------------------------|------------------------|----------------------------------|--|--------------------|------|--------|
| X(3823)                            | $3823.5 \pm 2.8$       | $4 \pm 6$              | ?/2 <sup>--</sup>                | $B \rightarrow K(\chi_{c1}\gamma)$   | Belle              | 2012 | N.Y.C  |
| X(3872)                            | $3871.52 \pm 0.20$     | $1.3 \pm 0.6$          | 1 <sup>++</sup> /2 <sup>++</sup> | $B \rightarrow K(\pi^+\pi^-J/\psi)$<br>$p\bar{p} \rightarrow (\pi^+\pi^-J/\psi) + \dots$ | Belle, BABAR, LHCb | 2003 | OK     |
|                                    |                        |                        |                                  | $B \rightarrow K(J/\psi\omega)$  | CDF, DØ, CMS       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(D^{*0}\bar{D}^0)$   | Belle, BABAR       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(J/\psi\gamma)$  | Belle, BABAR       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(\psi(2S)\gamma)$  | BABAR              |      | N.C.   |
| X(3915)                            | $3915.6 \pm 3.1$       | $28 \pm 10$            | 0/2 <sup>?+</sup>                | $B \rightarrow K(J/\psi\omega)$  | Belle, BABAR       | 2004 | OK     |
| X(3940)                            | $3942^{+9}_{-8}$       | $37^{+27}_{-17}$       | ? <sup>?</sup> +                 | $e^+e^- \rightarrow e^+e^-(J/\psi\omega)$<br>$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$     | Belle, BABAR       | 2007 | N.Y.C. |
| Y(4008)                            | $4008^{+121}_{-49}$    | $226 \pm 97$           | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma J/\psi\pi^+\pi^-$   | Belle              | 2007 | N.C.   |
| Z <sub>1</sub> (4050) <sup>+</sup> | $4051^{+24}_{-43}$     | $82^{+51}_{-55}$       | ?                                | $B \rightarrow K(\pi^+\chi_{c1}(1P))$  | Belle              | 2008 | N.C.   |
| Y(4140)                            | $4143 \pm 3.0$         | $15^{+11}_{-7}$        | ? <sup>?</sup> +                 | $B \rightarrow K(\phi J/\psi)$   | CDF                | 2009 | N.C.   |
| X(4160)                            | $4156^{+29}_{-25}$     | $139^{+113}_{-65}$     | ? <sup>?</sup> +                 | $e^+e^- \rightarrow J/\psi(D\bar{D}^*)$  | Belle              | 2007 | N.Y.C. |
| Z <sub>2</sub> (4250) <sup>+</sup> | $4248^{+185}_{-45}$    | $177^{+321}_{-72}$     | ?                                | $B \rightarrow K(\pi^+\chi_{c1}(1P))$  | Belle              | 2008 | N.C.   |
| Y(4260)                            | $4263 \pm 5$           | $108 \pm 14$           | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma J/\psi\pi^+\pi^-$   | BABAR, CLEO, Belle | 2005 | OK     |
| Y(4274)                            | $4274.4^{+8.4}_{-6.7}$ | $32^{+22}_{-15}$       | ? <sup>?</sup> +                 | $e^+e^- \rightarrow J/\psi\pi^0\pi^0$<br>$B \rightarrow K(\phi J/\psi)$                  | CLEO               |      |        |
| X(4350)                            | $4350.6^{+4.6}_{-5.1}$ | $13.3^{+18.4}_{-10.0}$ | 0, 2 <sup>++</sup>               | $e^+e^- \rightarrow e^+e^-(\phi J/\psi)$   | CDF                | 2010 | N.C.   |
| Belle                              |                        |                        |                                  |  | Belle              | 2009 | N.Y.C. |
| Y(4360)                            | $4353 \pm 11$          | $96 \pm 42$            | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma\psi(2S)\pi^+\pi^-$  | BABAR, Belle       | 2007 | OK     |
| Z(4430) <sup>+</sup>               | $4443^{+24}_{-18}$     | $107^{+113}_{-71}$     | ?                                | $B \rightarrow K(\pi^+\psi(2S))$   | Belle              | 2007 | N.C.   |
| X(4360)                            | $4634 \pm 12$          | $92^{+41}_{-32}$       | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma\Lambda_c^+\Lambda_c^-$  | Belle              | 2007 | N.Y.C. |
| Y(4660)                            | $4664 \pm 12$          | $48 \pm 15$            | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma\psi(2S)\pi^+\pi^-$  | Belle, BABAR       | 2007 | OK     |

**NEW Results from BaBar**

# Exotic charmonium-like states interpretation




**Molecular state:**  
loosely bound state of a pair of mesons.  
The dominant binding mechanism should be pion exchange. Being weakly bound, mesons tend to decay as if they were free.




**Tetraquark:**  
Bound state of four quarks, i.e. diquark-antidiquark  
Strong decays proceed via rearrangement processes.

NA Tornqvist PLB 590, 209 (2004)  
ES Swanson PLB 598, 197 (2004)  
E Braaten & T Kusunoki  
PRD 69 074005 (2004)  
CY Wong PRC 69, 055202 (2004)  
MB Voloshin PLB 579, 316 (2004)  
F Close & P Page PLB 578, 119 (2004) ....



**Charmonium hybrids**  
States with an excited gluonic degree of freedom  
Lattice and model predictions for the lowest lying hybrid:  
 $m \sim 4200$  MeV

L Maiani et al PRD 71, 014028 (2005)  
T-W Chiu & TH Hsieh PRD 73, 111503 (2006)  
D Ebert et al PLB 634, 214 (2006)  
...




**Conventional charmonium**

P Lacock et al (UKQCD) PLB 401, 308 (1997)  
SL Zhu PLB 625, 212 (2005)  
FE Close, PR Page PLB 628, 215 (2005)  
E Kou, O Pene PLB 631, 164 (2005)  
...

C Meng & KT Chao PRD 75, 114002 (2007)  
W Dunwoodie & V Ziegler PRL 100 062006 (2008)  
O Zhang, C Meng & HQ Zheng arXiv:0901.1553  
...


# Study of the $J/\psi\omega$ in two-photon interactions




# $\gamma\gamma \rightarrow J/\psi\omega$ (Analysis Motivation): look for X(3915) (1)

The X(3915) was seen by Belle and then confirmed by BaBar in  $B \rightarrow X(3915)K$ , with the  $X(3915) \rightarrow J/\psi\omega$

PRL 94, 182002 (2005)




PRD, 82 011101 R (2010)



B decays

Belle also observed the X(3915) in the process  $\gamma\gamma \rightarrow X(3915) \rightarrow J/\psi\omega$

PRL, 104 092001 (2010)



$\gamma\gamma$  interactions

Interpretation of X(3915) as the  $\chi_{c0}(2P)$  or  $\chi_{c2}(2P)$  state has been suggested  
But the  $\Gamma_{\gamma\gamma}(X(3915))B(X(3915) \rightarrow J/\psi\omega)$  reported by Belle is unexpectedly large compared to other excited charmonia.

The X(3872) was seen in 2003 by Belle and soon confirmed by several experiments

### X(3872) Properties

Narrow ( $\Gamma \leq 2.3$  MeV according to PDG 2012) with mass  
 $m(X)=3871.61 \pm 0.25$  MeV/c<sup>2</sup>

Observed in X(3872) J/ $\psi\pi^+\pi^-$ , dipion mass is “ $\rho$ -like” but possible “ $\rho-\omega$ ” interference


Also seen in B decays:  $X(3872) \rightarrow D^0\bar{D}^{*0}$ ,

$X(3872) \rightarrow J/\psi\omega$ ,  $X(3872) \rightarrow J/\psi\gamma$ ,

$X(3872) \rightarrow \psi(2S)\gamma$  seen by BaBar but not by Belle

C is positive from the observation of the decay  $J/\psi\gamma$ ,

Spin-parity identified as either  $J^{PC}=1^{++}$  or  $2^{-+}$ ; isospin I=0



Phys. Rev. Lett. 91, 262001 (2003)

### X(3872) interpretation

Conventional Charmonium:  $\chi_{c1}(2^3P_1)$  ( $1^{++}$ ) or  $\eta_{c2}(1^1D_2)$  ( $2^{-+}$ )

$D^0\bar{D}^{*0}$  Molecular interpretation:


$m(D^0) + m(\bar{D}^{*0}) = 3871.73 \pm 0.29$  MeV/c<sup>2</sup>

Compatible with  $J^{PC}=1^{++}$  assignment;

The observation of  $\gamma\gamma \rightarrow X(3872)$  would imply  $J^{PC}=2^{-+}$  and favor the charmonium interpretation

# $\gamma\gamma \rightarrow J/\psi\omega$ : new BaBar Results

BaBar with  $520 \text{ fb}^{-1}$  collected at the Y(nS) sample ( $n=2,3,4$ ) confirmed the evidence of the X(3915) ( $7.6 \sigma$ ) in  $\gamma\gamma \rightarrow X(3915) \rightarrow J/\psi\omega$




PRL 104, 092001 (2010)

|  | <i>BABAR</i>             | Belle              |
|--|--------------------------|--------------------|
| Mass ( $\text{MeV}/c^2$ )  | $3919.4 \pm 2.2 \pm 1.6$ | $3915 \pm 3 \pm 2$ |
| Width (MeV)  | $13 \pm 6 \pm 3$         | $17 \pm 10 \pm 3$  |
| $\Gamma_{\gamma\gamma} \times \mathcal{B} (\text{J}=0) \text{ (eV)}$ | $52 \pm 10 \pm 3$        | $61 \pm 17 \pm 8$  |
| $\Gamma_{\gamma\gamma} \times \mathcal{B} (\text{J}=2) \text{ (eV)}$ | $10.5 \pm 1.9 \pm 0.6$   | $18 \pm 5 \pm 2$   |

If  $\Gamma_{\gamma\gamma} = \mathcal{O}(1 \text{ keV})$  (typical  $c\bar{c}$ )  
 then  $\mathcal{B}(J/\psi\omega) > (1-6)\%$   
 which is relatively large compared to  
 charmonium model predictions

No evidence of the X(3872) → limit for  $J=2$  hypothesis  
 $\Gamma_{\gamma\gamma}(\text{X}(3872)) \times \mathcal{B}(\text{X}(3872) \rightarrow J/\psi\omega) (\text{J}=2) < 1.7 \text{ eV}$

# Search for resonances decaying to $\eta_c \pi^+ \pi^-$ in two-photon interactions





- BaBar using  $474 \text{ fb}^{-1}$  studied the process  $\gamma\gamma \rightarrow X \rightarrow \eta_c(1S) \pi^+ \pi^-$  where  $X$  stands for one of the resonances  $\chi_{c2}(1P)$ ,  $\eta_{c2}(2S)$ ,  $X(3872)$ ,  $X(3915)$  or  $\chi_{c2}(2P)$ ; the  $\eta_c(1S)$  is reconstructed using the decay  $\eta_c(1S) \rightarrow K_s^0 K\pi$ ;  $K_s^0 \rightarrow \pi^+ \pi^-$
- Predictions for  $B(\eta_{c2}(2S) \rightarrow \eta_{c2}(1S) \pi^+ \pi^-) \sim 2.2\%$  obtained from  $\Gamma(\eta_{c2}(2S) \rightarrow \eta_c(1S) \pi^+ \pi^-) / \Gamma(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) \sim 2.9$



M.B.Voloshin Mod. Phys. Lett A 17:1533-1538, 2002 ×

- If the  $X(3872)$  is the  $1^1D_2$  state  $\eta_{c2}$  the branching fraction  $B(X(3872) \rightarrow \eta_c \pi^+ \pi^-)$  could be significantly larger than  $B(X(3872) \rightarrow J/\psi \pi^+ \pi^-)$
- The quantum numbers  $J^{PC}=2^{-+}$  of the  $\eta_{c2}$  are consistent with the CDF result for the  $X(3872)$



S.L. Olsen (Belle Collaboration)  
Int. J. Mod. Phys. A497 20, 240 (2005)

PRL 98, 132002 (2007)



## Signal extraction in two steps

**BABAR**  
preliminary**BABAR**  
preliminary

\chi\_{c2}(1P)

**BABAR**  
preliminary**BABAR**  
preliminary

\eta\_c(2S)

**BABAR**  
preliminary**BABAR**  
preliminaryX(3872)  
 $X(3915)$   
 $\chi_{c2}(2P)$ 

- $m(K_S^0 K^\pm \pi^\mp)$  distribution parameters of the combinatoric background from a one-dimensional fit to  $m(K_S^0 K^\pm \pi^\mp)$
- Two-dimensional fit in  $m(K_S^0 K^\pm \pi^\mp)$  and  $m(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)$

No resonant signal observed  
observed contribution from non-resonant

$$\gamma\gamma \rightarrow X \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$$


| Resonance       | $\Gamma_{\gamma\gamma} \mathcal{B}(\text{eV})$ |      |
|-----------------|--|------|
|                 | Central value                                  | UL   |
| $\chi_{c2}(1P)$ | $7.2^{+5.5}_{-4.4} \pm 2.9$                    | 15.7 |
| $\eta_c(2S)$    | $65^{+47}_{-44} \pm 18$                        | 133  |
| $X(3872)$       | $-4.5^{+7.7}_{-6.7} \pm 2.9$                   | 11.1 |
| $X(3915)$       | $-13^{+12}_{-12} \pm 8$                        | 16   |
| $\chi_{c2}(2P)$ | $-16^{+15}_{-14} \pm 6$                        | 19   |

Using  $\mathcal{B}(\chi_{c2}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)$  and  
 $\mathcal{B}(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)$  we obtain:

**First Measurement**

$$\begin{aligned} \mathcal{B}(\chi_{c2}(1P) \rightarrow \eta_c(1S) \pi \pi) &< 2.2\% @ 90\% \text{ CL} \\ \mathcal{B}(\eta_c(2S) \rightarrow \eta_c(1S) \pi \pi) &< 7.4\% @ 90\% \text{ CL} \end{aligned}$$

# Search for $Z_1(4050)^+$ and $Z_2(4250)^+$ states in $\bar{B}^0 \rightarrow \chi_{c1} K^- \pi^+$ and $B^+ \rightarrow \chi_{c1} K_S^0 \pi^+$




# The charged Z family

- Belle observed a broad, charged charmonium-like states in  $(c\bar{c})K\pi$  Dalitz plot analyses
  - $Z(4430)^+$  in  $B \rightarrow \psi(2S)\pi^+ K$
  - $Z_1(4050)^+$  and  $Z_2(4250)^+$  in  $\bar{B}^0 \rightarrow \chi_{c1}\pi^+ K^-$
- Quark content at least  $c\bar{c}u\bar{d}$ : no simple  $q\bar{q}$  meson

Maiani: arXiv: 0708.3997  
 Karliner & Lipkin arXiv: 0802.0649


PRL 100, 142001 (2008)



$Z(4430)^+ \rightarrow \psi(2S)\pi^+$

Not confirmed by BaBar experiment  
 that studied also the  $J/\psi\pi^-K^+$  and the  
 $J/\psi\pi^-K_S^0$  channel

PRD 78, 072004 (2008)




$Z_1(4050)^+ \rightarrow \chi_{c1}\pi^+$


$Z_2(4250)^+ \rightarrow \chi_{c1}\pi^+$

PRD 79, 112001 (2009)

- BaBar using 429 fb<sup>-1</sup> studied the processes  $B^+ \rightarrow \chi_{c1} K_S^0 \pi^+$  and  $\bar{B}^0 \rightarrow \chi_{c1} K^- \pi^+$  to search for Z<sub>1</sub> and Z<sub>2</sub> found by Belle ( $\chi_{c1} \rightarrow J/\psi \gamma$ )
- Binned  $\chi^2$  fits to the background-subtracted and efficiency-corrected K $\pi$  mass spectra in terms of S, P and D wave amplitudes



| Channel                                     | S wave         | P wave                           | D wave         | $\chi^2/NDF$ |
|---|----------------|----------------------------------|----------------|--------------|
| $\bar{B}^0 \rightarrow \chi_{c1} K^- \pi^+$ | $40.4 \pm 2.2$ | $37.9 \pm 1.3$<br>$10.3 \pm 1.5$ | $11.4 \pm 2.0$ | 58/54        |
| $B^+ \rightarrow \chi_{c1} K_S^0 \pi^+$     | $42.4 \pm 3.5$ | $37.1 \pm 3.2$<br>$10.4 \pm 2.5$ | $10.1 \pm 3.1$ | 55/54        |




To represent the angular structure we compute the efficiency-corrected Legendre polynomial moments  $\langle Y_L^0 \rangle$  in each K $\pi$  mass interval by correcting for efficiency and then weighting each event by the  $Y_L^0(\cos\theta)$  functions

- Using the information from the  $K\pi$  system a description of the  $\chi_{c1}\pi$  mass distribution is studied. A MC simulation for  $B \rightarrow \chi_{c1} K\pi$  has been performed.
- The best  $\chi^2/NDF$  obtained is for  $L_{MAX}=5$  (where  $L$  is the order of the Legendre polynomial moments)

**Black dots: data**


**Red curve:** MC simulation with  $L_{MAX}=5$  (angular weights)

**Dotted blue curve:** Results of the simulation with no angular weights




- ✓ The result of the simulation with  $L_{MAX}=5$  is superimposed on the data (red curve).
- ✓ The excellent description of the data indicated that the angular information from the  $K\pi$  channel with  $L_{MAX}=5$  is able to account for the structures observed in the  $\chi_{c1}K\pi$  projection.
- ✓ **This indicates the absence of significant structure in the exotic  $\chi_{c1}\pi^+$  channel**

Fit the  $\chi_{c1}\pi$  mass spectrum using two scalar Breit-Wigners with parameters fixed to the Belle measurement ( $m(Z_1^+) = 4051 \text{ GeV}/c^2$ ,  $\Gamma(Z_1^+) = 82 \text{ MeV}$ ,  $m(Z_2^+) = 4248 \text{ GeV}/c^2$ ,  $\Gamma(Z_2^+) = 177 \text{ MeV}$ )



Belle: maximal resonant activity in window  $1.0 < m^2_{K\pi} < 1.75 (\text{GeV}/c^2)^2$



For two resonances fit

$$\mathcal{B}(\bar{B}^0 \rightarrow Z_1^+ K^-) \times \mathcal{B}(Z_1^+ \rightarrow \chi_{c1}\pi^+) < 1.8 \times 10^{-5}$$

$$\mathcal{B}(\bar{B}^0 \rightarrow Z_2^+ K^-) \times \mathcal{B}(Z_2^+ \rightarrow \chi_{c1}\pi^+) < 4.0 \times 10^{-5}$$


For a single resonance fit

$$\mathcal{B}(\bar{B}^0 \rightarrow Z^+ K^-) \times \mathcal{B}(Z^+ \rightarrow \chi_{c1}\pi^+) < 4.7 \times 10^{-5}$$

In all cases we obtained very low ( $< 2\sigma$ ) statistical significances

Within (large) uncertainties, limit compatible with the Belle result

# Study of the reaction $e^+e^- \rightarrow J/\psi\pi^+\pi^-$ via initial-state radiation



## The $Y(4260)$

$J^{PC} = 1^{--}$

Discovered at BaBar in ISR-production of  $J/\psi \pi^+\pi^-$  events in 2005 ( $233 \text{ fb}^{-1}$ )


Phys. Rev. Lett. 95 (2005) 142001

$$m_Y = (4259 \pm 8^{+2}_{-6}) \text{ MeV}/c^2$$

$$\Gamma_Y = (88 \pm 23^{+6}_{-4}) \text{ MeV}$$


Confirmations by CLEO-c, CLEO-III and Belle with some spread in the resonance parameters.

Phys. Rev. Lett. 99, 182004 (2007)



$Y(4008)$

All the  $1^{--}$  slots in the charmonium spectrum were already filled



The Belle result suggested the existence of another broad structure the  $Y(4008)$



$$M_{Y_{4260}} = 4247 \pm 12^{+17}_{-32}$$


$$\Gamma_{Y_{4260}} = 108 \pm 19 \pm 10$$

$$M_{Y_{4008}} = 4008 \pm 40^{+114}_{-28}$$

$$\Gamma_{Y_{4008}} = 226 \pm 44 \pm 87$$

The  $\psi(2S)$  region (3.5-4 GeV/c<sup>2</sup>)

We investigated the region below 4 GeV/c<sup>2</sup> ( $\psi(2S)$  and  $\psi(3770)$  region) of the  $J/\psi \pi^+\pi^-$  final state for the first time, since in this region we observe an excess of events




The excess of events can be attributed to different sources such as  $\psi(2S)$  tail,  $J/\psi \pi^+\pi^-$  continuum or  $\psi(3770)$  decay to  $J/\psi \pi^+\pi^-$ , since BES (PLB 605, 63(2005) and CLEO (PRL 96, 082004 (2006) reported such decay mode for the  $\psi(3770)$


In order to understand the possible contribution coming from the  $\psi(2S)$  tail a detailed study of the  $\psi(2S)$  lineshape has been performed

$$\sigma(e^+e^- \rightarrow \gamma\psi(2S)) = 14.5 \pm 0.7 \text{ pb}$$


$$\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.29 \pm 0.05 \text{ keV}$$



The conclusion is that it is not possible to discount the possibility of a contribution from  $e^+e^- \rightarrow J/\psi \pi^+\pi^-$  continuum cross section in this region

The Y(4260) region (up to 5.5 GeV/c<sup>2</sup>)

We performed an extended-maximum-likelihood fit to the signal region  $J/\psi \pi^+\pi^-$  distribution and simultaneously to the background distribution  
The fit is corrected for efficiency and luminosity



$$m(Y(4260)) = 4244 \pm 5 \pm 4 \text{ MeV}/c^2$$

$$\Gamma(Y(4260)) = 114^{+16}_{-15} \pm 7 \text{ MeV}$$


$$\Gamma_{e^+e^-} \times B(J/\psi \pi^+\pi^-) = 9.2 \pm 0.8 \pm 0.7 \text{ eV}$$

arXiv:1204.2158  
SUBMITTED TO PRD(RC)


- No evidence of the additional state claimed by Belle ~4 GeV/c<sup>2</sup>
- Excess of events above 3.74 GeV/c<sup>2</sup> might result from the  $\psi(2S)$  tail and a possible  $J/\psi \pi^+\pi^-$  continuum contribution
- Very clear signal for the Y(4260)
- The small excess of events ~ 4.5 GeV/c<sup>2</sup> attributed to statistical fluctuation

# $\pi^+\pi^-$ invariant mass distribution in the Y(4260) decay (1)

$4.15 < m(J/\psi\pi^+\pi^-) < 4.45 \text{ GeV}/c^2$



- The distribution seems to peak around the  $f_0(980)$  mass; however the peak is displaced from the usual  $f_0(980)$  position, since it is around 940 MeV/c<sup>2</sup>
- The fact that the peak is displaced and the particular shape of the  $m(\pi^+\pi^-)$  mass distribution seems to suggest a possible interference between the  $f_0(980)$  and  $m(\pi^+\pi^-)$  continuum




- We define  $\theta_\pi$  as the angle between the  $\pi^+$  direction and that of the recoil  $J/\psi$  both in the dipion rest frame
- The distribution is consistent with the  $S$ -wave behaviour ( $\chi^2/\text{NDF}=12.3/9$ ; Probability =19.7%)

# $\pi^+\pi^-$ invariant mass distribution in the Y(4260) decay (2)

- To test the possibility of an interference effect between the  $f_0(980)$  and the  $m(\pi^+\pi^-)$  continuum the  $f_0(980)$  line shape is taken from the BaBar analysis  $D_s^+ \rightarrow \pi^+\pi^-\pi^+$

PRD 79, 032003 (2009)



we interpolate on these distributions and use this amplitude to describe the  $f_0(980)$  in the following model



$$|\sqrt{pol} + e^{i\phi} F_{f_0(980)}|^2$$

where  $pol$  is a polynomial function used to describe the  $m(\pi^+\pi^-)$  continuum and  $F_{f_0(980)}$  is the interpolated function;  $\phi$  allows for a phase difference between these amplitudes

The result indicates that there is a real  $f_0(980)$  contribution to the decay of the  $Y(4260)$  to  $J/\psi\pi^+\pi^-$  but its contribution is not dominant


$$\frac{\mathcal{B}(Y_{4260} \rightarrow J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)}{\mathcal{B}(Y_{4260} \rightarrow J/\psi\pi^+\pi^-)} = (17 \pm 13)\%$$

# Study of the reaction $e^+e^- \rightarrow \psi(2S)\pi^+\pi^-$ via initial-state radiation



# Y(4350) and Y(4660) $\rightarrow \pi^+ \pi^- \psi(2S)$ (Motivation).


Y(4260) discovered in ISR  $\pi^+ \pi^- J/\psi$ . How about  $\pi^+ \pi^- \psi(2S)$  in ISR?



In  $\psi(2S)\pi^+\pi^-$  BaBar observed the Y(4350)  
 Single resonance fit =>  
 $m = (4324 \pm 24) \text{ MeV}/c^2$ ,  $\Gamma = (172 \pm 33) \text{ MeV}$   
 Incompatible with  $\psi(4415)$ ;  
 Poorly described by Y(4260)





PRL 99, 142002 (2007)



| State   | M, MeV/c <sup>2</sup> | $\Gamma_{\text{tot}}$ , MeV |
|---------|-----------------------|-----------------------------|
| Y(4350) | $4324 \pm 24$         | $172 \pm 33$                |
| Belle   | $4361 \pm 9 \pm 9$    | $74 \pm 15 \pm 10$          |
| Belle   | $4664 \pm 11 \pm 5$   | $48 \pm 15 \pm 3$           |

# ISR $\pi^+\pi^-\psi(2S)$ : New BaBar Result

- BaBar update using the full dataset, including the Y(2S) and Y(3S) data
- $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$




| Parameters   | First Solution<br>[constructive interference] | Second Solution<br>[destructive interference] |
|--|---|---|
| Mass Y(4360)(MeV/c <sup>2</sup> )                    | $4340 \pm 16 \pm 9$                           |   |
| Width Y(4360)(MeV)                                   | $94 \pm 32 \pm 13$                            |   |
| $\mathcal{B} \times \Gamma_{ee}(Y(4360))(\text{eV})$ | $6.0 \pm 1.0 \pm 0.5$                         | $7.2 \pm 1.0 \pm 0.6$                         |
| Mass Y(4660)(MeV/c <sup>2</sup> )                    | $4669 \pm 21 \pm 3$                           |   |
| Width Y(4660)(MeV)                                   | $104 \pm 48 \pm 10$                           |   |
| $\mathcal{B} \times \Gamma_{ee}(Y(4660))(\text{eV})$ | $2.7 \pm 1.3 \pm 0.5$                         | $7.5 \pm 1.7 \pm 0.7$                         |
| $\phi(^{\circ})$                                     | $12 \pm 27 \pm 4$                             | $-78 \pm 12 \pm 3$                            |

Confirm the BELLE enhancement  
 $\sim 4.6 \text{ GeV}/c^2$   
The obtained parameters for the Y(4360) and Y(4660) are consistent with the Belle results



- ✓ Charmonium spectroscopy has been revitalized by the discovery of many new states above the open charm threshold. A review of some of these new states has been presented.
- ✓ Many experimental results have been shown, with just enough data to whet the appetite, but at a statistical level which does not permit a clear understanding of the observed signals;
- ✓ More data are required, possibly from LHCb, but also from the SuperB or BELLE-II projects, should they materialize in the future;

# BACK-UP SLIDES




# The BaBar detector and data sample



BaBar is a powerful  $b$  factory: 467 million  $B\bar{B}$  pairs in the total data sample

BaBar is also a  $c$  factory: 1.3 million Charm events per  $\text{fb}^{-1}$




Offpeak (10.54GeV) + Scan above  $Y(4S)$ :  $53.9 \text{ fb}^{-1}$

$Y(4S)$ :  $432 \text{ fb}^{-1}$   
 $Y(3S)$ :  $30.2 \text{ fb}^{-1}$   
 $Y(2S)$ :  $14.5 \text{ fb}^{-1}$


# Charmonium production at the B-factories

## B decays




States of any quantum numbers can be formed

## Double charmonium production




## $2\gamma$ production




$J^{PC}=0^{\pm+}, 2^{\pm+} \dots$

## Initial State Radiation (ISR)



$J^{PC}=1^-$

# All in all



# Charmonium-like States (2)


| State                              | $m(\text{MeV}/c^2)$    | $\Gamma(\text{MeV})$   | $J^{PC}$                         | Process (mode)                                    | Experiment         | Year | Status |
|------------------------------------|------------------------|------------------------|----------------------------------|---|--------------------|------|--------|
| X(3823)                            | $3823.5 \pm 2.8$       | $4 \pm 6$              | ?/2 <sup>-+</sup>                | $B \rightarrow K(\chi_{c1}\gamma)$                | Belle              | 2012 | N.Y.C  |
| X(3872)                            | $3871.52 \pm 0.20$     | $1.3 \pm 0.6$          | 1 <sup>++</sup> /2 <sup>++</sup> | $B \rightarrow K(\pi^+\pi^-J/\psi)$               | Belle, BABAR, LHCb | 2003 | OK     |
|                                    |                        |                        |                                  | $p\bar{p} \rightarrow (\pi^+\pi^-J/\psi) + \dots$ | CDF, DØ, CMS       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(J/\psi\omega)$                   | Belle, BABAR       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(D^{*0}\bar{D}^0)$                | Belle, BABAR       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(J/\psi\gamma)$                   | Belle, BABAR       |      |        |
|                                    |                        |                        |                                  | $B \rightarrow K(\psi(2S)\gamma)$                 | Belle, BABAR       |      |        |
| X(3915)                            | $3915.6 \pm 3.1$       | $28 \pm 10$            | 0/2 <sup>?</sup> +               | $B \rightarrow K(J/\psi\omega)$                   | Belle, BABAR       | 2004 | OK     |
| X(3940)                            | $3942^{+9}_{-8}$       | $37^{+27}_{-17}$       | ? <sup>?</sup> +                 | $e^+e^- \rightarrow e^+e^-(J/\psi\omega)$         | Belle, BABAR       |      |        |
| Y(4008)                            | $4008^{+121}_{-49}$    | $226 \pm 97$           | 1 <sup>--</sup>                  | $e^+e^- \rightarrow J/\psi(D\bar{D}^*)$           | Belle              | 2007 | N.Y.C. |
| Z <sub>1</sub> (4050) <sup>+</sup> | $4051^{+24}_{-43}$     | $82^{+51}_{-55}$       | ?                                | $e^+e^- \rightarrow \gamma J/\psi\pi^+\pi^-$      | Belle              | 2007 | N.C.   |
| Y(4140)                            | $4143 \pm 3.0$         | $15^{+11}_{-7}$        | ? <sup>?</sup> +                 | $B \rightarrow K(\pi^+\chi_{c1}(1P))$             | Belle              | 2008 | N.C.   |
| X(4160)                            | $4156^{+29}_{-25}$     | $139^{+113}_{-65}$     | ? <sup>?</sup> +                 | $B \rightarrow K(\phi J/\psi)$                    | CDF                | 2009 | N.C.   |
| Z <sub>2</sub> (4250) <sup>+</sup> | $4248^{+185}_{-45}$    | $177^{+321}_{-72}$     | ?                                | $e^+e^- \rightarrow J/\psi(D\bar{D}^*)$           | Belle              | 2007 | N.Y.C. |
| Y(4260)                            | $4263 \pm 5$           | $108 \pm 14$           | 1 <sup>--</sup>                  | $B \rightarrow K(\pi^+\chi_{c1}(1P))$             | Belle              | 2008 | N.C.   |
|                                    |                        |                        |                                  | $e^+e^- \rightarrow \gamma J/\psi\pi^+\pi^-$      | BABAR, CLEO, Belle | 2005 | OK     |
|                                    |                        |                        |                                  | $e^+e^- \rightarrow J/\psi\pi^0\pi^0$             | CLEO               |      |        |
| Y(4274)                            | $4274.4^{+8.4}_{-6.7}$ | $32^{+22}_{-15}$       | ? <sup>?</sup> +                 | $B \rightarrow K(\phi J/\psi)$                    | CDF                | 2010 | N.C.   |
| X(4350)                            | $4350.6^{+4.6}_{-5.1}$ | $13.3^{+18.4}_{-10.0}$ | 0, 2 <sup>++</sup>               | $e^+e^- \rightarrow e^+e^-(\phi J/\psi)$          | Belle              | 2009 | N.Y.C. |
| Y(4360)                            | $4353 \pm 11$          | $96 \pm 42$            | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma\psi(2S)\pi^+\pi^-$     | BABAR, Belle       | 2007 | OK     |
| Z(4430) <sup>+</sup>               | $4443^{+24}_{-18}$     | $107^{+113}_{-71}$     | ?                                | $B \rightarrow K(\pi^+\psi(2S))$                  | Belle              | 2007 | N.C.   |
| X(4360)                            | $4634 \pm 12$          | $92^{+41}_{-32}$       | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma\Lambda_e^+\Lambda_e^-$ | Belle              | 2007 | N.Y.C. |
| Y(4660)                            | $4664 \pm 12$          | $48 \pm 15$            | 1 <sup>--</sup>                  | $e^+e^- \rightarrow \gamma\psi(2S)\pi^+\pi^-$     | Belle, BABAR       | 2007 | OK     |

Over 16 new resonances only 5 have been seen by different experiments




## ISR $\pi^+\pi^-\psi(2S)$

Y(4360)



Y(4660)



Statistics too low to draw conclusions